Candidate forename			Candidate surname			
Centre number			Candidate number			

OXFORD CAMBRIDGE AND RSA EXAMINATIONS ADVANCED GCE

G628 APPLIED SCIENCE

Sampling, Testing and Processing

WEDNESDAY 15 JUNE 2011: Afternoon DURATION: 1 hour 30 minutes

SUITABLE FOR VISUALLY IMPAIRED CANDIDATES

Candidates answer on the question paper.

OCR SUPPLIED MATERIALS:

Insert (inserted)

OTHER MATERIALS REQUIRED:

Electronic calculator Ruler (cm/mm)


READ INSTRUCTIONS OVERLEAF

INSTRUCTIONS TO CANDIDATES

- Write your name, centre number and candidate number in the boxes on the first page. Please write clearly and in capital letters.
- Use black ink. Pencil may be used for graphs and diagrams only.
- Read each question carefully. Make sure you know what you have to do before starting your answer.
- Write your answer to each question in the space provided. Additional paper may be used if necessary but you must clearly show your candidate number, centre number and question number(s).
- Answer <u>ALL</u> the questions.

INFORMATION FOR CANDIDATES

- Candidates may not bring the Pre-released Case Study into the examination room.
- The number of marks is given in brackets [] at the end of each question or part question.
- The total number of marks for this paper is <u>90</u>.



This means, for example, you should:

- ensure that text is legible and that spelling, punctuation and grammar are accurate so that the meaning is clear;
- organise information clearly and coherently, using specialist vocabulary when appropriate.
- A calculator may be used for this paper.
- You are advised to show all the steps in any calculations.

Answer ALL the questions.

Questions 1 and 2 refer to the materials supplied to your Centre in the Pre-release Case Study.
You are supplied with fresh copies in the Insert.

This question is based on the article 'Apples'.

•	(a)		rieties into the United King	
		1		
		2		
				 [2]

(b)	Apples are susceptible to a number of diseases. Treatments use both <i>contact</i> and <i>systemic</i> products to eradicate diseases. State what is meant by each of these terms and explain a problem that occurs with each type of product, apart from a build up in the soil.
	contact
	systemic

(c)	col	nnah helps a farmer spray apple trees with a oper based fungicide. A fungicide is a material t destroys fungi.
	(i)	State what she must do before she starts this work.
		[1]
	(ii)	Some of the fungicide falls onto the soil around the trees and Hannah collects some soil samples in order to find its copper content. Why should she collect representative
		samples?
	(iii)	Hannah collected more soil samples a week later. Suggest and explain how the concentration of fungicide may be different from the earlier
		samples.
		[2]

(d)	A farmer wishes to find the best conditions for storing a new variety of apple. Using your knowledge of planning experiments, state why it is important to change only one variable when finding the 'best' storage conditions for new varieties.
	[17]

(e) Gas-liquid chromatography was used to identify some of the compounds present in an apple that cause its aroma. The chromatogram, Fig. 1.1, was obtained.




Fig. 1.1

For similar compounds the retention time is often related to the relative molecular mass (Mr). The larger the relative molecular mass, the longer the retention time. In Fig. 1.1 compound A has the shortest retention time and compound G the longest retention time. Table 1.1 gives information about some of the compounds present.

Table 1.1

NAME	M _r
methyl propanoate	88
methyl butanoate	102
ethyl butanoate	116
butyl ethanoate	116
ethyl hexanoate	144

Peak G is produced by ethyl hexanoate.

(i)	compounds w Suggest how t chromatogram	Peak D is a split peak caused by to compounds with similar retention Suggest how the conditions used chromatogram could be changed two peaks become separated.	n times. I to obtain the		
			[1]		
(ii)	State how man	ny compounds are sl n, Fig 1.1.	hown in the		
			[1]		
(iii)	State the name responsible fo	es of the TWO comports of	ounds		
	1	2	[1]		
(iv)	Use Table 1.1 to explain why either peaks B or C could be given by methyl butanoate.				
			[2]		

(f) A student retrieved some apples from cold storage and tested them to see how the percentage of sugar in them changed over a period of time. He used a refractometer to measure the refractive index of the apple juice. The refractive index of a sugar solution depends on the concentration of sugar present.

This is an outline of his method.

- clean and slice an apple and place pieces into a food processor
- collect and filter the juice
- place three drops of the juice into a refractometer and measure the refractive index
- record the results and use a data book to find the concentration of sugar
- repeat the method every two days with a different apple
- plot a graph of the results

(i)	State how the apples to be tested should be stored before use after removal from cold storage.	
		1

(11)	containing the juice?	
	1	
	2	[2]
(iii)	A knife was used to cut the first apple. Why was it necessary to clean the knife before it was used again?	ore
		[1]
(iv)	The results were plotted in a graph, Fig. 1.2 (opposite).	
	Explain why this is a poor graph and what could be done to improve it.	
		[2]
me	e method described in part (f) is an invasive ethod. ate what is meant by an <i>invasive method</i> .	
		 Г1

% of sugar




Fig. 1.2

(h) Unfortunately the results obtained in part (f) were not accurate as all the sugars were assumed to be sucrose. The juices also contain glucose and fructose. However, scientists have recently developed a method of analysing apples for their individual sugar content by use of infrared absorption spectroscopy. In this new method, infrared energy is shone at an apple and the energy not absorbed is collected and analysed. Some results of the analysis are shown for three apples in Table 1.2.

Table 1.2

	PERCENTAGE (%)			
SUGAR	APPLE 1	APPLE 2	APPLE 3	
sucrose	3.2	2.4	3.1	
glucose	3.0	2.2	3.4	
fructose	5.4	5.0	5.6	
TOTAL SUGARS	11.6	9.6	12.1	

(i) Each apple had three measurements taken, equally spaced around the apple. Suggest why this was done.

[1]

	mean percentage =	% [1
ii)	Suggest a reason, other than errors in measurement or calculation, for the lower percentage of sugars in APPLE 2.	
	·	 [1
		_ L
v)	This method uses infrared absorption spectroscopy. Briefly explain how this met works. A description of the apparatus is no required.	hod
v)	spectroscopy. Briefly explain how this met works. A description of the apparatus is no	hod
v)	spectroscopy. Briefly explain how this met works. A description of the apparatus is no	hoc

(ii) Find the mean percentage of the total sugars for the three apples.

(i) When apples are sliced, the enzymes contained in the plant tissues are released. In the presence of oxygen (in air) an enzyme catalyses the conversion of colourless phenolic compounds in the plants to brown pigments called melanins.

This browning process can be slowed by the addition of a solution of Vitamin C to the slices of apple. Oxygen then reacts with the Vitamin C present until all the Vitamin C is used up, when browning starts to occur.



Devise an experiment that will demonstrate the relative rate at which this browning occurs and the effect of Vitamin C on the process.

Your method should be in the form of a set of instructions for other students to follow The

[10	
[TOTAL: 39	

This question is based on the article 'Mercury – a non-biological element?'

2	(a)	pol pla del als Sa	ientists have investigated environmental llution by mercury from the Luoxi processing int in Wuchuan province, China. Mercury was posited on the ground from the atmosphere to from ground water. Imples of soil were collected at various stances from the processing plant.	S
		(i)	State ONE safety precaution that should be taken when collecting these samples, givin reason for your answer.	
		(ii)	Suggest how the soil samples should be stored to avoid cross-contamination.	[2]
				[1]

(iii) State FOUR details that should be written on the labels of each sample container.

[2]

l. _____

(iv) Soil samples at one site were taken from the surface and then at various depths.

The mercury concentration of each sample was plotted in the graph, Fig. 2.1.

concentration of mercury / μg g⁻¹




Fig. 2.1

From the graph the scientists concluded that the mercury compounds present must be water soluble rather than insoluble.

	graph. Fig. 2.1. shows a final co	ncentra
of 0.	graph, Fig. 2.1, shows a final cor 5 μg g ⁻¹ at a depth of 50 cm.	
of 0. The also	5 μg g ⁻¹ at a depth of 50 cm. scientists said that this concent the background level in soils av	tration
of 0. The also the	5 μg g ⁻¹ at a depth of 50 cm. scientists said that this concent	tration way fro

- (b) The soil samples from the mercury processing plant were prepared for analysis by
 - drying in air
 - grinding
 - passing the particles through a fine sieve.

(1)	before they were analysed, state and explain any difference that this might make to the percentage of mercury measured.			
		[2]		
(ii)	Explain why the samples were ground and then passed through a sieve.			
		[1]		

(iii) Atomic Absorption Spectroscopy (AAS) is used to find the concentration of mercury in a solution.

This can be found using the following equation.

$$\frac{[Hg]_1}{[Hg]_2} = \frac{A_1}{A_2}$$

In an experiment

A₁ = absorption of unknown solution = 0.270

A₂ = absorption of known solution = 0.360

[Hg]₁ = concentration of mercury in the unknown solution

 $[Hg]_2$ = concentration of mercury in the known solution = $8.00 \,\mu g \,dm^{-3}$

Use these values to calculate the concentration of mercury in the unknown solution.

_____ μ**g dm⁻³ [2]**

(iv)	The original soil extract solution had a volume of 1 dm ³ . This was too concentrated for AAS and was diluted 100 times with water before use in part (b)(iii). Using your answer from (b)(iii), calculate the concentration of mercury in the original soil extract solution.
(v)	μg dm ⁻³ [1]. The original soil sample solution was prepared from 5.00 g of dried soil. Use your answer from (b)(iv) to calculate the percentage of mercury in the soil sample.

percentage of mercury = ______% [2]

	(VI)	Suggest two reasons why AAS was the choser method.
		1
		2
		[2]
(c	coi So	analysis of surface water for insoluble solids ntaining mercury was also carried out. me of this water was filtered and the solid tained washed and dried.
	(i)	State why the filtered solid was washed with pure water.
		[1]
	(ii)	After drying the filtered solid for 24 hours, it was weighed and then reheated for a further 48 hours and then weighed again. Why was this second drying and reweighing carried out?
		[1]

	(iii)	The following results were obtained.	
mass	s of f	ilter paper and dried solid = 3.9642 g	
		mass of filter paper = 3.9597 g	
		∴ mass of dried solid =	_g
		Complete the results, showing the mass of dried solid.	[1]
	(iv)	The dried solid is described as the total suspended solid (TSS). Analysis of this TSS for mercury showed that the solid weighed in (c)(iii) contained 338 ng $(3.38 \times 10^{-7} \text{g})$ of mercury.	ıt
		Calculate the percentage of mercury in the TSS.	
		percentage of mercury = %	[2]

BLANK PAGE

TURN OVER FOR QUESTION 2 (d)

(d) When a detonator containing mercury fulminate explodes it produces a shockwave that then sets off a secondary explosive.
 A graph of the shockwave velocity against the density of the packed primary explosive is shown in Fig. 2.2.

shockwave velocity /km s⁻¹




Fig. 2.2

(i)	Use the graph to find the shockwave velocity when the density is 3.0 g cm ⁻³		
	shockwave velocity =	km s ⁻¹ [1]	
(ii)	Extend the graph to find the shockwave velocity when the density is 4.4 g cm ⁻³ , explaining the assumption that you have made in finding this answer.		
	shockwave velocity = km s ⁻¹		
	assumption		
		[2]	

- (e) Nowadays, less mercury is being used in laboratories because of its toxicity. It was used in barometers to measure atmospheric pressure.
 - (i) A simple form of barometer consists of a closed tube containing a column of mercury that is supported by atmospheric pressure.




Fig. 2.3

Atmospheric pressure, *P*, is given by the equation

$$P = h \rho g$$

where ρ = density of mercury, h = height of mercury supported and g = a constant.

change (if at all) when atmospheric pressure increases.		
		[;

(ii) Some students found that any liquid can be used to measure atmospheric pressure in this way. The height of liquid supported, h (in cm), depends on the density of the liquid, ρ (in g cm⁻³), and can be found by using the equation shown below, when the pressure is 1 atmosphere.

$$h = \frac{1034}{\rho}$$

Calculate the height of the column when silicone oil is used. Silicone oil has a density of 1.050 g cm⁻³.

$$h =$$
 cm [1]

- (f) Nessler's solution, developed in 1856, is still used as a quantitative test for dissolved ammonia. An outline of the preparation of Nessler's solution is given below.
 - Dissolve 100 g of mercury(II) iodide and 70 g of potassium iodide in 100 cm³ of distilled water.
 - Add this slowly, with stirring, to a cold solution of sodium hydroxide in 700 cm³ of distilled water.
 - Make the solution up to 1 dm³ using distilled water.
 - Leave it to stand and filter if necessary.

(i)	Although Nessler's solution is very toxic, suggest a reason why this is still used.			
		[1]		
(ii)	Explain why any water to be added must be ammonia free.			
		 [1]		

(iii)	The total volume of solution needs to be exactly 1 dm ³ . State why a volumetric flask is used for this purpose.
	[1]
(iv)	The solution has been made up to 1 dm ³ and a stopper placed in the flask. State and explain what should be done next before leaving the solution to stand.
	[2]
	[TOTAL: 34]

- 3 (a) Some farming families in rural America continue to make 'mountain molasses' from sugar cane. 'Mountain molasses' is a sweet dark coloured syrup that is used in place of refined sugar.
 - (i) An outline of this process is given below.
 - Sugar cane stalks are crushed to produce juice.
 - The juice is filtered and boiled for several hours.
 - Solid impurities float to the surface and are removed.
 - After cooling, the thick syrupy molasses is bottled.



Use the outline above to devise a more detailed method for the farming families to make 'mountain molasses' starting with sugar cane stalks.

You should use 20 dm³ of juice in your process, which will produce 2 dm³ of 'mountain molasses'. Your method should mention the size and type of the equipment used and any risks.

Hint: a bucket contains approximately of liquid.)	10 dm ³	

	[7]
(ii)	The sugar content of the prepared molasses was found to be 60%. However, molasses with
	a sugar content of 75% was needed.
	Suggest how the molasses could be treated to increase the percentage of sugar.
	increase the percentage of sugar.
	[1]
	L

- (b) In some countries, molasses and other sugarcontaining mixtures are fermented to produce alcohol, which can be used as a fuel. For economic fermentation the starting sugar solution should contain about 25% sugar.
 - (i) State how much water should be added to $2 \, dm^3$ of molasses, which contains 75% sugar, to make a solution for fermentation that contains 25% of sugar.

water added =	dm ³ [1
mator addod —	MIII I	

(ii) The solution that contained 25% sugar was then fermented and the products fractionally distilled to obtain alcohol.

The apparatus shown in Fig. 3.1, was used to obtain some alcohol.




Fig. 3.1

State the function of the pieces of equipment labelled A and B.

Α			
В			[2

(c) Huge quantities of sugar are used in the sweet industry. Apart from cost and health and safety concerns, state two OTHER factors that should be considered when a colouring agent is to be used in a sweet.

2	 	 	

(d) Food scientists analysed a sweet to see which colouring agents had been used.

They used thin layer chromatography and obtained a chromatogram, Fig. 3.2, for the food colours present.




Fig. 3.2

The R_f values for some food colours are shown in Table 3.1

Table 3.1

FOOD COLOUR	R _f VALUE
A	0.40
В	0.47
С	0.48
D	0.60

(i) Use the chromatogram, Fig. 3.2, to decide which one of these food colours is definitely present.

Explain your reasoning.

food colour = ______
explanation ______

 (ii) Two of the food colours in Table 3.1 have very similar R_f values.
 Suggest what should be done so that clear identification of these food colours is possible.

(iii) One of the spots is removed from the chromatogram and the food colouring extracted.

The mass spectrum of this food colour is then obtained.

What information does the molecular ion value in the mass spectrum tell you about this food colour?

______ [1]

[TOTAL: 17]

END OF QUESTION PAPER



Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.