

GCE 2004
June Series



Mark Scheme

Biology/Human Biology A *BYA6*

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation meeting attended by all examiners and is the scheme which was used by them in this examination. The standardisation meeting ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for the standardisation meeting each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed at the meeting and legislated for. If, after this meeting, examiners encounter unusual answers which have not been discussed at the meeting they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available from:

Publications Department, Aldon House, 39, Heald Grove, Rusholme, Manchester, M14 4NA
Tel: 0161 953 1170

or

download from the AQA website: www.aqa.org.uk

Copyright © AQA 2004 and its licensors

COPYRIGHT

AQA retains the copyright on all its publications. However, registered centres for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to centres to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

The Assessment and Qualifications Alliance (AQA) is a company limited by guarantee registered in England and Wales 3644723 and a registered charity number 1073334. Registered address AQA, Devas Street, Manchester M15 6EX. *Dr Michael Cresswell Director General.*

BYA6**Question 1**

- (a) Suitable accepted evidence, 1 mark for evidence and 1 mark for explanation –
EITHER e.g. guttation
(only) upward pressure could force liquid water out of leaves;
OR
Sap exuding from a cut, rooted stem;
(only) upward force could make this happen; 2
- (b) (*Note: max. two for any component*)
- (i) Evaporation from leaves during daytime only/mainly;
tension/negative pressure (on water) in xylem creates inward pull
(on walls of xylem vessel);
xylem vessels become narrower;
due to adhesion of water molecules (to walls of xylem vessels);
- (ii) root pressure gives outward force/push on walls of xylem vessels;
tree would become wider/stay same diameter;
xylem vessels become wider/stay same diameter; max 3

Total 5 marks

Question 2

- (a) (i) **Many** gill lamellae/gill filaments; (ignore refs to ‘highly divided’) 1
- (ii) Counter-current mechanism/blood and water flow in opposite directions;
Not enough time for equalisation of concentrations/maintains concentration
gradient over length of gills/never reaches equilibrium; max 2
- (b) Humidity reduces difference in concentration of water (vapour) between body and air;
reduces rate of diffusion (of water vapour)(as are proportional); 2

Total 5 marks

Question 3

- (a) (More) urea formed from surplus protein/eq./surplus protein deaminated; 1
- (b) (i) In group A, water lost in urine replaced by drinking (water)/converse for group B; 1
- (ii) *For group B:*
 More concentrated filtrate;
 Decreased reabsorption from collecting duct;
 More water lost in urine;
 Less respiratory water formed by group B;
 (Allow reverse arguments for all points for group C) max 2
- (c) Water withdrawn by osmosis/down water potential gradient; 1

Total 5 marks

Question 4

- (a) Structure resulting from aggregation of several polypeptide chains/tertiary structures/eq: 1
- (b) Low pH/(more)H⁺; due to (increased) CO₂ (increased) respiration;
 (ignore refs to buffering action of haemoglobin)
 (increased) dissociation of haemoglobin;
 Low oxygen tension in tissues/plasma;
 Oxygen diffuses from r.b.c. to tissues; max 3
- (c) Deaminated for use in respiration/used in protein synthesis/suitable e.g.; 1

Total 5 marks

Question 5

- (a) Compact molecule so can store much in one cell;
 Insoluble so will not affect water potential/will not 'move' from storage site
 Branched molecule so readily hydrolysed max 2
- (b) Adult needs energy for movement/flight/mating;
 Sugars easily respired
 No growth in this phase so no need for protein;
 Sufficient protein acquired during larval stage (for egg production); max 3

Total 5 marks

Question 6

- (a) Flattened squashed/deformation of lamellae/membranes/corpuscle;
Causes sodium ion channels in axon membrane to open/increases permeability of axon membrane/nerve ending/nerve fibres;
Movement of sodium ions into axon/nerve ending/nerve fibre
[If ions mentioned once, assume candidate is referring to ions throughout; if no mention of ions penalise once only]
- max 2
- (b) Correct ref to near vision involved;
tension in suspensory ligaments;
lens does not become sufficiently convex
rays of light refracted less/bent less;
(focusing near objects) requires maximum refraction/maximum bending;
or rays of light focused beyond retina;
Allow converse for a 'young person' if this is made clear.
- max 3
- Total 5 marks

Question 7

- (a) (i) Exopeptidase;
amino acids are being broken off from the end/one by one; 2
- (ii) Hydolysis; ® 'wrong' bond
(e.g. glycosidic)
water (molecules) used;
to break peptide bonds/ to break bonds between amino acids;
(Accept last two marks from a suitable diagram if one is given) 3
- (b) Small intestine/duodenum/ileum;
enzyme most active in neutral/alkaline conditions/between pH 7-8;
found in this region of gut 3
- (c) (i) Organs comprise several tissues;
capillary comprises only one tissue/endothelial/epithelial tissue; 2
- (ii) A – facilitated diffusion
as transport protein needed but ATP not needed; In all cases reject
'energy' unless
qualified
B – active transport
as (transport protein and) ATP needed;
C – (simple) diffusion
as neither ATP nor transport protein needed;
(Ignore all references to concentration gradients) 3
- (iii) creates low concentration of amino acids/Na⁺ in cell
concentration gradient established between lumen and cell
(of amino acids or Na⁺) 2
- Total 15 marks

Question 8

- (a) (i) Increase of breathing/increased rate of contraction of intercostals / diaphragm;
increases tidal volume/depth of breathing/stronger contractions of
intercostals and/or diaphragm; 2
- (ii) More oxygen/**oxygenated** blood to cardiac muscle;
more glucose to cardiac muscle;
allows increased **aerobic** respiration;
increased heart rate
increased stroke volume max 4
- (iii) Sympathetic increases (heart rate) parasympathetic decreases (heart rate) 1
- (b) (i) Binds to receptor/proteins; and opens Na^+ channels;
 Na^+ enter and make membrane potential less negative/depolarised 2
- (ii) (Vesicles containing) neurotransmitter only in presynaptic membrane/neurone;
receptor/proteins only in postsynaptic membrane/neurone;
so neurotransmitter diffuses down concentration gradient; max 2
- (iii) GABA opens K^+ and Cl^- channels;
 K^+ passes out and Cl^- passes in;
Membrane potential more negative/hyperpolarised;
Requires increased stimulation/must open more Na^+ channels/allow more Na^+ to enter;
To reach threshold; Max 4

Total 15 marks

Question 9**Quality of Communication**

The answers to all sections of this question require the use of continuous prose. Quality of language should be considered in crediting points in the scheme. In order to gain credit, answers should be expressed logically and unambiguously, using scientific terminology where appropriate.

- (a)
1. Deviation of a value from norm initiates corrective mechanisms;
 2. fluctuations in plasma glucose concentration detected by hypothalamus/islet cells in pancreas;
 3. initial decrease, no food given (in plasma glucose) stimulates (increased) secretion of glucagon;
 4. increases (in plasma glucose) stimulate (increased) secretion of insulin;
 5. correct ref. to role of α and/or β cells as secretors;
 6. correct ref. to interconversion of glycogen / glucose;
 7. increased/decreased uptake of glucose by cells (as appropriate)/correct ref to change in membrane permeability;

max 5

- (b)(i)
1. Sensors in skin/hypothalamus detect reduced temperature;
 2. heat gain centre activated/inhibition of heat loss centre;
 3. vasoconstriction/constriction of arterioles in skin surface; (**R** capillaries)
 4. dilation of shunt vessels/constriction of – capillary sphincter;
 5. less blood to skin surface/capillaries
 6. reduced heat loss by radiation;
 7. increased heat gain by increased metabolic rate/respiration/movement/shivering;
 8. decreased heat loss by putting on clothes/huddling/reduced sweating;

max 5

- (ii)
1. Body temp./ 37°C is optimum temp for enzymes;
 2. excess heat denatures enzymes/alters tertiary structure/alters shape of active site/enzyme;
 3. substrate cannot bind/eq.;
 4. reactions cease/slowed;
 5. too little reduces kinetic energy of molecules / molecules move more slowly;
 6. fewer collisions/fewer ES complexes formed?

max 5

Total 15 marks