
Surname			Other	Names			
Centre Number				Candida	ate Number		
Candidate Signa	ture						



General Certificate of Education June 2003 Advanced Subsidiary Examination



# CHEMISTRY CHM1 Unit 1 Atomic Structure, Bonding and Periodicity

Wednesday 4 June 2003 Morning Session

In addition to this paper you will require:	
a calculator.	

Time allowed: 1 hour

### Instructions

- Use blue or black ink or ball-point pen.
- Fill in the boxes at the top of this page.
- Answer **all** questions in **Section A** and **Section B** in the spaces provided. All working must be shown.
- Do all rough work in this book. Cross through any work you do not want marked.
- The Periodic Table/Data Sheet is provided on pages 3 and 4. Detach this perforated sheet at the start of the examination.

### **Information**

- The maximum mark for this paper is 60.
- Mark allocations are shown in brackets.
- This paper carries 30 per cent of the total marks for AS. For Advanced Level this paper carries 15 per cent of the total marks.
- You are expected to use a calculator where appropriate.
- The following data may be required. Gas constant  $R = 8.31 \,\mathrm{J \, K^{-1} \, mol^{-1}}$
- Your answers to the question in **Section B** should be written in continuous prose, where appropriate. You will be assessed on your ability to use an appropriate form and style of writing, to organise relevant information clearly and coherently, and to use specialist vocabulary, where appropriate.

### **Advice**

• You are advised to spend about 45 minutes on **Section A** and about 15 minutes on **Section B**.

	For Exam	iner's Use	
Number	Mark	Number	Mark
1			
2			
3			
4			
5			
Total (Column	1)	<b>→</b>	
Total (Column	2)	$\rightarrow$	
TOTAL			
Examine	r's Initials		

Copyright © 2003 AQA and its licensors. All rights reserved.

# The Periodic Table of the Elements

The atomic numbers and approximate relative atomic masses shown in the table are for use in the examination unless stated otherwise in an individual question.

20.2 **Ne** Krypton 36 **2**22.0 **Rn** Radon Helium Helium Argon Xenon Bromine 35 Fluorine Chlorine Astatine lodine **=** 32.5 C 126.9 79.0 **Se** Selenium 34 Polonium 84 Phosphorus Sulphur Tellurium 16.0 O Oxygen 32.1 **S** 5 210.0 **Po** Antimony 51 74.9 **As** As Arsenic 33 Nitrogen Bismuth 83 121.8 **Sb** > Germanium 32 **Sn** Carbon <u>28.1</u> 72.6 **Ge** Silicon ≥ 20 Aluminium Thallium 81 Gallium **Ga** Boron Indium 114.8 **n** ≡ 27.0 **A** Cadmium 48 **55.4 Zn** Zinc 200.6 **Hg** Mercury 80 112.4 **Cd** 30 **Ag** Silver Copper **Au** Gold 47 Palladium Platinum 78 **S8**.7 Nickel 195.1 106.4 **Pd** 46 58.9 Cobalt Rhodium Iridium 102.9 **Rh** 55.8 **Fe** Iron Ruthenium Osmium 190.2 **Os** Chromium Manganese 24 25 Molybdenum Technetium 42 Rhenium 75 Lithium ≔ Tungsten 74 183.9 **W** relative atomic mass Vanadium 23 Tantalum | 92.9 | **Nb** Niobium 180.9 **Ta** atomic number Zirconium 40 Titanium | Lanthanum | Hafnium | 57 \* 178.5 **#** Key Scandium Yttrium Actinium 138.9 **La 8**8.9 33 Calcium Strontium Magnesium Beryllium **Qa** Barinm Radium 88 Potassium 19 Caesium Hydrogen Rubidium Francium Sodium Lithium 39.1 **X** 132.9 **Cs** I 1.0

**************************************	140.1 <b>Ce</b>	Ce Pr 144.2 144.2 Ce Pr Nd Pr	144.2 <b>Nd</b>	ි <b>E</b>	150.4 <b>Sm</b>	152.0 <b>Eu</b>	157.3 <b>Gd</b>	158.9 <b>Tb</b>	162.5 <b>Dy</b>	6.4.9 <b>5.</b> 9	167.3 <b>Er</b>		173.0 <b>Yb</b>	175.0 <b>Lu</b>
. <b>38 – 71</b> Lanmanides	Cerium 58	Cerium         Praseodymium         Neodymium         Prome           58         59         60         61	Neodymium 60	thiur.	n Samarium 62	_	Gadolinium 54	Terbium 35	m Terbium Dysprosium H	Holmium 7		Thulium 69	Ytterbium 70	Lutetium 71
007	232.0 <b>Th</b>	32.0 231.0 238.0 237.0 <b>Th Pa U N</b>	238.0 <b>U</b>		239.1 <b>Pu</b>	243.1 <b>Am</b>	247.1 <b>Cm</b>	247.1 <b>BK</b>	252.1 <b>Cf</b>	52) <b>ES</b>	(257) <b>Fm</b>		(259) <b>No</b>	(260) <b>Lr</b>
7 <b>90 - 103</b> Actinides	Thorium 90	Thorium Protactinium Uranium 92	Uranium 92	Neptunium 93	_	Americium 95	Curium 96	Berkelium 97	Californium 98	nsteinium 3				Lawrencium 103

**Table 1** Proton n.m.r chemical shift data

Type of proton	δ/ppm
$RCH_3$	0.7–1.2
$R_2CH_2$	1.2–1.4
$R_3$ CH	1.4–1.6
$RCOCH_3$	2.1–2.6
$ROCH_3$	3.1–3.9
$RCOOCH_3$	3.7–4.1
ROH	0.5-5.0

**Table 2** Infra-red absorption data

Bond	Wavenumber/cm <sup>-1</sup>
С—Н	2850–3300
C—C	750–1100
C=C	1620–1680
C=O	1680–1750
С—О	1000-1300
O—H (alcohols)	3230–3550
O—H (acids)	2500–3000

# SECTION A

Answer all questions in the spaces provided.

1	(a)	(i)	Complete the electronic configuration of aluminium.  1s <sup>2</sup>	
		(ii)	State the block in the Periodic Table to which aluminium belongs.	
				(2 marks)
	(b)	Desc	cribe the bonding in metals.	
				(2 marks)
	(c)	Expl	ain why the melting point of magnesium is higher than that of sodium.	
				(3 marks)
	(d)	Expl	lain how metals conduct electricity.	
		•••••		
		•••••		
		•••••		(2 marks)



2	(a)	Give	the relative charge and relative mass of an electron.
		Rela	tive charge
		Rela	tive mass
			(2 marks)
	(b)	Isoto	opes of chromium include <sup>54</sup> Cr and <sup>52</sup> Cr
		(i)	Give the number of protons present in an atom of <sup>54</sup> Cr
		(ii)	Deduce the number of neutrons present in an atom of <sup>52</sup> Cr
		(iii)	Apart from the relative mass of each isotope, what else would need to be known for the relative atomic mass of chromium to be calculated?
			(3 marks)
	(c)		der to obtain a mass spectrum of a gaseous sample of chromium, the sample must be ionised.
		(i)	Give <b>two</b> reasons why it is necessary to ionise the chromium atoms in the sample.
			Reason 1
			Reason 2
		(ii)	State what is adjusted so that each of the isotopes of chromium can be detected in turn.
		(iii)	Explain how the adjustment given in part (c)(ii) enables the isotopes of chromium to be separated.
			(4 marks)

(d)	(i)	State what is meant by the term empirical formula.
	(ii)	A chromium compound contains 28.4% of sodium and 32.1% of chromium by mass, the remainder being oxygen.  Calculate the empirical formula of this compound.
		(4 marks)



# TURN OVER FOR THE NEXT QUESTION

<b>3</b> (a)		ample of ethanol vapour, $C_2H_5OH$ ( $M_r = 46.0$ ), was maintained at a pressure of $KPa$ and at a temperature of $366  \text{K}$ .
	(i)	State the ideal gas equation.
	(ii)	Use the ideal gas equation to calculate the volume, in cm <sup>3</sup> , that 1.36 g of ethanol vapour would occupy under these conditions. (The gas constant $R = 8.31 \mathrm{J  K^{-1}  mol^{-1}}$ )
		(5 marks)
(1.)	2.6	
(b)		nesium nitride reacts with water to form magnesium hydroxide and ammonia.
	(i)	Balance the equation, given below, for the reaction between magnesium nitride and water.
		$Mg_3N_2 + H_2O \rightarrow Mg(OH)_2 + NH_3$
	(ii)	Calculate the number of moles, and hence the number of molecules, of NH <sub>3</sub> in 0.263 g of ammonia gas. (The Avogadro constant $L=6.02\times10^{23}~\rm mol^{-1}$ )
		(4 marks)

(c) Sodium carbonate is manufactured in a two-stage process as shown by the equations below.

NaCl + NH<sub>3</sub> + CO<sub>2</sub> + H<sub>2</sub>O 
$$\rightarrow$$
 NaHCO<sub>3</sub> + NH<sub>4</sub>Cl 
$$2NaHCO_3 \rightarrow Na_2CO_3 + H_2O + CO_2$$

Calculate the maximum mass of sodium carbonate which could be obtained from of sodium chloride.	om 800 g
	•••••
	••••••
	••••••
	••••••
(4	1 marks)



TURN OVER FOR THE NEXT QUESTION

4	(a)		HF and HCl are molecules having a polar covalent bond. Their boiling points are K and 188K respectively.
		(i)	State which property of the atoms involved causes a bond to be polar.
		(ii)	Explain, in terms of the intermolecular forces present in each compound, why HF has a higher boiling point than HCl.
			(4 marks)
	(b)		n aluminium chloride reacts with chloride ions, as shown by the equation below, a rdinate bond is formed.
			$AlCl_3 + Cl^- \rightarrow AlCl_4^-$
		Expl	ain how this co-ordinate bond is formed.
		•••••	(2 marks)

(c)	Draw the shape of the PCl <sub>5</sub> molecule and of angles.	the PCl <sub>4</sub> <sup>+</sup> ion. State the value(s) of the bond
	$PCl_5$	$PCl_4^+$
	Bond angle(s)	Bond angle(s)
		(4 marks)



# TURN OVER FOR THE NEXT QUESTION

### **SECTION B**

Answer the question below in the space provided on pages 12 to 16 of this booklet.

5 (a) The table below gives the melting point for each of the Period 3 elements Na – Ar.

Element	Na	Mg	Al	Si	P	S	Cl	Ar
Melting point/K	371	923	933	1680	317	392	172	84

In terms of structure and bonding, explain why silicon has a high melting point, and why the melting point of sulphur is higher than that of phosphorus. (7 marks)

- (b) Draw a diagram to show the structure of sodium chloride. Explain, in terms of bonding, why sodium chloride has a high melting point. (4 marks)
- (c) Give the conditions under which, if at all, beryllium and magnesium react with water. For any reaction that occurs, state **one** observation you would make and write an equation. (4 marks)

## **END OF QUESTIONS**