



Mark Scheme (Results)

Summer 2022

Pearson Edexcel International Advanced
Subsidiary Level
In Chemistry (WCH12)
Paper 01: Energetics, Group Chemistry,
Halogenoalkanes and Alcohols

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2022

Question Paper Log Number P70965A

Publications Code WCH12_01_2206_MS

All the material in this publication is copyright

© Pearson Education Ltd 2022

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Using the mark scheme

Examiners should look for qualities to reward rather than faults to penalise. This does NOT mean giving credit for incorrect or inadequate answers, but it does mean allowing candidates to be rewarded for answers showing correct application of principles and knowledge. Examiners should therefore read carefully and consider every response: even if it is not what is expected it may be worthy of credit.

The mark scheme gives examiners:

- an idea of the types of response expected
- how individual marks are to be awarded
- the total mark for each question
- examples of responses that should NOT receive credit.

/ means that the responses are alternatives and either answer should receive full credit. () means that a phrase/word is not essential for the award of the mark, but helps the examiner to get the sense of the expected answer.

Phrases/words in **bold** indicate that the meaning of the phrase or the actual word is **essential** to the answer. ecf/TE/cq (error carried forward) means that a wrong answer given in an earlier part of a question is used correctly in answer to a later part of the same question.

Candidates must make their meaning clear to the examiner to gain the mark. Make sure that the answer makes sense. Do not give credit for correct words/phrases which are put together in a meaningless manner. Answers must be in the correct context.

| Question number | Answer | Mark |
|-----------------|---|------|
| 1 | <p>The only correct answer is A ($2\text{Al(s)} + 1\frac{1}{2}\text{O}_2\text{(g)} \rightarrow \text{Al}_2\text{O}_3\text{(s)}$)</p> <p><i>B is incorrect because oxygen exists as O₂ in its standard state</i></p> <p><i>C is incorrect because standard enthalpy change of formation is for the formation of 1 mol of a compound</i></p> <p><i>D is incorrect because standard enthalpy change of formation is for the formation of 1 mol of a compound and oxygen exists as O₂ in its standard state</i></p> | (1) |

| Question number | Answer | Mark |
|-----------------|---|------|
| 2 | <p>The only correct answer is A (gains electrons and decreases in oxidation number)</p> <p><i>B is incorrect because oxidising agents are reduced during a reaction so there is a decrease in oxidation number</i></p> <p><i>C is incorrect because oxidising agents are reduced during a reaction so they gain electrons</i></p> <p><i>D is incorrect because oxidising agents are reduced during a reaction so they gain electrons and there is a decrease in oxidation number</i></p> | (1) |

| Question number | Answer | Mark |
|-----------------|---|------|
| 3 | <p>The only correct answer is B ($\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$)</p> <p><i>A is incorrect because branching in the carbon chain reduces the boiling temperature of isomeric alcohols</i></p> <p><i>C is incorrect because alkanes do not have hydrogen bonding and have lower boiling temperatures than alcohols with a similar number of electrons</i></p> <p><i>D is incorrect because alkanes do not have hydrogen bonding and have lower boiling temperatures than alcohols with a similar number of electrons</i></p> | (1) |

| Question number | Answer | Mark |
|-----------------|---|------|
| 4 | <p>The only correct answer is B (potassium chlorate(III))</p> <p><i>A is incorrect because the oxidation number of chlorine in $KClO_2$ is +3</i></p> <p><i>C is incorrect because the oxidation number of chlorine in $KClO_2$ is +3</i></p> <p><i>D is incorrect because the oxidation number of chlorine in $KClO_2$ is +3</i></p> | (1) |
| Question number | Answer | Mark |
| 5(a) | <p>The only correct answer is D ($Ca + 2H_2O \rightarrow Ca(OH)_2 + H_2$)</p> <p><i>A is incorrect because CaO is not a product of the reaction</i></p> <p><i>B is incorrect because CaO_2 is an incorrect formula for calcium oxide</i></p> <p><i>C is incorrect because $CaOH$ is an incorrect formula for calcium hydroxide</i></p> | (1) |
| Question number | Answer | Mark |
| 5(b) | <p>The only correct answer is A (calcium oxidised, hydrogen reduced)</p> <p><i>B is incorrect because oxygen is not reduced</i></p> <p><i>C is incorrect because hydrogen is not oxidised and calcium is not reduced</i></p> <p><i>D is incorrect because hydrogen is not oxidised and oxygen is not reduced</i></p> | (1) |

| Question number | Answer | Mark |
|-----------------|--|------|
| 6 | <p>The only correct answer is A ($\text{BaCO}_3(\text{s}) + 2\text{H}^+(\text{aq}) \rightarrow \text{Ba}^{2+}(\text{aq}) + \text{CO}_2(\text{g}) + \text{H}_2\text{O}(\text{l})$)</p> <p><i>B is incorrect because Ba_2CO_3 is not the formula for barium carbonate</i></p> <p><i>C is incorrect because solid barium carbonate should not be split up into ions</i></p> <p><i>D is incorrect because hydrochloric acid is in solution and should be split up into ions and Cl^- are spectator ions</i></p> | (1) |

| Question number | Answer | Mark |
|-----------------|---|------|
| 7 | <p>The only correct answer is D (Reagent: $\text{NaOH}(\text{aq})$, Test for gas: damp red litmus paper turns blue)</p> <p><i>A is incorrect because hydrochloric acid does not react with ammonium ions</i></p> <p><i>B is incorrect because hydrochloric acid does not react with ammonium ions</i></p> <p><i>C is incorrect because ammonia is produced and it is alkaline so turns damp red litmus paper blue</i></p> | (1) |

| Question number | Answer | Mark |
|-----------------|--|------|
| 8 | <p>The only correct answer is D (violet)</p> <p><i>A is incorrect because iodine is produced in the reaction and it is brown in aqueous solution but violet in a non-polar organic solvent</i></p> <p><i>B is incorrect because chlorine gas is green</i></p> <p><i>C is incorrect because bromine is orange in a non-polar organic solvent</i></p> | (1) |

| Question number | Answer | Mark |
|-----------------|--|------|
| 9(a) | <p>The only correct answer is C (yellow to orange)</p> <p><i>A is incorrect because methyl orange is yellow in alkaline solution</i></p> <p><i>B is incorrect because methyl orange is yellow in alkaline solution and turns red when excess acid has been added</i></p> <p><i>D is incorrect because methyl orange turns red when excess acid is added</i></p> | (1) |

| Question number | Answer | Mark |
|-----------------|--|------|
| 9(b) | <p>The only correct answer is C (22.80, 22.35, 22.40 (cm³))</p> <p><i>A is incorrect because the first titre should be higher than the other two titres</i></p> <p><i>B is incorrect because the second and third titres should be concordant and lower than the first titre</i></p> <p><i>D is incorrect because the second and third titres should be lower than the first titre</i></p> | (1) |

| Question number | Answer | Mark |
|-----------------|--|------|
| 9(c) | <p>The only correct answer is B (0.0668 (mol dm⁻³))</p> <p><i>A is incorrect because the volumes have been used the wrong way round</i></p> <p><i>C is incorrect because the mole ratio of 2 : 1 has not been used</i></p> <p><i>D is incorrect because the mole ratio has been used as 2 H₂SO₄ : 1 NaOH</i></p> | (1) |

| Question number | Answer | Mark |
|-----------------|--|------|
| 10(a) | <p>The only correct answer is D (decreasing the activation energy of the reaction)</p> <p><i>A is incorrect because only an increase in temperature causes the average kinetic energy of the molecules to increase</i></p> <p><i>B is incorrect because only a decrease in temperature causes the average kinetic energy of the molecules to decrease</i></p> <p><i>C is incorrect because if the activation energy increased, the rate of decomposition would decrease</i></p> | (1) |

| Question number | Answer | Mark |
|-----------------|---|------|
| 10(b) | <p>The only correct answer is D (0.833 (mol dm⁻³))</p> <p><i>A is incorrect because the volume of oxygen has not been converted into moles</i></p> <p><i>B is incorrect because the 2 : 1 mole ratio has been used the wrong way around</i></p> <p><i>C is incorrect because the 2 : 1 mole ratio has not been used</i></p> | (1) |

| Question number | Answer | Mark |
|-----------------|--|------|
| 11 | <p>The only correct answer is A (change in equilibrium position: left, enthalpy change: endothermic)</p> <p><i>B is incorrect because an exothermic reaction would decrease the concentration of iodine</i></p> <p><i>C is incorrect because the equilibrium position would shift to the left</i></p> <p><i>D is incorrect because the equilibrium position would shift to the left</i></p> | (1) |

| Question number | Answer | Mark |
|-----------------|--|------|
| 12 | <p>The only correct answer is C (nucleophile)</p> <p><i>A is incorrect because electrophiles attack atoms with a slight negative charge</i></p> <p><i>B is incorrect because free radicals attack neutral atoms</i></p> <p><i>D is incorrect because oxidising agents remove electrons from a species</i></p> | (1) |

| Question number | Answer | Mark |
|-----------------|--|------|
| 13 | <p>The only correct answer is B (P and Q only)</p> <p><i>A is incorrect because Q is also primary alcohol and will be oxidised to a carboxylic acid</i></p> <p><i>C is incorrect because R is a secondary alcohol and will be oxidised to a ketone</i></p> <p><i>D is incorrect because R is a secondary alcohol and will be oxidised to a ketone and S is a tertiary alcohol so is not easily oxidised</i></p> | (1) |

| Question number | Answer | Mark |
|-----------------|---|------|
| 14(a) | <p>The only correct answer is C (concentrated phosphoric(V) acid)</p> <p><i>A is incorrect because acidified potassium manganate(VII) converts an alkene into a diol</i></p> <p><i>B is incorrect because aqueous bromine reacts with an alkene to form a bromoalcohol</i></p> <p><i>D is incorrect because phosphorus(V) chloride reacts with an alcohol to form a chloroalkane</i></p> | (1) |


| Question number | Answer | Mark |
|-----------------|---|------|
| 14(b) | <p>The only correct answer is C ($C_6H_{11}OH + [O] \rightarrow C_6H_{10}O + H_2O$)</p> <p><i>A is incorrect because [O] is needed from the oxidising agent and hydrogen gas would not be produced</i></p> <p><i>B is incorrect because [O] is needed from the oxidising agent and hydrogen atoms would not be produced</i></p> <p><i>D is incorrect because the oxidising agent is not oxygen gas</i></p> | (1) |

| Question number | Answer | Mark |
|-----------------|---|------|
| 14(c) | <p>The only correct answer is C (3750-3200, 1669-1645)</p> <p><i>A is incorrect because there is a C-H bond in both compounds</i></p> <p><i>B is incorrect because there is a C-H bond in both compounds and there is no C=O in cyclohexene</i></p> <p><i>D is incorrect because there is no C=O in cyclohexene</i></p> | (1) |

(Total for Section A = 20 marks)

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|--|--|------------|
| 15(a) | <p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> • Size (and charge) calcium ion / Ca^{2+} has a larger (ionic) radius than a magnesium ion (but the same charge) or magnesium ion / Mg^{2+} has a smaller (ionic) radius / is smaller than a calcium ion (but the same charge) (1) • Polarising power so calcium ion / Ca^{2+} causes less polarisation / distortion or magnesium ion / Mg^{2+} causes more polarisation / distortion (1) • What is polarised of the carbonate ion / CO_3^{2-} / anion / negative ion / C–O bonds / C=O bonds / CO bonds (1) | <p>Penalise omission of ‘ion’ or just magnesium / Mg / calcium / Ca without charge once only in M1 and M2 Allow reverse argument for magnesium ions in M1 and M2</p> <p>Allow size for radius or just ‘bigger / smaller’ Allow ionic radius / size of cation increases down the group / decreases up the group Allow calcium carbonate has a larger cation Allow calcium ions have a lower charge density Allow calcium ions have more shells of electrons</p> <p>Ignore effective nuclear charge / mass : charge ratio Ignore atomic radius Ignore omission of same charge</p> <p>Do not award M1 if mention of different / incorrect charges on magnesium and calcium ions</p> <p>Allow polarising power of cation decreases down the group Allow Ca^{2+} causes less weakening of bonds for polarisation</p> <p>Do not award just ‘the carbonate ion is less polarisable’ for M2, although this can score M3</p> <p>Allow electron cloud for ion Do not award reference to nitrate / N-O bonds Do not award reference to breaking unspecified bonds / (ionic) bond between cation and anion Do not award references to intermolecular forces</p> | (3) |

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|--|--|------|
| 15(b)(i) | <ul style="list-style-type: none"> • calculation of mol of CO₂ (1) • calculation of mol of HCl (1) • calculation of volume of HCl and corresponding volume unit (1) | <p>Example of calculation:</p> <p>mol CO₂ = $\frac{100}{24000} = 0.0041667 / 4.1667 \times 10^{-3}$ (mol)</p> <p>mol HCl = $2 \times 0.0041667 = 0.0083333 / 8.3333 \times 10^{-3}$ (mol) TE on M1</p> <p>vol HCl = $\frac{0.0083333}{0.500} \times 1000 = (16.667)$ = 16.6 cm³ / 0.0166 dm³</p> <p>Do not award incorrect units e.g. cm⁻³ / dm⁻³</p> <p>Allow 16.67 / 16.7 cm³ as the theoretical volume of CO₂ is 100.02 / 100.2 cm³ Do not award 17 cm³ as the theoretical volume of CO₂ is 102 cm³ so would exceed the measurable volume of the syringe</p> <p>Allow any number between 16 and 16.7 cm³ / 0.016 and 0.0167 dm³ inclusive TE on M2 Ignore SF except 1 SF</p> <p>Correct answer with units and no working scores (3)</p> <p>Accept fractions / correct working not evaluated for M1 and M2 e.g. 1/240, 1/120</p> | (3) |

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|--|--|------|
| 15(b)(ii) | <ul style="list-style-type: none"> • tangent drawn at $t = 0$ • gradient • units | <p>Example of working:</p>  <p>Tangent must touch the curve for at least 24 s (2 small squares horizontally) and extend to at least 20 cm³</p> <p>Gradient = $\frac{100}{360} = 0.27778$ (expected value 0.25 to 0.33 for tangent at $t = 0$)</p> <p>TE on tangent drawn at any time value</p> <p>If no tangent drawn, allow a selected point and y/x value e.g. $32/120 = 0.27$</p> <p>Stand alone mark $\text{cm}^3 \text{ s}^{-1}$ or cm^3 / s or $\frac{\text{cm}^3}{\text{s}}$</p> <p>Allow $\text{cm}^3 \text{ s}^{-1}$</p> | (3) |

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|---|--|------|
| 15(b)(iii) | <p>An answer that makes reference to the following point:</p> <ul style="list-style-type: none"> initial rate halves and final volume of CO₂ halves / is 45 (cm³) | <p>Allow initial rate decreases and final volume of CO₂ decreases</p> <p>Do not award any specific decrease (e.g. decrease by a factor of 4) except for half</p> | (1) |

(Total for Question 15 = 10 marks)

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|---|---|------|
| 16(a) | <p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> • atomic radius increases or distance between the nucleus and outer electrons increases or there are more shells / energy levels of inner electrons between the nucleus and the outer shell electrons (1) • so there is less attraction (by the nucleus with a higher charge) for the bonding electrons / shared pair of electrons (1) | <p>Allow size of atoms increases / gets bigger Allow just 'more shells of electrons' Allow effective nuclear charge decreases Do not award nuclear charge decreases Do not award reference to ions / ionic radius for M1 only</p> <p>Allow greater shielding between the nucleus and the bonding electrons / shared pair of electrons Note – bonding / shared pair can be mentioned anywhere in the answer</p> | (2) |







| Question Number | Answer | Additional Guidance | Mark |
|-----------------|---|--|------|
| 16(b) | <p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> three oxidation numbers of I: $\text{IO}_3^- = (+)5$ $\text{I}^- = -1$ $\text{I}_2 = 0$ (1) two different species / ions / compounds (of iodine) are oxidised and reduced (to form the same species) or there is not one species / ion / compound that is being oxidised and reduced or 2 different oxidation states are not produced from one oxidation state (of iodine) or only one species / oxidation state of iodine is formed (1) | <p>Allow oxidation numbers written near species in the equation</p> <p>Ignore oxidation numbers of H and O</p> <p>Do not award O.N. I_2 is neutral / I^- is -5</p> <p>Allow I_2 / iodine is oxidised and reduced in the reverse reaction</p> <p>Allow (iodine in) IO_3^- is only being reduced or (iodine in) I^- is only being oxidised</p> <p>Ignore just 'the reaction is only oxidation / reduction'</p> <p>Ignore just 'comproportionation'</p> <p>Ignore just 'I / iodine is not simultaneously oxidised and reduced'</p> | (2) |

| Question Number | Answer | Additional Guidance | Mark | | | | | | |
|-----------------|--|--|-----------------|--|-------|--------------------------------|------|---|-----|
| 16(c) | <ul style="list-style-type: none"> sulfur dioxide / sulfur (IV) oxide / SO_2 produced from HBr and hydrogen sulfide / H_2S produced from HI | <p>Example of table:</p> <table border="1"> <tr> <td>Hydrogen halide</td> <td>Compound produced with the lowest oxidation number of sulfur</td> </tr> <tr> <td>(HBr)</td> <td>sulfur dioxide / SO_2</td> </tr> <tr> <td>(HI)</td> <td>hydrogen sulfide / H_2S</td> </tr> </table> <p>Ignore Br_2 and I_2</p> <p>Note - If name and formula are given, both must be correct</p> | Hydrogen halide | Compound produced with the lowest oxidation number of sulfur | (HBr) | sulfur dioxide / SO_2 | (HI) | hydrogen sulfide / H_2S | (1) |
| Hydrogen halide | Compound produced with the lowest oxidation number of sulfur | | | | | | | | |
| (HBr) | sulfur dioxide / SO_2 | | | | | | | | |
| (HI) | hydrogen sulfide / H_2S | | | | | | | | |

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|---|---|------------|
| 16(d) | <p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> • all hydrogen halides have London forces (and dipole-dipole forces between molecules) (1) • the strength of the London forces increases as the number of electrons increases (so the boiling temperature increases from HCl to HI) or the strength of the London forces increases as the polarisability of the molecules increases from HCl to HI (1) • (only) HF has hydrogen bonding (between molecules) (1) • hydrogen bonding is (much) stronger than London forces / dipole-dipole forces (so HF has the highest boiling temperature) (1) | <p>Allow van der Waals' forces / dispersion forces / attractions between instantaneous dipoles and induced dipoles for London forces or a description of London forces</p> <p>Ignore London forces omitted from HF Do not award this mark if ions mentioned in answer Do not award this mark if breaking H-Cl, H-Br or H-I bonds</p> <p>Ignore the strength of the London forces increases as the size of the molecule / M_r increases</p> <p>Do not award M3 if hydrogen bonding in any other hydrogen halide</p> <p>Allow more heat energy is needed to overcome hydrogen bonding than London forces Allow hydrogen bonding is the strongest intermolecular force / bond</p> | (4) |


| Question Number | Answer | Additional Guidance | Mark |
|-----------------|---|---|------|
| 16(e) | <p>An answer that makes reference to the following point:</p> <ul style="list-style-type: none"> • calculation of mol of AgCl (1) • calculation of mass of Cl⁻ (1) • calculation of percentage of Cl⁻ (1) <p>OR</p> <ul style="list-style-type: none"> • calculation of % by mass of Cl in AgCl (1) • calculation of mass of Cl in residue (1) • calculation of percentage of Cl⁻ (1) | <p>Example of calculation:</p> <p>Method 1</p> $\text{mol AgCl} = \frac{0.226}{143.4} = 0.0015760 / 1.5760 \times 10^{-3}$ $\text{mass Cl}^{-} = 1.5760 \times 10^{-3} \times 35.5 = 0.055948 \text{ (g)}$ $\% \text{ Cl}^{-} = \frac{0.055948}{0.098} \times 100 = 57.09 / 57.1 / 57 \text{ (\%)}$ <p>Method 2</p> $\% \text{ by mass of Cl in AgCl} = \frac{35.5}{143.4} \times 100 = 24.756 \text{ (\%)}$ $\text{mass of Cl} = 24.756 \times 0.226 = 0.055948 \text{ (g)}$ $\% \text{ by mass of Cl}^{-} \text{ in residue} = \frac{0.055948}{0.098} \times 100 = 57.09 / 57.1 / 57 \text{ (\%)}$ <p>Correct answer with no working scores (3)</p> <p>Allow TE at each stage</p> <p>Allow alternative methods</p> <p>Ignore SF except 1 SF</p> <p>Accept fractions / correct working not evaluated for M1 and M2</p> | (3) |

(Total for Question 16 = 12 marks)

| Question Number | Answer | Additional Guidance | Mark | | | | | | |
|---|--|--|----------------|----------------|---|-----------|---|---------|-----|
| 17(a) | <ul style="list-style-type: none"> both classifications correct | <p>Example of table:</p> <table border="1" data-bbox="1330 284 1899 689"> <thead> <tr> <th data-bbox="1330 284 1666 355">Halogenoalkane</th> <th data-bbox="1666 284 1899 355">Classification</th> </tr> </thead> <tbody> <tr> <td data-bbox="1330 355 1666 523">  </td> <td data-bbox="1666 355 1899 523">secondary</td> </tr> <tr> <td data-bbox="1330 523 1666 689">  </td> <td data-bbox="1666 523 1899 689">primary</td> </tr> </tbody> </table> <p>Allow 2° for secondary Allow 1° for primary Ignore halogenoalkane</p> | Halogenoalkane | Classification |  | secondary |  | primary | (1) |
| Halogenoalkane | Classification | | | | | | | | |
|  | secondary | | | | | | | | |
|  | primary | | | | | | | | |

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|--|--|------|
| 17(b) | <p>An explanation that makes reference to the following points:</p> <ul style="list-style-type: none"> 2-chloro-2-methylpropane should react faster than 1-iodobutane because it is tertiary (1) 1-iodobutane should react faster than 2-chloro-2-methylpropane because the C–I bond enthalpy is lower than C–Cl (1) it is not possible to predict the relative effects of these two opposing factors / structure and bond enthalpies (1) | <p>Stand alone mark Allow tertiary / branched chain / more branched halogenoalkanes have a higher rate of hydrolysis than primary halogenoalkanes Do not award secondary for 2-chloro-2-methylpropane Do not award tertiary carbocation</p> <p>Stand alone mark Allow just ‘the C–I bond is weaker / has a lower bond enthalpy than C–Cl’ Ignore just ‘bonds in 1-iodobutane are weaker’ Do not award H–I / H–Cl bonds</p> <p>Conditional on two opposing factors If M1 and M2 scored, allow ‘so it is not possible to predict the relative rate of hydrolysis’</p> | (3) |

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|--|--|------|
| 17(c)(i) | <ul style="list-style-type: none"> ammonia (gas) would escape (from the condenser when heated under reflux) or to prevent ammonia (gas) escaping (from the condenser when heated under reflux) | <p>Ignore just ‘ammonia will evaporate’ Ignore concentrated alcoholic Ignore references to safety Ignore just ‘gas / reactant escapes’ Do not award any other substance escaping</p> | (1) |

| Question Number | Answer | Additional Guidance | Mark |
|--|---|--|------|
| 17(c)(ii) | <ul style="list-style-type: none"> • lone pair on N of NH₃ and curly arrow from lone pair on N to, or towards C (1) • dipole on C-Br and curly arrow from C-Br bond to, or just beyond, Br (1) • lone pair on N of NH₃ and curly arrow from lone pair on N to, or towards H (1) • curly arrow from N-H bond to, or towards N (1) | Penalise missing lone pair on N once only Penalise negative charge on NH ₃ once only Penalise half-arrow heads once only Do not award any charge / dipole on H Ignore any changes to final products | (4) |
| Example of mechanism: <div style="text-align: center; margin-top: 20px;">  </div> | | | |

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|--|--|------|
| 17(d) | <ul style="list-style-type: none"> <li data-bbox="360 293 1115 325">• calculation of amounts of KBr and H₂SO₄ (1) <li data-bbox="360 533 1115 676">• calculation of amount of C₂H₅OH and statement or implication that this is the limiting quantity (1) <li data-bbox="360 756 1115 788">• calculation of maximum mass of C₂H₅Br formed (1) | <p data-bbox="1167 252 1473 284">Example of calculation:</p> <p data-bbox="1167 304 1648 368">amount KBr = $\frac{14.90}{119} = 0.12521$ (mol)</p> <p data-bbox="1167 379 1677 443">amount H₂SO₄ = $\frac{16.35}{98.1} = 0.16667$ (mol)</p> <p data-bbox="1167 454 1865 486">Allow use of 98 for <i>M_r</i> of H₂SO₄ giving 0.16684 (mol)</p> <p data-bbox="1167 528 1834 592">amount C₂H₅OH = $\frac{4.65}{46} = 0.10109 / 0.10 / 0.1$ (mol)</p> <p data-bbox="1167 603 1906 703">and any indication that the limiting reagent is C₂H₅OH e.g. by use of mol of ethanol in M3</p> <p data-bbox="1167 751 1834 852">(maximum amount C₂H₅Br formed = 0.10109 (mol) maximum mass C₂H₅Br formed = 0.10109 x 108.9 = 11.008 / 11.01 / 11.0 / 11 (g)</p> <p data-bbox="1167 863 1845 895">Allow use of 109 for <i>M_r</i> of C₂H₅Br giving 11.018 (g)</p> <p data-bbox="1167 938 1861 970">Ignore SF except 1 SF, but allow 0.1 for mol C₂H₅OH</p> | (3) |

| Question Number | Acceptable Answers | Additional Guidance | Mark | | | | | | | | | | | | |
|--|---|--|---|---|---|-----|---|-----|---|---|---|---|---|---|-----|
| 17(e)* | <p>This question assesses a student’s ability to show a coherent and logically structured answer with linkages and fully-sustained reasoning.</p> <p>Marks are awarded for indicative content and for how the answer is structured and shows lines of reasoning.</p> <p>The following table shows how the marks should be awarded for indicative content.</p> <table border="1" data-bbox="349 616 864 959"> <thead> <tr> <th data-bbox="349 616 600 767">Number of indicative marking points seen in answer</th> <th data-bbox="600 616 864 767">Number of marks awarded for indicative marking points</th> </tr> </thead> <tbody> <tr> <td data-bbox="349 767 600 807">6</td> <td data-bbox="600 767 864 807">4</td> </tr> <tr> <td data-bbox="349 807 600 847">5–4</td> <td data-bbox="600 807 864 847">3</td> </tr> <tr> <td data-bbox="349 847 600 887">3–2</td> <td data-bbox="600 847 864 887">2</td> </tr> <tr> <td data-bbox="349 887 600 927">1</td> <td data-bbox="600 887 864 927">1</td> </tr> <tr> <td data-bbox="349 927 600 959">0</td> <td data-bbox="600 927 864 959">0</td> </tr> </tbody> </table> <p>The following table shows how the marks should be awarded for structure and lines of reasoning.</p> | Number of indicative marking points seen in answer | Number of marks awarded for indicative marking points | 6 | 4 | 5–4 | 3 | 3–2 | 2 | 1 | 1 | 0 | 0 | <p>Guidance on how the mark scheme should be applied:</p> <p>The mark for indicative content should be added to the mark for lines of reasoning. For example, an answer with five indicative marking points that is partially structured with some linkages and lines of reasoning scores 4 marks (3 marks for indicative content and 1 mark for partial structure and some linkages and lines of reasoning).</p> <p>If there are no linkages between points, the same five indicative marking points would yield an overall score of 3 marks (3 marks for indicative content and no marks for linkages).</p> | (6) |
| Number of indicative marking points seen in answer | Number of marks awarded for indicative marking points | | | | | | | | | | | | | | |
| 6 | 4 | | | | | | | | | | | | | | |
| 5–4 | 3 | | | | | | | | | | | | | | |
| 3–2 | 2 | | | | | | | | | | | | | | |
| 1 | 1 | | | | | | | | | | | | | | |
| 0 | 0 | | | | | | | | | | | | | | |

| | Number of marks awarded for structure of answer and sustained line of reasoning |
|---|---|
| Answer shows a coherent and logical structure with linkages and fully sustained lines of reasoning demonstrated throughout. | 2 |
| Answer is partially structured with some linkages and lines of reasoning. | 1 |
| Answer has no linkages between points and is unstructured. | 0 |

Comment:

Look for the indicative marking points first, then consider the mark for structure of answer and sustained line of reasoning

In general it would be expected that 5 or 6 indicative points would get 2 reasoning marks, and 3 or 4 indicative points would get 1 mark for reasoning, and 0, 1 or 2 indicative points would score zero marks for reasoning.

General points to note


If there is any incorrect chemistry, deduct mark(s) from the reasoning. If no reasoning mark(s) awarded do not deduct mark(s).

| | | | |
|--|---|--|--|
| | <p>Indicative content</p> <ul style="list-style-type: none"> • IP1 – Similarity Both reactions involve hydroxide ions / OH⁻ • IP2 – Type of reaction Reaction with aqueous solution is substitution and reaction with ethanolic solution is elimination • IP3 – Type of reagent (OH⁻ is a) nucleophile in aqueous solution and a base in ethanolic solution • IP4 – Products In aqueous solution propan-2-ol / an alcohol forms and in ethanolic solution propene / an alkene forms • IP5 – Equation in aqueous solution CH₃CHBrCH₃ + OH⁻ → CH₃CHOHCH₃ + Br⁻ • IP6 – Equation in ethanolic solution CH₃CHBrCH₃ + OH⁻ → CH₃CH=CH₂ + H₂O + Br⁻ | <p>Penalise use of incorrect halogenoalkane once only</p> <p>Allow OH⁻ shown with both reactions anywhere in the answer e.g. in the equations Allow both reactions need heat (under reflux)</p> <p>Ignore displacement for substitution</p> <p>Ignore dehydration for elimination Do not award dehydrogenation for elimination</p> <p>If IP2 and IP3 not awarded, allow 1 IP for just ‘nucleophilic substitution’ or ‘elimination by a base’</p> <p>This can be scored from the equations</p> <p>In IP5 and IP6, allow displayed formulae / any combination of displayed and structural formulae / skeletal formula Allow KOH / KBr / K⁺ + OH⁻ / K⁺ + Br⁻ The equations must be balanced Ignore state symbols even if incorrect Ignore mechanisms even if incorrect</p> | |
|--|---|--|--|

(Total for Question 17 = 18 marks)

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|---|---|------|
| 18(a)(i) | <ul style="list-style-type: none"> (temperature) 298 K / 25°C and (pressure) 1 atm / 100 kPa / 101 kPa / 1 x 10⁵ Pa / 1.01 x 10⁵ Pa | Allow 'a specified / stated temperature' Ignore just 'room temperature' Do not award 298°K Do not award incorrect pressure units e.g. 101 Pa | (1) |

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|---|--|------|
| 18(a)(ii) | <ul style="list-style-type: none"> molecular formula for 2,2,4-trimethylpentane (1) rest of equation correct conditional on use of C₈H₁₈ or use of structural / displayed / skeletal formula for 2,2,4-trimethylpentane (1) | Example of equation: $\text{C}_8\text{H}_{18} + 12\frac{1}{2}\text{O}_2 \rightarrow 8\text{CO}_2 + 9\text{H}_2\text{O}$ Accept 25/2 for 12½ Allow multiples e.g. $2\text{C}_8\text{H}_{18} + 25\text{O}_2 \rightarrow 16\text{CO}_2 + 18\text{H}_2\text{O}$ Ignore state symbols even if incorrect | (2) |

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|---|--|------|
| 18(a)(iii) | <ul style="list-style-type: none"> <li data-bbox="360 699 987 842">• y axis labelled enthalpy and products line drawn at a lower level than reactants line (1) <li data-bbox="360 1034 987 1066">• downwards arrow labelled with $\Delta_c H$ (1) | <p data-bbox="1032 252 1308 284">Example of diagram;</p>  <p data-bbox="1032 676 1883 778">Allow energy / H / enthalpy level as label for y axis Do not award ΔH / enthalpy change / energy change as label for y axis</p> <p data-bbox="1032 804 1883 874">Allow names / formulae of reactants and products but both must be there e.g. $C_8H_{18} + O_2$ for reactants and $CO_2 + H_2O$ for products</p> <p data-bbox="1032 900 1839 1002">Ignore missing / incorrect balancing numbers if formulae given Ignore label / missing label on x axis Ignore activation energy hump(s)</p> <p data-bbox="1032 1027 1839 1145">M2 Conditional on reactants higher than products Allow label as ΔH / -5461 / other label that indicates enthalpy change of combustion / reaction</p> <p data-bbox="1032 1171 1861 1235">Do not award double headed arrow / or just a line with no arrow / arrow labelled $-\Delta H$</p> | (2) |

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|--|--|------|
| 18(a)(iv) | <ul style="list-style-type: none"> • calculation of energy given out by 1 g (1) • calculation of energy given out by 1 cm³ (1) • calculation of energy given out by 1 dm³ (1) | <p>Example of calculation:</p> <p>Method 1 enthalpy change / g = $\frac{5461}{114} = 47.904$ (kJ)</p> <p>enthalpy change / cm³ = $47.904 \times 0.692 = 33.149$ (kJ) TE on M1</p> <p>enthalpy change / dm³ = $33.149 \times 1000 = 33\,149 / 33.149 \times 10^3$ (kJ) TE on M2</p> <p>Method 2 mass of 2,2,4-trimethylpentane in 1 dm³ = $0.692 \times 1000 = 692$ (g) (1) mol in 1 dm³ = $\frac{692}{114} = 6.0702$ (mol) (1) TE on M1 enthalpy change / dm³ = $6.0702 \times 5461 = 33\,149 / 33.149 \times 10^3$ (kJ) (1) TE on M2</p> <p>Allow alternative methods Correct answer with some working scores (3) Ignore SF except 1 SF Ignore minus sign Ignore units, even if incorrect</p> | (3) |

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|--|---|------|
| 18(b)(i) | <ul style="list-style-type: none"> • calculation of heat evolved • calculation of moles of ethanol used • working for heat evolved per mole • value of $\Delta_c H$ to 2 / 3 SF and negative sign and units | <p>Example of calculation:</p> <p>heat evolved = $100.0 \times 4.18 \times 13.2$ $= 5517.6 \text{ (J)} / 5.5176 \text{ kJ}$ Do not award $100.305 \times 4.18 \times 13.2 = 5534.4 \text{ (J)}$</p> <p>amount of ethanol = $\frac{0.305}{46} = 0.0066304 / 6.6304 \times 10^{-3} \text{ (mol)}$</p> <p>heat evolved per mole = $\frac{5.5176}{6.6304 \times 10^{-3}} \text{ (= 832.17)}$</p> <p>TE on M1 and M2</p> <p>$\Delta_c H = -830 / -832 \text{ kJ mol}^{-1}$ Allow units kJ/mol or $\frac{\text{kJ}}{\text{mol}}$ or kJ mol^{-1}</p> <p>Ignore letter case in units e.g. k or K, J or j Accept $-830000 / -832000 \text{ J mol}^{-1}$ TE on M3</p> <p>Ignore SF except 1 SF in M1, M2 and M3 Correct answer with some working to 2/3 SF with sign and units scores (4)</p> | (4) |

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|---|---|------|
| 18(b)(ii) | <ul style="list-style-type: none"> calculation of percentage error | <p>Example of calculation: $\frac{(2 \times 0.05 \times 100)}{13.2} = 0.75758 (\%)$</p> <p>Allow 0.7576 / 0.758 / 0.76 / 0.8 Correct answer with no working scores (1) Ignore signs Do not award 0.75 / 0.757 / 0.80</p> | (1) |

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|---|--|------|
| 18(b)(iii) | <p>An answer that makes reference to any two of the following points:</p> <ul style="list-style-type: none"> heat loss (to the surroundings) (1) incomplete combustion (of ethanol) (1) some ethanol evaporates (1) calculation does not take into account the heat capacity of the beaker (1) | <p>Allow insufficient oxygen for combustion Ignore not all of the ethanol was burned</p> <p>Ignore product(s) / water evaporates</p> <p>Allow some heat is used to heat up the beaker Ignore thermometer Ignore ethanol was impure Ignore water was not stirred Ignore no lid on beaker</p> | (2) |

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|--|--|------|
| 18(c) | <ul style="list-style-type: none"> <li data-bbox="360 328 1084 432">• calculation of bond energies of O–H and C–H (1) <li data-bbox="360 477 1084 512">• calculation of bond energy of C–O (1) <li data-bbox="360 628 1084 663">• calculation of bond energy of C–C (1) | <p data-bbox="1167 252 1473 284">Example of calculation:</p> <p data-bbox="1167 292 1301 323">Method 1</p> <p data-bbox="1167 328 1765 360">bond energy O–H = $928/2 = (+)464$ (kJ mol⁻¹)</p> <p data-bbox="1167 368 1218 400">and</p> <p data-bbox="1167 405 1778 437">bond energy C–H = $1740/4 = (+)435$ (kJ mol⁻¹)</p> <p data-bbox="1167 477 1727 544">bond energy C–O = $2105 - (3 \times 435) - 464$ = $(+)336$ (kJ mol⁻¹)</p> <p data-bbox="1167 552 1294 584">TE on M1</p> <p data-bbox="1167 624 1800 691">bond energy C–C = $3322 - (5 \times 435) - 464 - 336$ = $(+)347$ (kJ mol⁻¹)</p> <p data-bbox="1167 699 1301 730">Method 2</p> <p data-bbox="1167 735 1711 767">$3322 - 2105 = 1217 = \text{C–C} + 2 \times \text{C–H}$ (1)</p> <p data-bbox="1167 772 1868 804">bond energy $2 \times \text{C–H} = 1740/2 = (+)870$ (kJ mol⁻¹) (1)</p> <p data-bbox="1167 809 1711 841">C–C = $1217 - 870 = (+)347$ (kJ mol⁻¹) (1)</p> <p data-bbox="1167 880 1671 912">M3 TE on M1 and M2 in both methods</p> <p data-bbox="1167 952 1749 984">Correct answer with some working scores (3)</p> | (3) |

| Question Number | Answer | Additional Guidance | Mark |
|-----------------|--|--|------|
| 18(d) | <p>An answer that makes reference to the following points:</p> <ul style="list-style-type: none"> • there are fewer (gas) molecules on the right hand side / more (gas) molecules on the left hand side (1) • so the equilibrium position will shift to the right / product side and the equilibrium yield of ethanol will increase (1) | <p>Ignore effect of temperature Ignore effect on rate of reaction</p> <p>Allow moles for molecules Allow 2 (gas) molecules on the left and 1 (gas) molecule on the right Allow higher pressure favours the side with fewer (gas) molecules</p> <p>Allow forward reaction is favoured and the equilibrium yield of ethanol will increase</p> | (2) |

(Total for Question 18 = 20 marks)

Total for Section C = 20 marks

Total for paper = 80 marks

