

Mark Scheme (Final)

Summer 2008

GCE

GCE Chemistry (6246/01B)

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Using the mark scheme

- 1 / means that the responses are alternatives and either answer should receive full credit.
- 2 () means that a phrase/word is not essential for the award of the mark, but helps the examiner to get the sense of the expected answer.
- 3 [] words inside square brackets are instructions or guidance for examiners.
- 4 Phrases/words in **bold** indicate that the meaning of the phrase or the actual word is **essential** to the answer.
- 5 ecf/TE/cq (error carried forward) means that a wrong answer given in an earlier part of a question is used correctly in answer to a later part of the same question.

In general, an inference should follow an observation.

Apparatus and Materials

Apparatus

Each candidate will require:

1. 50.0 cm³ burette, stand and clamp, with small funnel for filling, white tile and a small beaker for draining burette;
2. two 250 cm³ conical flasks;
3. 25.0 cm³ pipette and safety filler;
4. seven test tubes and one boiling tube in a test tube rack;
5. one 10 cm³ and two 25 cm³ measuring cylinders;
6. a supply of dropping pipettes;
7. a 250 cm³ beaker of hot water at about 70 °C to be used as a water bath.

Materials

Each candidate will require:

- (a)* 200 cm³ of aqueous sodium thiosulphate of concentration 0.110 mol dm⁻³ labelled **Solution F**;
- (b)* 200 cm³ of aqueous potassium manganate(VII) of concentration 0.020 mol dm⁻³ labelled **Solution G**. The concentration of this solution is **not** to be disclosed to candidates;
- (c)* 10 cm³ of approximately 0.25 mol dm⁻³ aqueous aluminium chloride labelled **Solution of H**. The identity of this solution is **not** to be disclosed to candidates;
- (d)* 5 cm³ of approximately 0.25 mol dm⁻³ aqueous copper(II) chloride, labelled **Solution of I**. The identity of this solution is **not** to be disclosed to candidates;
- (e)* 5 cm³ of butanone labelled **J**. The identity of this compound is **not** to be disclosed to candidates;
- (f) 100 cm³ of dilute sulphuric acid of concentration approximately 1.0 mol dm⁻³, labelled **Dilute sulphuric acid**;
- (g) 100 cm³ of aqueous potassium iodide of concentration approximately 0.50 mol dm⁻³ labelled **Aqueous potassium iodide**;
- (h) 10 cm³ of dilute sodium hydroxide; concentration approximately 1.0 mol dm⁻³;
- (i) access to a small bottle of Universal Indicator solution;
- (j) 20 cm³ of dilute aqueous ammonia; concentration approximately 2.0 mol dm⁻³;
- (k) 5 cm³ of dilute nitric acid; concentration approximately 2.0 mol dm⁻³;
- (l) 5 cm³ of aqueous silver nitrate; concentration approximately 0.05 mol dm⁻³;
- (m) 5 cm³ of 2,4-dinitrophenylhydrazine solution. This may be made by adding 0.1 g of the solid reagent to 45 cm³ of water and 5 cm³ of concentrated hydrochloric acid, stirring and filtering if necessary. Alternatively centres may prepare this reagent using their own procedure providing the reagent gives a positive test with butanone;
- (n) 5 cm³ dilute sulphuric acid; concentration approximately 1.0 mol dm⁻³ (for Question 3);
- (o) 5 cm³ of aqueous potassium dichromate(VI); concentration approximately 0.20 mol dm⁻³;
- (p) 10 cm³ of aqueous sodium hydroxide; concentration approximately 0.50 mol dm⁻³. Label this solution **0.50 mol dm⁻³ sodium hydroxide for Q3(c)**;
- (q) 10 cm³ of iodine/potassium iodide solution made by adding 2 g iodine to 6 g potassium iodide dissolved in 100 cm³ water and labelled **aqueous iodine**;
- (r) 20 cm³ of freshly prepared aqueous starch; concentration approximately 1% labelled **starch**;
- (s) a supply of distilled water.

For home centres (ONLY), the chemicals identified with an asterisk (*) will be sent by a firm of manufacturing chemists.

Question Number	Correct Answer	Acceptable Answers	Reject	Mark															
1.(a)	<p>Table 1 Check subtractions and averaging arithmetic, correcting if necessary. All volumes recorded to 0.05 cm³ (1) <i>ALLOW one slip but withhold this mark if any readings are in the wrong boxes.</i> <i>ALLOW 0 as initial volume NOT 50 as initial volume</i> All subtractions correct (1)</p> <p><i>[✓✓top RHS of Table 1]</i></p> <p>Mean titre For correct averaging of chosen values / choosing identical values and for recording the average correct to 2 or 3 dps or to nearest 0.05 cm³ (1) Do not penalise missing 2/3rd dp if already penalised in Table 1.</p> <p><i>[✓ by the mean in space or near the dotted line in paragraph below]</i></p> <p>Accuracy If the candidate has made an arithmetical error in the Table 1 volumes used in the mean or in averaging the examiner must calculate a new average.</p> <ul style="list-style-type: none"> • For an averaging error simply calculate a new value using the candidate's chosen titres. • If a wrongly subtracted titre has been used in the mean then choose any two identical titres or take an average of the closest two titres. <p>Calculate the difference(d) between the candidate's mean titre and that of the examiner or supervisor.</p> <p>Examiner's titre = 22.70 cm³ (to be confirmed at standardisation)</p> <p>Award marks for accuracy as follows.</p> <table border="1" data-bbox="260 1765 858 1895"> <tr> <td>Difference</td> <td>±0.20</td> <td>±0.30</td> <td>±0.40</td> <td>±0.50</td> </tr> <tr> <td>(d) =</td> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <td>Mark</td> <td>4</td> <td>3</td> <td>2</td> <td>1</td> </tr> </table>	Difference	±0.20	±0.30	±0.40	±0.50	(d) =					Mark	4	3	2	1			10
Difference	±0.20	±0.30	±0.40	±0.50															
(d) =																			
Mark	4	3	2	1															

Question Number	Correct Answer	Acceptable Answers	Reject	Mark
2.(a)	Observations White precipitate (1) Dissolves / disappears (in excess NaOH) / colourless solution (1) Inference Zinc / Zn^{2+} , aluminium / Al^{3+} } any two (1) lead(II) / Pb^{2+} Ignore Cd^{2+} / Sn^{2+} / Sn^{4+} / Sb^{3+}	Soluble in excess/ goes clear	Symbols Zn, Al, Pb.	3

Question Number	Correct Answer	Acceptable Answers	Reject	Mark
2.(b)	Observation Any red (1) Inferences Aluminium / Al^{3+} (1) Acidic (since only 3+ ion of Al, Zn, Pb) Acidic - Stand alone mark (1)			3

Question Number	Correct Answer	Acceptable Answers	Reject	Mark
2.(c)	Observations White precipitate (1) Insoluble in excess NH_3 (1) Inference $Al(OH)_3$ / $[Al(H_2O)_3(OH)_3]$ /aluminium hydroxide (1)			3

Question Number	Correct Answer	Acceptable Answers	Reject	Mark
2.(d)	Observation White precipitate (1) Inference chloride / Cl^- (1)		Chlorine / Cl	2

Question Number	Correct Answer	Acceptable Answers	Reject	Mark
2.(e)	$AlCl_3$	Al_2Cl_6		1

Question Number	Correct Answer	Acceptable Answers	Reject	Mark
2.(f)	Observations (any) blue precipitate (1) Dissolves/ disappears in excess (1) Deep(er) blue solution (1) Inferences $Cu(OH)_2$ / $[Cu(H_2O)_4(OH)_2]$ (1) $[Cu(H_2O)_2(NH_3)_4]^{2+}$ (1)	Soluble in excess/ goes clear $[Cu(NH_3)_4]^{2+}$	$[Cu(NH_3)_6]^{2+}$	5

Question Number	Correct Answer	Acceptable Answers	Reject	Mark
2.(g)	$CuCl_2$			1

Question Number	Correct Answer	Acceptable Answers	Reject	Mark
3.(a)	Observation Yellow / orange precipitate (1) Inference Carbonyl / C=O / >C=O / both of aldehyde or ketone (1)	Yellow-orange		2

Question Number	Correct Answer	Acceptable Answers	Reject	Mark
3.(b)	Observation Stays orange / no change (1) Inferences Ketone / not aldehyde if follows A or K in (a) (1) Not oxidised / no redox / does not reduce $\text{Cr}_2\text{O}_7^{2-}$ (1)	No reaction	Just "nothing" Reject cq on wrong colour	3

Question Number	Correct Answer	Acceptable Answers	Reject	Mark
3.(c)	Observation (pale) Yellow precipitate (1) Inferences triiodomethane / Iodoform / CHI_3 (1) Methyl ketone / CH_3CO (1)	Cream ppte	CH_3I Methyl secondary alcohol / ethanol / ethanal	3

Question Number	Correct Answer	Acceptable Answers	Reject	Mark
3.(d)(i)	<i>m/e</i> 72 (1) Structure $\text{CH}_3\text{—CH}_2\text{—C=O}$ / $\text{C}_2\text{H}_5\text{—C=O}$ (1) CH_3 CH_3 Ignore positive charge		$\text{CH}_3\text{COCH}_2\text{CH}_3$ Species with negative charge	2

Question Number	Correct Answer	Acceptable Answers	Reject	Mark
3.(d)(ii)	$\text{CH}_3\text{CH}_2\text{CO}^+$ / $\text{CH}_2\text{COCH}_3^+$		Formula with no positive charge $\text{C}_3\text{H}_5\text{O}^+$	1

Question Number	Correct Answer	Acceptable Answers	Reject	Mark
4.	1 ✓ (Add Na ₂ SO ₄ to all five): the one that gives white ppt is BaCl ₂ 2 ✓ Add BaCl ₂ to other four solutions. White ppt with AgNO ₃ 3 ✓ Add AgNO ₃ to remaining three solutions 4 ✓ White ppts with NaCl and ZnCl ₂ 5 ✓ Brown ppt with NH ₃ / remaining one is NH ₃ 6 ✓ Add NH ₃ to NaCl and ZnCl ₂ 7 ✓ White ppt with ZnCl ₂	No white ppt with NH ₃		7

OR

Question Number	Correct Answer	Acceptable Answers	Reject	Mark
4.	1 ✓ (Add Na ₂ SO ₄ to all five): the one that gives white ppt is BaCl ₂ 2 ✓ Add BaCl ₂ to other four solutions. White ppt with AgNO ₃ 3 ✓ Add remaining three solutions to AgCl ppt 4 ✓ AgCl dissolves in NH ₃ 5 ✓ Add AgNO ₃ to remaining two solutions 6 ✓ White ppt with both ZnCl ₂ and NaCl 7 ✓ Add excess NH ₃ : ZnCl ₂ ppt dissolves			7

OR

Question Number	Correct Answer	Acceptable Answers	Reject	Mark
4.	1 ✓ (Add Na ₂ SO ₄ to all five): the one that gives white ppt is BaCl ₂ 2 ✓ Add BaCl ₂ to other four solutions. White ppt with AgNO ₃ 3 ✓ Add remaining three solutions to AgCl ppt 4 ✓ AgCl dissolves in NH ₃ 5 ✓ Add NH ₃ to NaCl and ZnCl ₂ 6 ✓ White ppt with ZnCl ₂ 7 ✓ No ppt with NaCl/ NaCl remaining			7

OR

Question Number	Correct Answer	Acceptable Answers	Reject	Mark
4.	1 ✓ (Add Na ₂ SO ₄ to all five): two white ppts - BaCl ₂ and AgNO ₃ 2 ✓ Distinguish between ppts 3 ✓ Add AgNO ₃ to remaining three solutions 4 ✓ White ppts with NaCl and ZnCl ₂ 5 ✓ Brown ppt with NH ₃ / remaining one is NH ₃ 6 ✓ Add NH ₃ to NaCl and ZnCl ₂ 7 ✓ White ppt with ZnCl ₂	No white ppt with NH ₃		7