

# Mark Scheme Summer 2009

GCE

## GCE Chemistry (8CH07) International Supplement 2

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information please call our Customer Services on + 44 1204 770 696, or visit our website at [www.edexcel.com](http://www.edexcel.com).

If you have any subject specific questions about the content of this Mark Scheme that require the help of a subject specialist, you may find our **Ask The Expert** email service helpful.

Ask The Expert can be accessed online at the following link:

<http://www.edexcel.com/Aboutus/contact-us/>

(If you are calling from outside the UK please dial + 44 1204 770 696 and state that you would like to speak to the **Science** subject specialist).

Summer 2009

Publications Code US021182

All the material in this publication is copyright

© Edexcel Ltd 2009

## Contents

1.	6CH07/01 Mark Scheme	5
----	----------------------	---



## 6CH07/01

Question Number	Correct Answer	Reject	Mark
1 (a)(i)	No (colour) change (to flame) OR no flame colour <b>Accept</b> No colour	White flame	1

Question Number	Correct Answer	Reject	Mark
1 (a)(ii)	Effervescence / bubbling / fizzing <i>IGNORE</i> hissing		1

Question Number	Correct Answer	Reject	Mark
1 (a)(iii)	<b>Observation:</b> (Lime water) turns milky / cloudy or white precipitate (formed) (1) <b>Accept</b> White solid (formed) / chalky  <b>Inference:</b> carbon dioxide / CO <sub>2</sub> (1)	Turns white	2

Question Number	Correct Answer	Reject	Mark
1 (a)(iv)	<b>Observation:</b> White precipitate (formed) (1) <b>Accept</b> White solid / crystal (formed) <i>IGNORE</i> references to heat given out and to precipitate insoluble in excess <b>Inference:</b> Magnesium hydroxide / Mg(OH) <sub>2</sub> (1)	White substance  Confirms magnesium present	2

Question Number	Correct Answer	Reject	Mark
1 (b)(i)	Lithium / Li <sup>+</sup> (1) Strontium / Sr <sup>2+</sup> (1) <b>Accept</b> Calcium / Ca <sup>2+</sup> (1)	Rubidium Li, Sr, Ca (penalise use of element symbol once only)	2

Question Number	Correct Answer	Reject	Mark
1 (b)(ii)	Dissolves (in the ammonia) (to form a colourless solution) <b>Accept</b> Soluble <i>IGNORE</i> references to dilute ammonia	Partially dissolves	1

Question Number	Correct Answer	Reject	Mark
1 (b)(iii)	<b>Observation:</b> Brown or red-brown or orange (1) <b>Inference:</b> Bromine / Br <sub>2</sub> (1)	Red  Bromide (for bromine)	2

Question Number	Correct Answer	Reject	Mark
1 (b)(iv)	<b>From:</b> Orange or yellow <b>To:</b> blue or green or blue-green		1

Question Number	Correct Answer	Reject	Mark
1 (b)(v)	<b>Mark two points independently</b> (Hydrogen) bromide oxidized / bromine oxidation number increased (from -1 to 0) / changes from -1 to 0 / Bromide loses an electron / (hydrogen) bromide is a reducing agent (1)  sulfuric acid reduced / sulfur oxidation number decreases (from (+)6 to (+)4) / changes from (+)6 to (+)4 / sulfate gains electrons / sulfuric acid is an oxidizing agent (1) <b>Accept</b> (+)VI to (+)IV sulfate reduced		2

Question Number	Correct Answer	Reject	Mark
2 (a)(i)	<b>Vertical</b> line at 3.5 minutes intersects extrapolated top line (1) <b>Horizontal</b> extrapolated lower line and 66-69 minus 20-22 = $\Delta T$ (1)	incorrect or no extrapolation line joining points at 3 & 4 minutes & extrapolated to intersect top line (0)	2

Question Number	Correct Answer	Reject	Mark
2 (a)(ii)	$(1 \times 50 \times 10^{-3}) = 0.0500$ <i>IGNORE sf</i>		1

Question Number	Correct Answer	Reject	Mark
2 (a)(iii)	$65.4 \times 0.05 = 3.27$ (g) / 3.3 (g) <b>Accept</b> $65 \times 0.05 = 3.25$ (g) / 3.3 (g)		1

Question Number	Correct Answer	Reject	Mark
2 (a)(iv)	Heat capacity negligible <b>Accept:</b> low specific heat capacity or zinc absorbs less heat than solution	Mass negligible No heat absorbed by zinc All heat absorbed by solution	1

Question Number	Correct Answer	Reject	Mark														
2 (a)(v)	<p>50 x 4.18 x <math>\Delta T</math> (1) (<math>\Delta T</math> CQ on (a)(i)) Penalise use of incorrect mass here only. <b>IGNORE</b> <math>c = 4.2 \text{ Jg}^{-1}\text{C}^{-1}</math></p> <table border="1"> <thead> <tr> <th><math>\Delta T</math></th> <th>Heat energy (kJ)</th> </tr> </thead> <tbody> <tr><td>44</td><td>9.20</td></tr> <tr><td>45</td><td>9.41</td></tr> <tr><td>46</td><td>9.61</td></tr> <tr><td>47</td><td>9.82</td></tr> <tr><td>48</td><td>10.0(3)</td></tr> <tr><td>49</td><td>10.2(4)</td></tr> </tbody> </table> <p>(units if given must be consistent) (1) <b>IGNORE</b> sf except 1 sf</p>	$\Delta T$	Heat energy (kJ)	44	9.20	45	9.41	46	9.61	47	9.82	48	10.0(3)	49	10.2(4)		2
$\Delta T$	Heat energy (kJ)																
44	9.20																
45	9.41																
46	9.61																
47	9.82																
48	10.0(3)																
49	10.2(4)																

Question Number	Correct Answer	Reject	Mark																					
2 (a)(vi)	<p>= – answer to (a)(v) <math>\div</math> answer to (a)(ii) (1) For 0.05 mol:</p> <table border="1"> <thead> <tr> <th><math>\Delta T</math></th> <th>Heat energy (kJ)</th> <th><math>\Delta H / \text{kJ mol}^{-1}</math></th> </tr> </thead> <tbody> <tr><td>44</td><td>9.20</td><td>-180</td></tr> <tr><td>45</td><td>9.41</td><td>-190</td></tr> <tr><td>46</td><td>9.61</td><td>-190</td></tr> <tr><td>47</td><td>9.82</td><td>-200</td></tr> <tr><td>48</td><td>10.0(3)</td><td>-200</td></tr> <tr><td>49</td><td>10.2(4)</td><td>-200</td></tr> </tbody> </table> <p>CQ on moles from 2 (a)(ii) Sign <b>and</b> 2 sf (1) [this mark may be awarded for any <b>calculated</b> value]</p>	$\Delta T$	Heat energy (kJ)	$\Delta H / \text{kJ mol}^{-1}$	44	9.20	-180	45	9.41	-190	46	9.61	-190	47	9.82	-200	48	10.0(3)	-200	49	10.2(4)	-200		2
$\Delta T$	Heat energy (kJ)	$\Delta H / \text{kJ mol}^{-1}$																						
44	9.20	-180																						
45	9.41	-190																						
46	9.61	-190																						
47	9.82	-200																						
48	10.0(3)	-200																						
49	10.2(4)	-200																						

Question Number	Correct Answer	Reject	Mark
2 (b)(i)	Ensure equilibration or steady temperature or same temperature (as surroundings)	More accurate temperature	1

Question Number	Correct Answer	Reject	Mark
2 (b)(ii)	To allow for cooling / a cooling correction / to compensate for heat loss	Temperature correction To determine maximum temperature change More accurate temperature / $\Delta T$	1

Question Number	Correct Answer	Reject	Mark
2 (b)(iii)	Low heat capacity Good insulator Poor heat conductor Low mass Absorbs less heat	Low <b>specific</b> heat capacity	1

Question Number	Correct Answer	Reject	Mark
2 (b)(iv)	Ensure uniform temperature <b>Accept</b> to spread out heat (uniformly) <i>IGNORE</i> references to mixing reagents, increasing reaction rate, enabling reactants to react and temperature accuracy.		1

Question Number	Correct Answer	Reject	Mark
2 (b)(v)	Burette / pipette / measuring cylinder / volumetric or graduated flask	Beaker / conical flask	1

Question Number	Correct Answer	Reject	Mark
2 (b)(vi)	Lid on polystyrene cup/ Increase insulation <b>Accept</b> Put cup in a beaker	Magnetic stirrer	1

Question Number	Correct Answer	Reject	Mark
2 (c)	Zn>Pb>Cu OR Zinc <b>displaces</b> both so is most reactive (1) The more exothermic / negative ( <b>accept</b> 'the larger') the $\Delta H$ the greater the difference in reactivity (so lead more reactive than copper) (1) If the order of reactivity is reversed maximum 1	Answers in <b>just</b> terms of reactivity or electrochemical series Generalised answers References to energy or enthalpy <b>required</b> for the reaction	2




Question Number	Correct Answer	Reject	Mark
3 (a)(i)	<b>Observation:</b> Steamy/misty /white fumes (1) <b>Explanation:</b> Hydrogen chloride / HCl formed OR chloroalkene / chloro- compound formed OR Substitution reaction with OH (1)	Smoke or solid  Hydrochloric acid  Chloroalkane  Just OH / alcohol group reacts (with PCl <sub>5</sub> )	2

Question Number	Correct Answer	Reject	Mark
3 (a)(ii)	<b>Observation:</b> Purple to colourless or brown (1)  <b>Explanation:</b> Addition to C=C /alkene OR oxidation of C=C /alkene OR OH / alcohol group oxidised (1) <b>Accept</b> Reacts with C=C to form diol or with OH to form an aldehyde or a carboxylic acid OR manganate(VII) / permanganate / MnO <sub>4</sub> <sup>-</sup> to MnO <sub>2</sub> (if brown) or Mn(II) / Mn <sup>2+</sup> (if decolourized)	Just 'decolourized'  'Reacts' alone instead of addition or oxidation 'Due to the presence of C=C /alkene / OH' A oxidised	2


Question Number	Correct Answer	Reject	Mark
3 (a)(iii)	<b>Observation:</b> Orange or yellow or brown ( <b>accept</b> red-brown) to colourless (1) <b>Explanation:</b> (Bromine) addition to C=C /alkene (Bromine) reacts with C=C /alkene to form dibromoalkanol / dibromo compound (1)	'pink' instead of purple 'clear' instead of colourless Just 'decolourized' Reaction alone instead of addition  dibromoalkane	2

Question Number	Correct Answer	Reject	Mark
3 (b)	$  \begin{array}{cccc}  \text{H} & \text{H} & \text{H} & \text{H} \\    &   &   &   \\  \text{H}-\text{C} & -\text{C} & -\text{C} & -\text{C}-\text{O}-\text{H} \\    &   &   &   \\  \text{Br} & \text{Br} & \text{H} & \text{H}  \end{array}  $ <p>OR</p> $  \begin{array}{cccc}  \text{H} & \text{H} & \text{H} & \text{H} \\    &   &   &   \\  \text{H}-\text{C} & -\text{C} & -\text{C} & -\text{C}-\text{O}-\text{H} \\    &   &   &   \\  \text{Br} & \text{O} & \text{H} & \text{H} \\  &   & & \\  & \text{H} & &   \end{array}  $ <p>Accept</p> $  \begin{array}{cccc}  \text{H} & \text{H} & \text{H} & \text{H} \\    &   &   &   \\  \text{H}-\text{C} & -\text{C} & -\text{C} & -\text{C}-\text{O}-\text{H} \\    &   &   &   \\  \text{O} & \text{Br} & \text{H} & \text{H} \\    & & & \\  \text{H} & & &   \end{array}  $ <p>Accept OH for O—H</p>		1

Question Number	Correct Answer	Reject	Mark
4 (a)	Funnel with neck & tap (1) <i>IGNORE</i> stopper Organic layer above aqueous layer (1) Stand alone See diagrams	Conical / filter / Buchner funnel with tap Funnel too full to be stoppered	2



YES



NO

Question Number	Correct Answer	Reject	Mark
4 (b)(i)	(Organic & aqueous) layers are immiscible <b>OR</b> consequence of not shaking e.g. layers form <b>Accept</b> 'to ensure layers mix <i>IGNORE</i> references to rate	Just 'to mix reagents' Explanations in terms of density differences	1


Question Number	Correct Answer	Reject	Mark
4 (b)(ii)	Neutralize (excess) acid / H <sup>+</sup> <b>Accept</b> remove acid / H <sup>+</sup> React with acid <i>IGNORE</i> Use of HCl for hydrochloric acid release of CO <sub>2</sub>	Just 'neutralize / neutralization	1

Question Number	Correct Answer	Reject	Mark
4 (b)(iii)	Carbon dioxide / CO <sub>2</sub> / gas is formed (1) Release pressure / pressure builds up (1)		2

Question Number	Correct Answer	Reject	Mark
4 (b)(iv)	Drying agent or to remove water	Dehydrating agent	1

Question Number	Correct Answer	Reject	Mark
4 (b)(v)	To pour off the liquid leaving the solid behind	Pour / pour carefully / transfer	1

Question Number	Correct Answer	Reject	Mark
4 (c)(i)	Bulk of the thermometer bulb adjacent to the outlet leading to the condenser (see diagram)		1



Question Number	Correct Answer	Reject	Mark
4 (c)(ii)	Water in through the lower tube and out through the upper If words are used (water in & water out) ignore the direction of any arrows		1

Question Number	Correct Answer	Reject	Mark
4 (d)	<p>Mass of alcohol = <math>5 \times 0.805 = 4.025</math> (g) (1)</p> <p>Moles of alcohol = <math>4.025 \div 88 = 0.0457</math></p> <p>= moles of 2-chloro-2-methylbutane</p> <p>Mass 2-chloro-2-methylbutane (100% yield) = <math>0.0457 \times 106.5 = 4.87</math></p> <p>70% yield = <math>4.87 \times \frac{70}{100} = 3.41</math> g (1)</p> <p>ignore sf except for 1 sf</p> <p>If the molar masses are transposed penalise once (answer = 2.32 g)</p> <p>Correct answer and some working (2)</p>		2



Further copies of this publication are available from  
Edexcel UK Regional Offices at [www.edexcel.org.uk/sfc/feschools/regional/](http://www.edexcel.org.uk/sfc/feschools/regional/)  
or International Regional Offices at [www.edexcel-international.org/sfc/academic/regional/](http://www.edexcel-international.org/sfc/academic/regional/)

For more information on Edexcel qualifications, please visit [www.edexcel-international.org/quals](http://www.edexcel-international.org/quals)  
Alternatively, you can contact Customer Services at [www.edexcel.org.uk/ask](http://www.edexcel.org.uk/ask) or on + 44 1204 770 696

Edexcel Limited. Registered in England and Wales no.4496750  
Registered Office: One90 High Holborn, London, WC1V 7BH