

Write your name here	
Surname	Other names
Centre Number	Candidate Number
<input type="text"/>	<input type="text"/>
Edexcel GCE	
Chemistry	
Advanced Subsidiary	
Unit 3B: Chemistry Laboratory Skills I Alternative	
Thursday 13 May 2010 – Morning Time: 1 hour 15 minutes	Paper Reference 6CH07/01
Candidates may use a calculator.	Total Marks
<input type="text"/>	<input type="text"/>

Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
– *there may be more space than you need.*

Information

- The total mark for this paper is 50.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*
- You will be assessed on your ability to organise and present information, ideas, descriptions and arguments clearly and logically, including your use of grammar, punctuation and spelling.
- A Periodic Table is printed on the back cover of this paper.

Advice

- Read each question carefully before you start to answer it.
- Keep an eye on the time.
- Try to answer every question.
- Check your answers if you have time at the end.

N35693A

©2010 Edexcel Limited.
7/7/7/2/



Turn over ►

edexcel 
advancing learning, changing lives

Answer ALL the questions. Write your answers in the spaces provided.

1 Compound A is a white solid that contains one Group 1 cation and one anion.

(a) (i) Describe how you would carry out a flame test on compound A.

(3)

.....

.....

.....

.....

.....

.....

(ii) In a flame test, compound A gives a red flame. Deduce the formula of the cation present.

(1)

.....

(b) On prolonged strong heating, compound A forms a white solid, B, and a gas. The gas turns limewater milky.

(i) Identify, by name or formula, the compound that is dissolved in water to make limewater.

(1)

.....

(ii) Suggest the formula for the anion in compound A. Justify your answer.

(2)

.....

.....

.....



(c) When water is added to the white solid, **B**, it dissolves completely and exothermically to form solution **C**.

(i) Identify, by name or formula, the anion present in **B**.

(1)

(ii) Identify, by name or formula, the anion present in **C**.

(1)

(iii) Suggest a test for the anion present in **C**. Give the result of your test.

(2)

Test

Result

(d) Suggest the **formula** of compound **A**.

(1)

(Total for Question 1 = 12 marks)



2 This question is about two isomeric halogenoalkanes, **P** and **Q**.

(a) A hot aqueous solution of silver nitrate is added to each halogenoalkane. Both halogenoalkanes react to form a yellow precipitate.

(i) Identify, by name or formula, this yellow precipitate.

(1)

(ii) The isomers have relative molecular mass 169.9. Deduce the molecular formula of the isomers.

(1)

(iii) Halogenoalkane **P** forms the yellow precipitate faster than halogenoalkane **Q**. Draw a displayed formula for halogenoalkane **P**.

(1)

(iv) Give the name or structural formula of the alcohol, **R**, formed by the reaction of halogenoalkane, **P**, with hot aqueous silver nitrate.

(1)



(b) When **R** is boiled with a mixture of potassium dichromate(VI) and dilute sulfuric acid, the organic product **S** forms.

(i) Give the colour change you would expect to see.

(2)

From **to**

(ii) Give the **name** of **S**.

(1)

(iii) Give the type of reaction involved in the conversion of **R** to **S**.

(1)

(Total for Question 2 = 8 marks)



BLANK PAGE



3 The purity of a sample of potassium iodate(V) was determined by titration.

The steps of the experimental procedure are as follows.

1. 0.100 g of the sample was dissolved in water in a beaker and the solution made up to 100 cm³ in an appropriate flask.
2. A 10.0 cm³ portion of this solution of potassium iodate(V) was transferred to a conical flask.
3. An excess of both potassium iodide solution and sulfuric acid were then added to the conical flask. This produced a solution, **T**, containing iodine.
4. Solution **T** was titrated with 0.0200 mol dm⁻³ sodium thiosulfate solution using a suitable indicator.
5. Steps 2, 3 and 4 were repeated twice.

(a) (i) Name the piece of apparatus used to remove the 10.0 cm³ portions of potassium iodate(V) solution (step 2).

(1)

(ii) Name the indicator you would use for the titration and give the colour change you would expect to see (step 4).

(2)

Indicator

Colour change from to

(b) The following results were obtained for the titrations.

Titration number	1	2	3
Final burette reading / cm ³	19.50	33.20	46.95
Initial burette reading / cm ³	5.00	19.50	33.20
Titre / cm ³			

(i) Complete the table by calculating the titres.

(1)




(ii) Explain why the correct value for the mean titre is 13.73 cm³.

(1)

(iii) Calculate the number of moles of sodium thiosulfate in the mean titre.

(1)


(c) The ionic equation for the reaction between iodine and sodium thiosulfate in the titration is shown below.



Calculate the number of moles of iodine in solution **T** using this equation and your answer to (b)(iii).

(1)

(d) The ionic equation for the reaction of iodate(V) ions with iodide ions is shown below.



Using this equation and your answer to (c), calculate the number of moles of iodate(V) ions which reacted to produce solution **T**.

(1)



(e) (i) Name the appropriate flask used in step 1.

(1)

(ii) Describe how you would make up exactly 100 cm³ of potassium iodate(V) solution in this flask, ready for step 2.

(3)

(iii) Calculate the number of moles of potassium iodate(V) in 100 cm³ of the solution, using your answer to (d).

(1)

(iv) Calculate the mass of potassium iodate(V) in the sample.

[Assume the molar mass of potassium iodate(V) is 214 g mol⁻¹]

(1)

(v) Calculate the percentage purity of the sample.

(1)

(f) Suggest the most significant hazard in step 3.

(1)


(Total for Question 3 = 16 marks)



- 4 An experiment to prepare 0.100 mol of 1-bromobutane uses the reaction of butan-1-ol with hydrogen bromide.

Hydrogen bromide is formed in the reaction mixture from potassium bromide and moderately concentrated sulfuric acid.

The process has an 80 % yield after purification of the 1-bromobutane.



The steps of the experimental procedure are as follows.

1. Add measured amounts of potassium bromide and butan-1-ol to 10 cm³ of water into a 50 cm³ two-necked flask.
2. Fit the two-necked flask with a reflux condenser and a tap funnel.
3. Immerse the flask in a beaker of cold water and add 10 cm³ of concentrated sulfuric acid from the tap funnel, a few drops at a time.
4. Remove the flask from the cold water and close the tap on the tap funnel. Heat the mixture under reflux for 30 minutes.
5. Allow the mixture to cool. Then set up the apparatus for distillation. Boil the mixture and collect the distillate in a measuring cylinder.
6. Transfer the distillate to a separating funnel. The distillate consists of two layers, an aqueous layer and impure 1-bromobutane. Separate the two layers.
7. Wash the impure 1-bromobutane with concentrated hydrochloric acid and separate the two layers.
8. Wash the 1-bromobutane layer with sodium hydrogencarbonate solution, releasing any gas formed.
9. Collect the 1-bromobutane layer in a conical flask and add anhydrous sodium sulfate.
10. Decant the 1-bromobutane into a 50 cm³ flask.

Data

Property	Butan-1-ol	1-bromobutane	Water
Density / g cm ⁻³	0.81	1.3	1.0
Molar mass / g mol ⁻¹	74	137	18
Boiling temperature / °C	117.3	101.7	100.0



(a) (i) Show, by calculation, that 0.125 mol of butan-1-ol is needed to make 0.100 mol of 1-bromobutane.

(2)

(ii) Calculate the volume of 0.125 mol of butan-1-ol, in cm^3 .

(2)

(iii) Calculate the minimum mass of potassium bromide required in step 1.

[The molar mass of potassium bromide is 119 g mol^{-1}]


(1)



(b) Complete and label the diagram below of the apparatus assembled in steps 1, 2 and 3.

[You may assume that the apparatus is suitably clamped.]

(4)



(c) (i) State, with a reason, whether the upper or lower layer contains 1-bromobutane in step 6.

(1)

(ii) The product is washed with concentrated hydrochloric acid in step 7 to remove unreacted butan-1-ol. In step 8, why is the product then washed with sodium hydrogencarbonate solution and what causes a build up of gas?

(2)

(d) (i) What further step is necessary to purify the 1-bromobutane obtained in step 10?

(1)

(ii) How does the step in (d)(i) give information about the purity of the product?

(1)

(Total for Question 4 = 14 marks)

TOTAL FOR PAPER = 50 MARKS



BLANK PAGE



BLANK PAGE



N 3 5 6 9 3 A 0 1 5 1 6

The Periodic Table of Elements

		1	2	3	4	5	6	7	0 (8)										
		<div style="border: 1px solid black; padding: 2px; display: inline-block;"> 1.0 H hydrogen 1 </div>																	
		Key relative atomic mass atomic symbol name atomic (proton) number																	
		(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)
6.9	9.0	Li lithium 3	Be beryllium 4	Sc scandium 21	Ti titanium 22	V vanadium 23	Cr chromium 24	Mn manganese 25	Fe iron 26	Co cobalt 27	Ni nickel 28	Cu copper 29	Zn zinc 30	B boron 5	C carbon 6	N nitrogen 7	O oxygen 8	F fluorine 9	He helium 2
23.0	24.3	Na sodium 11	Mg magnesium 12	Y yttrium 39	Zr zirconium 40	Nb niobium 41	Mo molybdenum 42	Tc technetium [98]	Ru ruthenium 44	Rh rhodium 45	Pd palladium 46	Ag silver 47	Cd cadmium 48	Al aluminium 13	Si silicon 14	P phosphorus 15	S sulfur 16	Cl chlorine 17	Ar argon 18
39.1	40.1	K potassium 19	Ca calcium 20	La* lanthanum 57	Hf hafnium 72	Ta tantalum 73	W tungsten 74	Re rhenium 75	Os osmium 76	Ir iridium 77	Pt platinum 78	Au gold 79	Hg mercury 80	Ge germanium 32	As arsenic 33	Se selenium 34	Br bromine 35	Kr krypton 36	Xe xenon 54
85.5	87.6	Rb rubidium 37	Sr strontium 38	La* lanthanum 57	Zr zirconium 40	Nb niobium 41	Mo molybdenum 42	Tc technetium [98]	Ru ruthenium 44	Rh rhodium 45	Pd palladium 46	Ag silver 47	Cd cadmium 48	In indium 49	Sb antimony 51	Te tellurium 52	I iodine 53	Xe xenon 54	Rn radon 86
132.9	137.3	Cs caesium 55	Ba barium 56	La* lanthanum 57	Hf hafnium 72	Ta tantalum 73	W tungsten 74	Re rhenium 75	Os osmium 76	Ir iridium 77	Pt platinum 78	Au gold 79	Hg mercury 80	Tl thallium 81	Pb lead 82	Bi bismuth 83	Po polonium 84	At astatine 85	Rn radon 86
[223]	[226]	Fr francium 87	Ra radium 88	Ac* actinium 89	Rf rutherfordium 104	Db dubnium 105	Sg seaborgium 106	Bh bohrium 107	Hs hassium 108	Mt meitnerium 109	Ds darmstadtium 110	Rg roentgenium 111	Elements with atomic numbers 112-116 have been reported but not fully authenticated						
140	141	Ce cerium 58	Pr praseodymium 59	Nd neodymium 60	Pm promethium 61	Eu europium 63	Gd gadolinium 64	Tb terbium 65	Dy dysprosium 66	Ho holmium 67	Er erbium 68	Tm thulium 69	Yb ytterbium 70	Lu lutetium 71	Fm fermium 100	Md mendelevium 101	No nobelium 102	Lr lawrencium 103	Th thorium 90
232	[231]	Pa protactinium 91	U uranium 92	Np neptunium 93	Pu plutonium 94	Am americium 95	Cm curium 96	Bk berkelium 97	Cf californium 98	Es einsteinium 99	Fm fermium 100	Md mendelevium 101	No nobelium 102	Lr lawrencium 103	[254]	[256]	[254]	[257]	[257]

* Lanthanide series

* Actinide series

