

**OXFORD CAMBRIDGE AND RSA EXAMINATIONS****Advanced GCE****CHEMISTRY****Unifying Concepts in Chemistry**

Friday

**24 JANUARY 2003**

Afternoon

**2816/01**

1 hour 15 minutes

Candidates answer on the question paper.

Additional materials:

*Data Sheet for Chemistry*

Scientific calculator

Candidate Name

Centre Number

Candidate  
Number

--	--	--

**TIME** 1 hour 15 minutes**INSTRUCTIONS TO CANDIDATES**

- Write your name in the space above.
- Write your Centre number and Candidate number in the boxes above.
- Answer all the questions.
- Write your answers in the spaces provided on the question paper.
- Read each question carefully and make sure you know what you have to do before starting your answer.

**INFORMATION FOR CANDIDATES**

- The number of marks is given in brackets [ ] at the end of each question or part question.
- You will be awarded marks for the quality of written communication where this is indicated in the question.
- You may use a scientific calculator.
- You may use the *Data Sheet for Chemistry*.
- You are advised to show all the steps in any calculations.


FOR EXAMINER'S USE		
Qu.	Max.	Mark
1	17	
2	17	
3	12	
4	14	
<b>TOTAL</b>	<b>60</b>	

**This question paper consists of 10 printed pages and 2 blank pages.**

Answer all the questions.

- 1 Syngas is a mixture of carbon monoxide and hydrogen gases, used as a feedstock for the manufacture of methanol.

A dynamic equilibrium was set up between carbon monoxide, CO, hydrogen, H<sub>2</sub>, and methanol, CH<sub>3</sub>OH. The equilibrium system is shown by Equilibrium 1.1 below.



Equilibrium 1.1

The equilibrium concentrations of the three components of this equilibrium are shown below.

component	CO(g)	H <sub>2</sub> (g)	CH <sub>3</sub> OH(g)
equilibrium concentration /mol dm <sup>-3</sup>	3.1 × 10 <sup>-3</sup>	2.4 × 10 <sup>-2</sup>	2.6 × 10 <sup>-5</sup>

- (a) State two features of a system that is in *dynamic equilibrium*.

.....  
 .....  
 ..... [2]

- (b) (i) Write the expression for K<sub>c</sub> for this equilibrium system.

[2]

- (ii) Calculate the numerical value of K<sub>c</sub> for this equilibrium.

[2]

- (c) The pressure was increased whilst keeping the temperature constant. The system was left to reach equilibrium. The equilibrium position of Equilibrium 1.1 shifted to the right.

- (i) Explain why the equilibrium moved to the right.

.....  
 ..... [2]

- (ii) What is the effect, if any, on K<sub>c</sub>?

..... [1]

(iii) State and explain the effect on the rates of the forward and reverse reactions

- when the pressure was first changed
- when the system reached equilibrium.

.....  
.....  
.....  
.....  
.....  
.....

[4]

(d) The temperature was increased whilst keeping the pressure constant. The system was left to reach equilibrium. The value of  $K_c$  for Equilibrium 1.1 decreased.

(i) Explain what happens to the equilibrium position of Equilibrium 1.1.

.....  
.....  
.....  
.....  
.....  
.....

[2]

(ii) Deduce the sign of the enthalpy change for the forward reaction shown in Equilibrium 1.1. Explain your reasoning.

.....  
.....

[1]

(iii) Explain how the partial pressure of  $\text{CH}_3\text{OH(g)}$  would change as the system moves towards equilibrium.

.....  
.....  
.....

[1]

[Total: 17]

- 2** Nitrous oxide,  $\text{N}_2\text{O}$ , is a colourless gas with a mild, pleasing odour and sweet taste. It is widely used as a propellant in aerosol cans of whipped cream.

- (a) Nitrous oxide is formed when ammonium nitrate,  $\text{NH}_4\text{NO}_3$ , is gently heated.



- (i) What mass of  $\text{N}_2\text{O}$  is formed by heating 100 g of  $\text{NH}_4\text{NO}_3$ ?

[3]

- (ii) What happens to the oxidation number of each nitrogen from  $\text{NH}_4\text{NO}_3$  in this reaction?

.....  
 .....  
 .....


[2]

- (b) When heated strongly, nitrous oxide decomposes into its elements.



This reaction is first order with respect to  $\text{N}_2\text{O}$ .

The graph below shows how nitrous oxide decomposes with time at constant temperature.



- (i) Explain how the graph confirms that this reaction is first order with respect to N<sub>2</sub>O.

.....  
.....  
.....  
.....  
..... [3]

- (ii) Write the expression for the rate equation of this reaction.

[1]

- (iii) Use the graph to work out the rate of reaction, in mol dm<sup>-3</sup>s<sup>-1</sup>, at 70 seconds. Show clearly your working on the graph.

rate = ..... mol dm<sup>-3</sup>s<sup>-1</sup> [2]

- (iv) Calculate the rate constant for this reaction. State the units.

k = ..... units ..... [2]

- (v) What evidence is there that the mechanism of this reaction takes place in more than a single step?

.....  
.....  
.....  
..... [2]

- (c) N<sub>2</sub>O is occasionally injected into the engines of racing cars to give more power and exceptional acceleration. The N<sub>2</sub>O decomposes exothermically to N<sub>2</sub> and O<sub>2</sub>.

Suggest two reasons why this reaction provides an extra boost to the engine.

.....  
.....  
..... [2]

[Total: 17]

**3** In this question, one mark is available for the quality of written communication.

(a) Describe what is meant by the following terms used in acid-base chemistry.

- The Bronsted-Lowry theory of acids and bases.
  - Conjugate acid-base pairs.
  - Dilute and weak acids.

Illustrate your answer by choosing suitable examples of acids and bases. Write equations where appropriate.

[7]

### Quality of Written Communication [1]

(b) The acid dissociation constant  $K_a$  of hydrocyanic acid, HCN, is  $4.9 \times 10^{-10} \text{ mol dm}^{-3}$ .

(i) Write an expression for the acid dissociation constant of HCN.

[1]

(ii) Calculate the pH of a  $0.010 \text{ mol dm}^{-3}$  solution of hydrocyanic acid.

[3]

[Total: 12]

**4 Organic acids occur widely in nature.**

- (a) Butanoic acid,  $\text{CH}_3(\text{CH}_2)_2\text{COOH}$ , is a straight-chain organic acid, largely responsible for the odour of rancid butter.

Caprylic acid is another straight-chain organic acid. It is produced in the body in small amounts as an antifungal agent in human sweat.

- (i) Some caprylic acid was isolated from human sweat and analysed. The sample of caprylic acid had the percentage composition by mass:

C, 66.7%; H, 11.1%; O, 22.2%.  $M_r = 144$ .

Calculate the molecular formula of caprylic acid and suggest its structural formula.

[4]

- (ii) Tracker dogs are trained to follow odours such as the characteristic blend of organic acids in the sweat from a person's feet. A dog is able to detect extremely small quantities of these acids.

Sweat containing equal amounts of butanoic and caprylic acids produces more butanoic acid vapour than caprylic acid vapour.

Suggest a reason for this. Explain your answer.

.....  
.....  
.....  
.....

[2]

- (b) Compound A is a straight-chain organic acid. A chemist analysed a sample of acid A by the procedure below.

The chemist first prepared a  $250\text{ cm}^3$  solution of **A** by dissolving 10.8 g of **A** in water.

In a titration,  $25.00 \text{ cm}^3$   $0.500 \text{ mol dm}^{-3}$  NaOH were neutralised by exactly  $21.40 \text{ cm}^3$  of solution A.

- Calculate the pH of the NaOH(aq) used in the titration.  $K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{dm}^{-6}$ .
  - Use the results to calculate the molar mass of acid A and suggest its identity.

[8]

[Total: 14]

**BLANK PAGE**

**BLANK PAGE**

