

OXFORD CAMBRIDGE AND RSA EXAMINATIONS Advanced Subsidiary GCE

CHEMISTRY 2812

Chains and Rings

Wednesday

4 JUNE 2003

Morning

1 hour

Candidates answer on the question paper.

Additional materials:

Data Sheet for Chemistry

Scientific calculator

Candidate Name	Centre Number	Candidate Number

TIME 1 hour

INSTRUCTIONS TO CANDIDATES

- Write your name in the space above.
- Write your Centre number and Candidate number in the boxes above.
- Answer all the questions.
- Write your answers in the spaces provided on the question paper.
- Read each question carefully and make sure you know what you have to do before starting your answer.

INFORMATION FOR CANDIDATES

- The number of marks is given in brackets [] at the end of each question or part question.
- You will be awarded marks for the quality of written communication where this is indicated in the question.
- You may use a scientific calculator.
- You may use the Data Sheet for Chemistry.
- You are advised to show all the steps in any calculations.

FOR EXA	MINER	'S USE
Question Number	Max.	Mark
1	15	
2	12	
3	12	
4	6	
5	15	
TOTAL	60	


This question paper consists of 10 printed pages and 2 blank pages.

Answer all the questions.

1

Co	mpoi	und A is an alcohol with the following percentage composition by mass:
С,	60.09	%; H, 13.3%; O, 26.7%.
(a)	(i)	Calculate the empirical formula of compound A.
		[2]
	(ii)	The relative molecular mass of compound A is 60. Show that this agrees with the molecular formula C_3H_8O .
		[1]
(b)	Cor of e	mpound A is one of two possible isomers that are both alcohols. Draw the structure each isomer.
		[2]
(c)	Con	npound A was refluxed with an excess of an oxidising agent to form compound B .
		State a suitable oxidising agent for this reaction.
		[1]
	(ii)	State the colour change you would see during this oxidation.
		From [2]
	(iii)	Explain what is meant by the term reflux.
		[1]

(d) The infra red spectrum of compound B is shown below.




[Total: 15]

4

For Examine Use

2 Dibromoethene, C₂H₂Br₂, has two structural isomers **C** and **D**, shown below.



Br C=C Br


(a) (i) Name compound C.

		[1]
(ii)	What is the empirical formula of compound D?	
	***************************************	[1]

(b) The reaction between either C or D and Br₂ can be used to show the presence of the C=C double bond.

(i) State what you would see when isomer C reacts with Br₂.

- _______[1]
- (ii) State the type of mechanism for the reaction between isomer C and Br₂.
- (c) Isomers C and D can both behave as alkenes. For each of the following reactions, state the conditions, if any, and identify the organic products that can be formed.
 - (i) Isomer C reacts with H₂.



organic product

conditions[2]

5

For Examiner's Use

(ii) Isomer C reacts with HBr.

[2]

(iii) Isomer D reacts with H₂O.

onditions

......[3]

[Total: 12]

3

6

For	
Examine	į
Use	

In t	nis question, one mark is available for the quality of written communication.
(a)	Explain why methane does not react with reagents such as nucleophiles or electrophiles.
	[2]
(b)	Name and describe the mechanism for the reaction between methane and bromine.
	State the type of bond fission involved in the mechanism and explain why it is difficult to produce a single organic product.
	In your answer, include any relevant equations and conditions.
	•••••••••••••••••••••••••••••••••••••••
	••••
	•••••
	[9]
	Quality of Written Communication [1]
	(Total: 12)

7

For
Examine
Use

a) ((i)	Write a balanced equation for the hydrolysis of 1-bromobutane, C ₄ H ₉ Br.
((ii)	Describe, with the aid of curly arrows and relevant dipoles, the mechanism for th hydrolysis of 1-bromobutane with hydroxide ions, OH ⁻ .
		Try die Grand Commence Commenc
		[3
6	com eac	pared by an experiment using AgNO ₃ solution. Aqueous AgNO ₃ was added t
€ 8	com eac app	pared by an experiment using AgNO ₃ solution. Aqueous AgNO ₃ was added the halogenoalkane. The rates of hydrolysis were measured by timing the earance of the silver halide precipitate.
€ 8	com eac app	pared by an experiment using AgNO ₃ solution. Aqueous AgNO ₃ was added the halogenoalkane. The rates of hydrolysis were measured by timing the earance of the silver halide precipitate. Place the three halogenoalkanes in the order of their rates of hydrolysis with the
e	com eac app	pared by an experiment using AgNO ₃ solution. Aqueous AgNO ₃ was added the halogenoalkane. The rates of hydrolysis were measured by timing the earance of the silver halide precipitate. Place the three halogenoalkanes in the order of their rates of hydrolysis with the compound with the fastest rate first.
e	com eac app	pared by an experiment using AgNO ₃ solution. Aqueous AgNO ₃ was added the halogenoalkane. The rates of hydrolysis were measured by timing the earance of the silver halide precipitate. Place the three halogenoalkanes in the order of their rates of hydrolysis with the compound with the fastest rate first. fastest
6 8 8	com eac app	pared by an experiment using AgNO ₃ solution. Aqueous AgNO ₃ was added the halogenoalkane. The rates of hydrolysis were measured by timing the earance of the silver halide precipitate. Place the three halogenoalkanes in the order of their rates of hydrolysis with the compound with the fastest rate first. fastest
6 8	corr eac app (i)	spared by an experiment using AgNO ₃ solution. Aqueous AgNO ₃ was added the halogenoalkane. The rates of hydrolysis were measured by timing the earance of the silver halide precipitate. Place the three halogenoalkanes in the order of their rates of hydrolysis with the compound with the fastest rate first. fastest
6 8	corr eac app (i)	spared by an experiment using AgNO ₃ solution. Aqueous AgNO ₃ was added to the halogenoalkane. The rates of hydrolysis were measured by timing the earance of the silver halide precipitate. Place the three halogenoalkanes in the order of their rates of hydrolysis with the compound with the fastest rate first. fastest
6 8	corr eac app (i)	Place the three halogenoalkanes in the order of their rates of hydrolysis with the compound with the fastest rate first. fastest

8

For Examiner's Use

5 The question below concerns some of the chemistry involved in the brewing of beer.

Yeast ferments glucose, $C_6H_{12}O_6$, anaerobically (without oxygen) to produce ethanol and carbon dioxide.

(a) Balance the following equation for the anaerobic fermentation of glucose.

$$C_6H_{12}O_6 \rightarrow C_2H_5OH + CO_2$$

[1]

- **(b)** A batch containing 36.0 kg of glucose, $C_6H_{12}O_6$, produced 2.30 kg of ethanol, C_2H_5OH .
 - (i) Calculate the number of moles of glucose used.

[2]

(ii) Use the equation in (a) to deduce the maximum number of moles of ethanol that can be produced from 36.0 kg of glucose.

$$M_{\rm r} \, {\rm C_2H_5OH} = 46$$

[1]

(iii) Calculate the actual number of moles of ethanol produced by this batch fermentation.

[1]

(iv) Calculate the percentage yield of ethanol.

[1]

(c) The ethanol present in an alcoholic drink passes into the bloodstream and the liver is involved in breaking down the ethanol. The first stage of this breakdown involves the following reaction.

$$CH_3CH_2OH + [O] \longrightarrow CH_3CHO + H_2O.$$
 reaction 5.1

CH₃CHO is then oxidised to other products.

(i) Name the functional group present in CH₃CHO.

(ii) Suggest one possible oxidation product that can be formed from CH₃CHO.

...... [1]

C

For Examiner Use

(d) Methylated spirit contains ethanol and methanol, CH₃OH. Methylated spirit is unfit for drinking because an oxidation product of methanol causes liver damage and blindness. The methanol is oxidised in the body in a similar oxidation to that of ethanol shown in **reaction 5.1**.

Write an equation for the oxidation of methanol.

______[2]

(e) Methanol can be added to petrol.

- (i) Write a balanced equation for the complete combustion of methanol.
- (ii) Suggest why methanol is added to petrol.
-[1]
- (f) Methanol, CH₃OH, can be reacted with carbon monoxide to produce ethanoic acid, CH₃COOH.

Construct an equation for this reaction.

(g) Much of the ethanoic acid produced is used in the manufacture of the monomer ethenyl ethanoate (vinyl acetate). The monomer is then used to produce the polymer, polyvinyl acetate (PVA), part of which is shown below.

(i) Draw a circle around the repeat unit in PVA.

[1]

(ii) Draw the structure of the monomer ethenyl ethanoate (vinyl acetate).

[1]

[Total: 15]

10

BLANK PAGE