## OXFORD CAMBRIDGE AND RSA EXAMINATIONS

**Advanced Subsidiary GCE** 

CHEMISTRY

2811

Foundation Chemistry

Wednesday

8 JUNE 2005

Morning

1 hour

RECOGNISING ACHIEVEMENT

Candidates answer on the question paper.
Additional materials:

Data Sheet for Chemistry
Scientific Calculator

Candidate Name	Centre Number	Candidate Number

### TIME 1 hour

### **INSTRUCTIONS TO CANDIDATES**

- Write your name in the space above.
- Write your Centre number and Candidate number in the boxes above.
- Answer all the questions.
- Write your answers in the spaces provided on the question paper.
- Read each question carefully and make sure you know what you have to do before starting your answer.

## INFORMATION FOR CANDIDATES

- The number of marks is given in brackets [ ] at the end of each question or part question.
- You will be awarded marks for the quality of written communication where this is indicated in the question.
- You may use a scientific calculator.
- You may use the Data Sheet for Chemistry.
- You are advised to show all the steps in any calculations.

FOR EXAMINER'S USE		
Qu.	Max.	Mark
1	16	
2	13	
3	8	
4	7	
5	16	
TOTAL	60	

2

#### For Examiner's Use

## Answer all the questions.

1 The element titanium, Ti, atomic number 22, is a metal that is used in the aerospace industry for both airframes and engines.

A sample of titanium for aircraft construction was analysed using a mass spectrometer and was found to contain three isotopes,  $^{46}$ Ti,  $^{47}$ Ti and  $^{48}$ Ti. The results of the analysis are shown in Table 1.1 below.

Table 1.1

isotope	<sup>46</sup> Ti	<sup>47</sup> Ti	<sup>48</sup> Ti
relative isotopic mass	46.00	47.00	48.00
percentage composition	8.9	9.8	81.3

(a)	(i)	Expla	in the term <i>iso</i>	topes.				
	(ii)	Comp	olete the table I		s of two of the			[1]
			isotope	protons	neutrons	electrons		
			<sup>46</sup> Ti					
			<sup>47</sup> Ti					[2]
(b)	titaı	nium.	information in answer to thre			ive atomic ma	ss of this san	iple of
						. ~	· J.	
						300	1	·
						ı		[2]
(c)	Coi	mplete	the electronic	configuration o	of a titanium ato	om.		
	1s <sup>2</sup>	2s²2p <sup>6</sup>	S					[1]

For Examiner's Use

(a)	ита	nium nas metaliic bonding.
	(i)	Explain what is meant by metallic bonding. Use a diagram in your answer.
	(ii)	How does metallic bonding allow titanium to conduct electricity?
		[1]
(e)	A s	tudent reacted 1.44 g of titanium with chlorine to form 5.70 g of a chloride X.
	(i)	How many moles of Ti atoms were reacted?
		[1]
	(ii)	How many moles of Cl atoms were reacted?
		[2]
	(iii)	Determine the empirical formula of <b>X</b> .
		[1]
	(iv)	Construct a balanced equation for the reaction between titanium and chlorine.
	<b>(</b> )	
	(v)	At room temperature, <b>X</b> is a liquid which does <b>not</b> conduct electricity. What does this information suggest about the bonding and structure in <b>X</b> ?
		[O]
		[2]
		[Total: 16]

4

For	
Examine 3	5
Hea	

( <b>a</b> ) Pre	edict the formula of radium chloride.
	[1]
	erre and Marie Curie extracted radium from radium chloride by reduction. plain what is meant by <i>reduction</i> , using this reaction as an example.
••••	<u> </u>
••••	[2]
(c) Ra	dium reacts vigorously when added to water.
Ra	(s) + $2H_2O(I) \rightarrow Ra(OH)_2(aq) + H_2(g)$
(i)	Use the equation to predict <b>two</b> observations that you would <b>see</b> during this reaction.
	[2]
(ii)	Prodict a nH value for this solution
(11)	Predict a pH value for this solution.
(11)	·
(d) Re	actions of the Group 2 metals involve removal of electrons. The electrons are
(d) Re rer tre	actions of the Group 2 metals involve removal of electrons. The electrons are noved more easily as the group is descended and this helps to explain the increasing and in reactivity.
(d) Re rer tre	actions of the Group 2 metals involve removal of electrons. The electrons are noved more easily as the group is descended and this helps to explain the increasing and in reactivity.  The removal of one electron from each atom in 1 mole of gaseous radium atoms is
(d) Re rer tre	actions of the Group 2 metals involve removal of electrons. The electrons are noved more easily as the group is descended and this helps to explain the increasing and in reactivity.  The removal of one electron from each atom in 1 mole of gaseous radium atoms is
(d) Re rer tre	actions of the Group 2 metals involve removal of electrons. The electrons are noved more easily as the group is descended and this helps to explain the increasing and in reactivity.  The removal of one electron from each atom in 1 mole of gaseous radium atoms is called the
(d) Re rer tre	actions of the Group 2 metals involve removal of electrons. The electrons are noved more easily as the group is descended and this helps to explain the increasing and in reactivity.  The removal of one electron from each atom in 1 mole of gaseous radium atoms is called the
(d) Reretre	actions of the Group 2 metals involve removal of electrons. The electrons are noved more easily as the group is descended and this helps to explain the increasing and in reactivity.  The removal of one electron from each atom in 1 mole of gaseous radium atoms is called the
(d) Reretre	actions of the Group 2 metals involve removal of electrons. The electrons are noved more easily as the group is descended and this helps to explain the increasing and in reactivity.  The removal of one electron from each atom in 1 mole of gaseous radium atoms is called the
(d) Reretre	actions of the Group 2 metals involve removal of electrons. The electrons are noved more easily as the group is descended and this helps to explain the increasing and in reactivity.  The removal of one electron from each atom in 1 mole of gaseous radium atoms is called the
(d) Reretre	actions of the Group 2 metals involve removal of electrons. The electrons are noved more easily as the group is descended and this helps to explain the increasing and in reactivity.  The removal of one electron from each atom in 1 mole of gaseous radium atoms is called the

For Examiner's Use

			3
ē	acic	i. He	In that a stomach-ache and needed to take something to neutralise excess stomache decided to take some Milk of Magnesia, which is an aqueous suspension of turn hydroxide, ${\rm Mg(OH)}_2$ .
(	(a)		main acid in the stomach is hydrochloric acid, HCI(aq), and the unbalanced ation for the reaction that takes place with Milk of Magnesia is shown below.
			Mg(OH) <sub>2</sub> (s) +HC $l$ (aq) $\longrightarrow$ MgC $l$ <sub>2</sub> (aq) +H <sub>2</sub> O(l)
		Bala abo	ance the equation by adding numbers where necessary in the unbalanced equation ve.
(	(b)	0.10	student's stomach contained 500 cm <sup>3</sup> of stomach fluid with an acid concentration of 08 mol dm <sup>-3</sup> . The student swallowed some Milk of Magnesia containing 2.42 g OH) <sub>2</sub> . He wondered whether this dose was sufficient to neutralise the stomach acid.
		Ass	ume that all the acid in the stomach fluid was 0.108 mol dm <sup>-3</sup> hydrochloric acid.
		(i)	How many moles of HCl were in the 500 cm <sup>3</sup> of stomach fluid?
		(ii)	Calculate the mass of Mg(OH) <sub>2</sub> necessary to neutralise this stomach fluid.
	ı	(iii)	[3] Determine whether the student swallowed too much, too little, or just the right amount of Milk of Magnesia to neutralise the stomach acid.
(	(c)		ewing chalk has been used for many years to combat excess stomach acid and
			gestion tablets often contain calcium carbonate, CaCO <sub>3</sub> . Suggest, with the aid of an ation, how these tablets work.

[Total: 8]

6

4

For Examir.er's Use

	lorine d iodir	is used in the preparation of many commercially important materials such as bleach ne.
(a)	Blea aqu	ach is a solution of sodium chlorate(I), NaOCI, made by dissolving chlorine in eous sodium hydroxide.
		$Cl_2(g) + 2NaOH(aq) \rightarrow NaOCl(aq) + NaCl(aq) + H_2O(l)$
		ermine the changes in oxidation number of chlorine during the preparation of bleach comment on your results.
	••••	
	•••••	
		[3]
(b)	very	ne is extracted commercially from seawater with chlorine gas. Seawater contains y small quantities of dissolved iodide ions, which are oxidised to iodine by the orine gas.
	(i)	Write an ionic equation for the reaction that has taken place.
		[2]
	(ii)	Use your understanding of electronic structure to explain why chlorine is a stronger oxidising agent than iodine.
		[2]
		[Total: 7]

8

For Examiner's Use

(c) The atomic radii of nitrogen and oxygen are shown below.

element	nitrogen	oxygen
atomic radius /nm	0.075	0.073

Explain why a nitrogen atom is larger than an oxygen atom.
[4]
Quality of Written Communication [1]

[Total: 16]

## **END OF QUESTION PAPER**

For Examiner's Use

5 In this question, one mark is available for the quality of use and organisation of scientific terms.

Nitrogen and oxygen are elements in Period 2 of the Periodic Table. The hydrogen compounds of oxygen and nitrogen, H<sub>2</sub>O and NH<sub>3</sub>, both form hydrogen bonds.

(a) (i) Draw a diagram containing two H<sub>2</sub>O molecules to show what is meant by hydrogen bonding. On your diagram, show any lone pairs present and relevant dipoles.

		[3]
	(ii)	State and explain <b>two</b> anomalous properties of water resulting from hydrogen bonding.
		[4]
(b)	The	'dot-and-cross' diagram of an ammonia molecule is shown below.
		Ha N aH
	Pred	dict, with reasons, the bond angle in an ammonia molecule.
	*****	
	*****	
	••••	
	•••••	