

Mark Scheme 2816/01
January 2006

Question	Expected Answers	Marks
1 (a)	partial dissociation: $\text{HCOOH} = \text{H}^+ + \text{HCOO}^-$ ✓	[1]
(b) (i)	$\text{pH} = -\log(1.55 \times 10^{-3}) = 2.81/2.8$ ✓ [H ⁺] deals with negative indices over a very wide range/ pH makes numbers manageable /removes very small numbers ✓	[2]
(ii)	$K_a = \frac{[\text{H}^+(\text{aq})][\text{HCOO}^-(\text{aq})]}{[\text{HCOOH}(\text{aq})]}$ ✓ (state symbols not needed)	[1]
(iii)	$K_a = \frac{[\text{H}^+(\text{aq})]^2}{[\text{HCOOH}(\text{aq})]} = \frac{(1.55 \times 10^{-3})^2}{0.015}$ ✓ $= 1.60 \times 10^{-4} \text{ (mol dm}^{-3}\text{)} \checkmark$ $\text{p}K_a = -\log K_a = -\log(1.60 \times 10^{-4}) = 3.80$ ✓	[3]
(iv)	Percentage dissociating = $\frac{(1.55 \times 10^{-3}) \times 100}{0.015} = 10.3\%$ / 10% ✓ (working not required)	[1]
(c) (i)	$\text{HCOOH} + \text{NaOH} \longrightarrow \text{HCOONa} + \text{H}_2\text{O}$ ✓ state symbols not needed	[1]
(ii)	$n(\text{HCOOH}) = 0.0150 \times 25.00/1000 = 3.75 \times 10^{-4}$ ✓ volume of NaOH(aq) that reacts is 30 cm^3 ✓ so $[\text{NaOH}] = 3.75 \times 10^{-4} \times 1000/30 = 0.0125 \text{ mol dm}^{-3}$ ✓	[2]
(iii)	$K_w = [\text{H}^+(\text{aq})][\text{OH}^-(\text{aq})]$ ✓ $\text{pH} = -\log(1 \times 10^{-14}/0.0125) = 12.10/12.1$ ✓ (calc 12.09691001)	[3]
(iv)	metacresol purple ✓ pH range coincides with pH change during sharp rise OR pH 6-10 /coincides with equivalence point/end point ✓	[2]
		Total: 16

Question	Expected Answers	Marks												
2 (a)	$K_c = \frac{[HI]^2}{[H_2][I_2]}$ ✓	[1]												
(b) (i)	<table style="border: none; margin-left: 20px;"> <tr> <td style="padding-right: 20px;">H₂</td> <td style="padding-right: 20px;">I₂</td> <td>HI</td> </tr> <tr> <td>0.30</td> <td>0.20</td> <td>0</td> </tr> <tr> <td>0.14</td> <td>0.04</td> <td>0.32</td> </tr> <tr> <td></td> <td style="text-align: center;">✓</td> <td style="text-align: center;">✓</td> </tr> </table>	H ₂	I ₂	HI	0.30	0.20	0	0.14	0.04	0.32		✓	✓	[2]
H ₂	I ₂	HI												
0.30	0.20	0												
0.14	0.04	0.32												
	✓	✓												
(b) (ii)	$K_c = \frac{0.32^2}{0.14 \times 0.04} = 18.28571429$ ✓ = 18 (to 2 sig figs) ✓ no units ✓ (or ecf based on answers to (i) and/or (a))	[3]												
(c)	K_c is constant ✓ Composition of mixture is the same ✓	[2]												
(d)	(Forward) reaction is exothermic (ora) ✓ because equilibrium moves to the left / K_c is less ✓	[2]												
(e) (i)	$I_2(aq) + H_2S(g) \longrightarrow 2HI(aq) + S(s)$ species and balance ✓ state symbols: accept (s) for I ₂ ; (aq) for H ₂ S ✓	[2]												
(e) (ii)	amount I ₂ reacted = 1.89 mol / HI formed = 3.44 mol ✓ theoretical amount HI produced = 3.78 mol/484 g ✓ $\% \text{ yield} = \frac{3.44 \times 100}{3.78} \text{ or } \frac{440 \times 100}{484} = 91.0 \%$ ✓	[3]												
(e) (iii)	$[HI] = \frac{3.44 \times 1000}{750} = 4.58/4.59 \text{ mol dm}^{-3}$ ✓ pH = -log 4.59 = -0.66 ✓	[2]												
		Total: 17												

Question	Expected Answers	Marks
3	<p>From graph, constant half-life ✓ Therefore 1st order w.r.t. $[\text{CH}_3\text{COCH}_3]$ ✓</p> <p>From table, rate doubles when $[\text{H}^+]$ doubles ✓ Therefore 1st order w.r.t. $[\text{H}^+]$ ✓</p> <p>From table, rate stays same when $[\text{I}_2]$ doubles ✓ Therefore zero order w.r.t. $[\text{I}_2]$ ✓ Order with no justification does not score.</p> <p>rate = $k[\text{H}^+][\text{CH}_3\text{COCH}_3]$ ✓ (from all three pieces of evidence)</p> $k = \frac{\text{rate}}{[\text{H}^+][\text{CH}_3\text{COCH}_3]} = \frac{2.1 \times 10^{-9}}{0.02 \times 1.5 \times 10^{-3}} \checkmark$ <p>= $7.0 \times 10^{-5} \checkmark \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1} \checkmark$ accept 7×10^{-5}</p> <p>rate determining step involves species in rate equation ✓</p> <p>two steps that add up to give the overall equation ✓</p> <p>The left hand side of a step that contains the species in rate-determining step ✓ i.e., for marking points 2 and 3: $\text{CH}_3\text{COCH}_3 + \text{H}^+ \longrightarrow [\text{CH}_3\text{COHCH}_3]^+$ $[\text{CH}_3\text{COHCH}_3]^+ + \text{I}_2 \longrightarrow \text{CH}_3\text{COCH}_2\text{I} + \text{HI} + \text{H}^+$</p> <p>organises relevant information clearly and coherently, using specialist vocabulary where appropriate Use of the following four words/phrases: constant, half-life, order, doubles/x2 ✓</p>	<p>[2]</p> <p>[2]</p> <p>[2]</p> <p>[4]</p> <p>[3]</p> <p>[1]</p>
		Total: 14

