

Surname	Centre Number	Candidate Number
Other Names		2



GCE A level

1095/01



S16-1095-01

CHEMISTRY – CH5

A.M. WEDNESDAY, 22 June 2016

1 hour 45 minutes

For Examiner's use only		
Question	Maximum Mark	Mark Awarded
Section A	1.	10
	2.	12
	3.	18
Section B	4.	20
	5.	20
Total	80	

1095
010001

ADDITIONAL MATERIALS

In addition to this examination paper, you will need:

- a calculator;
- an 8 page answer book;
- a copy of the **Periodic Table** supplied by WJEC.
Refer to it for any **relative atomic masses** you require.

INSTRUCTIONS TO CANDIDATES

Use black ink or black ball-point pen.

Write your name, centre number and candidate number in the spaces at the top of this page.

Section A Answer **all** questions in the spaces provided.

Section B Answer **both** questions in **Section B** in a separate answer book which should then be placed inside this question-and-answer book.

Candidates are advised to allocate their time appropriately between **Section A (40 marks)** and **Section B (40 marks)**.

INFORMATION FOR CANDIDATES

The number of marks is given in brackets at the end of each question or part-question.

The maximum mark for this paper is 80.

Your answers must be relevant and must make full use of the information given to be awarded full marks for a question.

The *QWC* label alongside particular part-questions indicates those where the Quality of Written Communication is assessed.


SECTION A

Answer all questions in the spaces provided.

1. (a) Elen carried out an investigation into the rate of reaction between propanone and iodine in an acidic solution. This is a multi-step reaction but the overall equation for the reaction is:



- (i) In the first part of the investigation she measured how the concentration of propanone changed with time. Her results are shown in the graph below.



Explain how the graph shows that the reaction is first order with respect to propanone. Use values from the graph to justify your answer. [2]

.....

.....

.....

.....

(ii) In the second part of the investigation Elen investigated how different initial concentrations of iodine and acid affected the rate of reaction. The following results were obtained.

$[\text{CH}_3\text{COCH}_3]$ / mol dm^{-3}	$[\text{I}_2]$ / mol dm^{-3}	$[\text{H}^+]$ / mol dm^{-3}	Initial rate / $\text{mol dm}^{-3} \text{ s}^{-1}$
1.5×10^{-3}	0.030	0.020	2.1×10^{-9}
1.5×10^{-3}	0.060	0.040	4.2×10^{-9}
1.5×10^{-3}	0.030	0.040	4.2×10^{-9}

I. Determine the orders of reaction with respect to I_2 and H^+ . [2]

I_2

.....

H^+

.....

II. Write the rate equation for the reaction. [1]

.....

III. Calculate the value of the rate constant in the rate equation and state its unit. [2]

$k =$

Unit

- (b) Another multi-step reaction is the one between nitrogen dioxide and carbon monoxide. The overall equation for the reaction is:



The rate equation for this reaction is as follows.

$$\text{rate} = k[\text{NO}_2]^2$$

The first step is the rate-determining step.

- (i) Explain what is meant by the *rate-determining step*. [1]

.....
.....

- (ii) Write equations to show a possible two-step mechanism for this reaction. [2]

.....
.....

Total [10]

10

BLANK PAGE

2. Acids can be considered to be strong or weak and concentrated or dilute.

- (a) For an aqueous solution of an acid, explain the difference between the meaning of the terms *weak acid* and *dilute acid*. [2]

.....

.....

.....

.....

- (b) The grids opposite show titration curves for the addition of aqueous sodium hydroxide solution to 25.0 cm³ of aqueous acid.

From the list below, choose which acids were used to give curves **A** and **B** giving reasons for your answer.

W	0.1 mol dm ⁻³ HCl
X	0.001 mol dm ⁻³ HCl
Y	0.1 mol dm ⁻³ CH ₃ COOH
Z	0.001 mol dm ⁻³ CH ₃ COOH

(K_a for CH₃COOH = 1.8×10^{-5} mol dm⁻³)

- (i) Curve **A** [2]

.....

.....

.....


- (ii) Curve **B** [3]

.....


.....

.....

Titration curves of acid-alkali reactions



Curve A



Curve B

- (iii) State, giving a reason, which of the following indicators would be **most** suitable for titration **B**. [2]

Indicator	pH range
methyl orange	3.4 – 4.8
chlorophenol red	4.8 – 6.4
thymol blue	8.0 – 9.6
brilliant cresyl blue	10.8 – 12.0

- (iv) Calculate the concentration of the aqueous sodium hydroxide solution used in titration **A**. [2]

Concentration = mol dm⁻³

- (c) Aqueous ammonia reacts with hydrochloric acid to form the salt ammonium chloride, NH₄Cl. Give a reason why the pH value for a solution of NH₄Cl is less than 7. [1]

Total [12]

12

BLANK PAGE

3. Read the passage below and then answer the questions in the spaces provided.

Hydrogen

Hydrogen might be the simplest of all the elements in terms of atomic structure, but a look at the chemistry of hydrogen enables us to gain a better understanding of many important chemical ideas. Several chemical definitions and standards are based on hydrogen chemistry – from standard electrodes to the pH scale.

- 5 Hydrogen is the first element in the Periodic Table and is named from the Greek word *hydrogenos* which means water maker. Hydrogen is the only element that has different names for its isotopes. ${}^1_1\text{H}$ is hydrogen, ${}^2_1\text{H}$ is deuterium and ${}^3_1\text{H}$ is tritium.

Acidity is expressed using the pH scale first devised by the Swedish chemist Sorenson.


$$\text{pH} = -\log[\text{H}^+]$$

- 10 The scale usually runs from 0–14 because 1 mol dm⁻³ H⁺ (acid) has a pH of 0 and 1 mol dm⁻³ OH⁻ (alkali) has a pH of 14. An aqueous solution is neutral when the concentrations of H⁺ and OH⁻ are equal. At 25 °C, the ionic product of water, K_w , has a numerical value of 1.0×10^{-14} . Pure water has a pH of 7, and is neutral. This neutral value of pH can be calculated from K_w . Since boiling water has a larger value of K_w than water at 25 °C, it follows that a substance
15 that is dissolved in boiling water to give a solution with a pH of 7 is slightly alkaline!

When measuring electrode potentials, it is potential differences which are measured. This means that the potential of one half-cell is compared with that of another. Again, hydrogen is the basis of the comparison. All electrode potentials are compared with that of the standard hydrogen electrode.

- 20 Looking at data for elements, we see that hydrogen often has the greatest or smallest quantity. For example when burned in air, hydrogen evolves more heat per unit mass than any other substance [$\Delta H_c^\ominus(\text{H}_2) = -286 \text{ kJ mol}^{-1}$]. Rockets such as the space shuttle, use a mixture of liquid hydrogen and liquid oxygen to propel them into orbit. Cars have been developed that run on hydrogen using fuel cells. The original airships were filled with hydrogen but its flammability
25 led to a catastrophic fire on the Hindenburg in 1937. Modern airships use helium.

Most hydrogen today is used for the processing of fossil fuels and in the production of ammonia.



- Other important uses include as a hydrogenating agent in making margarines, in the production of methanol, in the manufacture of hydrochloric acid and also in cryogenics. Hydrogen – the
30 light, flammable gas with its important industrial roles – does far more than just make water!

- End of passage -

- (a) Write an expression for the ionic product of water, K_w , (*line 12*) giving its unit, if any. [1]

Unit

- (b) The value for K_w at 100 °C is 5.13×10^{-13} . Use this to explain why an aqueous solution of a salt with a pH of 7 at this temperature is slightly alkaline (*line 15*). [3]

.....
.....

- (c) All electrode potentials are compared with the standard hydrogen electrode (*lines 18-19*). With the aid of a diagram or otherwise explain what is meant by the *standard hydrogen electrode*. [2]

.....
.....
.....

- (d) (i) Use the data given to calculate the standard enthalpy change of combustion of methane. [2]

Substance	CH ₄ (g)	CO ₂ (g)	H ₂ O(l)
Standard enthalpy change of formation, $\Delta H_f^\theta / \text{kJ mol}^{-1}$	-75	-394	-286

$$\Delta H_c^\theta = \dots\dots\dots \text{kJ mol}^{-1}$$

- (ii) Use this result to show that the statement in *line 21* is correct when comparing hydrogen and methane. [2]

.....

.....

- (e) Cars have been developed that run on hydrogen using fuel cells (*lines 23-24*). Explain the principles underlying the operation of the hydrogen fuel cell. [3]

QWC [1]

.....

.....

.....

.....

.....

.....

(f) In the production of ammonia (*lines 26-27*), nitrogen and hydrogen were mixed in a vessel and allowed to reach equilibrium at a given temperature. The initial partial pressure of nitrogen was 26 atm and that of hydrogen was 82 atm. The equilibrium partial pressure of the remaining nitrogen was 18 atm.

(i) Write an expression for the equilibrium constant, K_p , for this reaction. [1]

(ii) Calculate the equilibrium partial pressures of hydrogen and ammonia and use these to calculate a value for K_p at this temperature, giving the unit if any. [3]

$K_p =$

Unit

Total [18]

18

Total Section A [40]

SECTION B

Answer **both** questions in the separate answer book provided.

4. (a) Copper is a typical transition metal.

Characteristics of these metals include an ability to:

- form coloured ions
- show variable oxidation states
- form complex ions

(i) State **one other** chemical property of transition metals. [1]

(ii) Explain why copper(I) compounds are generally white. [2]

- (b) Copper compounds take part in several different types of reaction including ligand substitution and precipitation. Using copper compounds, give an example for both types of reaction, stating any observations. Give the formula for the copper-containing product for each example. [6]

QWC [1]

- (c) Brass is an alloy of copper and zinc.


A 2.05 g brass screw was dissolved in nitric acid and the solution formed was diluted to 100 cm³ in a volumetric flask. An excess of potassium iodide solution was added to 25.0 cm³ of this solution and the iodine produced was titrated against a 0.200 mol dm⁻³ solution of sodium thiosulfate. The iodine required 24.00 cm³ of the sodium thiosulfate solution for complete reaction.

(i) Name a suitable indicator for this titration. [1]

(ii) Calculate the percentage by mass of copper in the brass. Give your answer to **three** significant figures. [4]

(The ratio of Cu²⁺:S₂O₃²⁻ is 1:1)

- (d) The diagram below shows the apparatus that was used to measure the emf of a Cu^{2+}/Cu , Fe^{2+}/Fe electrochemical cell.



Some standard electrode potentials, E^\ominus , are given below.

System	E^\ominus/V
$\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Cu}(\text{s})$	+0.34
$\text{Fe}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Fe}(\text{s})$	-0.44
$\text{Ni}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Ni}(\text{s})$	-0.25

- Name the part of the cell labelled **A** and state its purpose. [2]
- State, giving a reason, which of the electrodes will be positively charged in the above cell. [1]
- Calculate the standard emf, in volts, for the above cell. [1]
- State whether or not you would expect nickel to react with iron(II) ions. Give a reason for your answer. [1]

Total [20]


5. (a) Group II elements can only show an oxidation state of II, however Group IV elements can show oxidation states of II and IV in their compounds.

- (i) State how the relative stability of these oxidation states changes as Group IV is descended and give a reason for this trend. [2]
- (ii) The characteristics of the Group IV elements and their compounds change significantly from carbon to lead. Show how this statement is true by comparing:
- the reactions, if any, of carbon dioxide and lead(II) oxide with acids and alkalis
 - the reduction-oxidation properties of carbon monoxide and lead(IV) oxide.

Your answer should include any relevant chemical equations.

[6]
QWC [1]

(b) Endothermic solid-solid reactions are rare in chemistry, but some do occur spontaneously. One such example is the reaction between barium hydroxide and ammonium chloride. The reaction can be represented as follows.



The entropy values of the compounds involved in this reaction are given below.

Compound	Ba(OH) ₂ ·8H ₂ O(s)	NH ₄ Cl(s)	NH ₃ (g)	H ₂ O(l)	BaCl ₂ ·2H ₂ O(s)
Entropy / J K ⁻¹ mol ⁻¹	427	95	192	70	203

- (i) Explain why there is an increase in entropy for this reaction. [1]
- (ii) Calculate the entropy change for this reaction. [1]
- (iii) Calculate the free energy change, ΔG , for the reaction at 25 °C and explain why this reaction is feasible. [3]

- (c) The enthalpy change of formation of barium chloride, BaCl_2 , can be determined indirectly using a Born-Haber cycle.

Use the data given below to calculate the enthalpy change of formation of barium chloride in kJ mol^{-1} . [4]

Process	$\Delta H^\theta / \text{kJ mol}^{-1}$
$\text{Ba(s)} \longrightarrow \text{Ba(g)}$	176
$\frac{1}{2}\text{Cl}_2(\text{g}) \longrightarrow \text{Cl(g)}$	121
$\text{Ba(g)} \longrightarrow \text{Ba}^+(\text{g}) + \text{e}^-$	502
$\text{Ba}^+(\text{g}) \longrightarrow \text{Ba}^{2+}(\text{g}) + \text{e}^-$	966
$\text{Cl(g)} + \text{e}^- \longrightarrow \text{Cl}^-(\text{g})$	-364
$\text{Ba}^{2+}(\text{g}) + 2\text{Cl}^-(\text{g}) \longrightarrow \text{BaCl}_2(\text{s})$	-2018

- (d) Write the **formulae** of the chlorine-containing species that are produced when chlorine reacts with warm aqueous sodium hydroxide. [2]

Total [20]

Total Section B [40]

END OF PAPER

BLANK PAGE

BLANK PAGE



GCE A level

1095/01-A



S16-1095-01A

**CHEMISTRY – PERIODIC TABLE
FOR USE WITH CH5**

A.M. WEDNESDAY, 22 June 2016

THE PERIODIC TABLE

Group 1 2 3 4 5 6 7 0

Period 1 2 3 4 5 6 7

Period	1	2	p Block						0
1	1.01 H Hydrogen 1							4.00 He Helium 2	
2	6.94 Li Lithium 3	9.01 Be Beryllium 4						20.2 Ne Neon 10	
3	23.0 Na Sodium 11	24.3 Mg Magnesium 12						35.5 Cl Chlorine 17	
4	39.1 K Potassium 19	40.1 Ca Calcium 20						79.9 Br Bromine 35	
5	85.5 Rb Rubidium 37	87.6 Sr Strontium 38						127 I Iodine 53	
6	133 Cs Caesium 55	137 Ba Barium 56						(222) Rn Radon 86	
7	(223) Fr Francium 87	(226) Ra Radium 88						(210) At Astatine 85	

Key	
Ar	relative atomic mass
Symbol	atomic number
Name	Z

d Block					
10.8 B Boron 5	12.0 C Carbon 6	14.0 N Nitrogen 7	16.0 O Oxygen 8	19.0 F Fluorine 9	20.2 Ne Neon 10
27.0 Al Aluminium 13	28.1 Si Silicon 14	31.0 P Phosphorus 15	32.1 S Sulfur 16	35.5 Cl Chlorine 17	40.0 Ar Argon 18
69.7 Ga Gallium 31	72.6 Ge Germanium 32	74.9 As Arsenic 33	79.0 Se Selenium 34	79.9 Br Bromine 35	83.8 Kr Krypton 36
65.4 Zn Zinc 30	63.5 Cu Copper 29	58.7 Ni Nickel 28	58.9 Co Cobalt 27	55.8 Fe Iron 26	54.9 Mn Manganese 25
112 Cd Cadmium 48	108 Ag Silver 47	106 Pd Palladium 46	103 Rh Rhodium 45	101 Ru Ruthenium 44	101 Tc Technetium 43
201 Hg Mercury 80	197 Au Gold 79	195 Pt Platinum 78	192 Ir Iridium 77	190 Os Osmium 76	186 Re Rhenium 75
204 Tl Thallium 81	207 Pb Lead 82	209 Bi Bismuth 83	209 Po Polonium 84	210 At Astatine 85	210 Rn Radon 86

f Block											
140 Ce Cerium 58	141 Pr Praseodymium 59	144 Nd Neodymium 60	150 Sm Samarium 62	(153) Eu Europium 63	157 Gd Gadolinium 64	163 Dy Dysprosium 66	165 Ho Holmium 67	167 Er Erbium 68	169 Tm Thulium 69	173 Yb Ytterbium 70	175 Lu Lutetium 71
232 Th Thorium 90	(231) Pa Protactinium 91	238 U Uranium 92	(242) Pu Plutonium 94	(243) Am Americium 95	(247) Cm Curium 96	(251) Cf Californium 98	(254) Es Einsteinium 99	(253) Fm Fermium 100	(256) Md Mendelevium 101	(254) No Nobelium 102	(257) Lr Lawrencium 103