

CAMBRIDGE INTERNATIONAL EXAMINATIONS

GCE Advanced Level

www.XtremePapers.com

MARK SCHEME for the October/November 2013 series

9336 FOOD STUDIES

9336/01

Paper 1 (Theory), maximum raw mark 100

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2013 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

Page 2	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2013	9336	01

1 (a) Anaemia

Causes

Iron deficiency anaemia - lack of iron-providing foods -
 lack of vitamin C -
 blood loss - e.g. surgery / childbirth / menstruation etc. -
 destruction of red blood cells -
 iron needed for formation of haemoglobin - in red blood cells -
 to transport oxygen - to oxidise glucose in cells - to release energy
 low haemoglobin - less oxygen - less energy released -

Megaloblastic anaemia - lack of folate and vitamin B12 -

Impaired DNA synthesis – prevents cell division
 Fewer / larger - irregular-shaped red blood cells -
 so less oxygen carried

Pernicious anaemia

lack of intrinsic factor in gut needed for absorption of vitamin B12 -
 B12 not absorbed

Symptoms

pallor - tiredness - dizziness - headaches etc.

Remedies

sources of iron - e.g. red meat / dark green vegetables / eggs /
 fortified cereals / dark chocolate etc.

sources of vitamin C - e.g. citrus fruit / blackcurrants /
 green vegetables / peppers etc.

sources of folate - e.g. green vegetables / liver / yeast extract

sources of vitamin B12 - e.g. meat / milk

Max of 1 food example for each correct nutrient

Do not accept supplements

10 points covering all areas

2 points = 1 mark

[5]

(b) Osteoporosis

Causes

loss of all bone components - especially calcium - in old people over 50 -
 particularly women - after menopause - body draws calcium from the bones -
 linked to reduced productions of oestrogen -
 not caused by a lack of calcium - or vitamin D

Symptoms

loss of bone density - shrinkage in height - aches in bones -
 bones become porous - fracture more readily

Remedies

Phosphorous – in all foods

plenty of dietary calcium - from milk / cheese / green vegetables / fortified breakfast
 cereals / white bread etc.

vitamin D for calcium absorption - milk / cheese / oily fish /
 fish liver oil / margarine / butter / sunshine etc.

regular exercise - as preventive measures -

adequate supply of calcium at younger age -

to ensure bones are strong

Accept 1 food example for Vitamin D and sunshine

10 points covering all areas

2 points = 1 mark

[5]

Page 3	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2013	9336	01

(c) Goitre

Causes

insufficient intake of iodine -
 common in areas a long way from the sea -
 where soil contains little iodine -
 needed by the thyroid gland - for formation of thyroxin - a hormone
 involved in the regulation of the rate of oxidation of nutrients in body cells

Symptoms

enlargement of thyroid gland -

Remedies

use of iodised table salt - has potassium iodide added -
 sea fish / salt water fish - vegetables grown near the sea -
 seaweed etc.

10 points covering all areas

2 points = 1 mark

[5]

(d) Night blindness

Causes

lack of vitamin A (retinol) - needed to synthesise rhodopsin -
 visual purple - a light-sensitive pigment - in retina of the eye -
 which helps eyes to adjust to dim light -

Or (Not awarded twice)

Symptoms

vision in dim light is impaired -

Remedies

Animal food - foods rich in vitamin A include milk / cheese / butter / egg yolk /
 fish liver oil / margarine etc.

plants - contain beta-carotene -

which can be converted into vitamin A in the body -

but is only one sixth as valuable as consuming vitamin A -

found in orangey-red fruits and vegetables -

e.g. carrots / papaya / red peppers

Max 1 food for retinol

Max 1 food for beta carotene

10 points covering both areas

2 points = 1 mark

[5]

(e) Scurvy

Causes

vitamin C (Ascorbic acid) deficiency - needed for absorption of iron

formation of haemoglobin - transport oxygen from lungs to cells -

to oxidise glucose - and release energy -

needed for the formation of collagen -

main protein of connective tissue - protects organs -

Page 4	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2013	9336	01

Symptoms

spontaneous bruising - as small blood vessels break - haemorrhaging under skin - gums become black - and spongy /swollen gums
teeth become loose - wounds heal slowly - fractures slow to heal -
caused by failure to form connective tissue -
may also have anaemia - failure to absorb iron -
inability to form red blood cells - change in bone structure

Remedies

regular intake important since vitamin C cannot be stored in the body -
e.g. blackcurrants, citrus fruit, rose hips, melon, strawberries, green vegetables, green peppers, new potatoes
take care to avoid loss of vitamin C when preparing, cooking and serving green vegetables -
water soluble - destroyed by heat -
oxidises etc. - raw fruit and vegetables contain most vitamin C -
some foods are fortified with vitamin C - some fruit juices etc. -
read packaging to compare with RDI etc.

Max 1 food example

10 points covering all areas 2 points = 1 mark [5]

2 (a) Digestion of food in the small intestine

Duodenum

Protein

trypsinogen - from pancreatic juice - is activated by enterokinase - in intestinal juice - to become trypsin -

trypsin converts protein into peptides

Fat

bile - made in liver - stored in the gall bladder - emulsifies fat/ breaks into small droplets - increases surface area

lipase - from pancreatic juice - converts fat to fatty acids - and glycerol

Carbohydrate

amylase - from pancreatic juice - converts starch to maltose -

Ileum

Protein

erepsin - from intestinal juice - converts peptides to amino acids -

Fat

lipase - from intestinal juice - completes breakdown of fat into fatty acids and glycerol

Carbohydrate

intestinal juice - contains maltase - converts maltose to glucose -

lactase - converts lactose to galactose and glucose -

sucrase / invertase - converts sucrose to glucose and fructose -

24 points 2 points = 1 mark [12]

Page 5	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2013	9336	01

(b) Absorption of nutrients in the ileum

Protein

amino acids absorbed through microvilli - by active transport - into blood capillaries - of villi - then to liver - via hepatic portal vein

Fat

glycerol and fatty acids - reform into fats - in walls of villi - pass into lacteal - then into lymphatic system - by diffusion - and active transport - now called chylomicrons

Carbohydrate

villi - finger-like projections - increase surface area of ileum -

(can show on a diagram) - well supplied with blood capillaries -

monosaccharides (glucose, galactose, fructose) absorbed directly into bloodstream - pass to liver - via hepatic portal vein

Minerals

only about one third of **calcium** is absorbed -

remainder is lost from the body in faeces - vitamin D aids absorption otherwise little would be absorbed - lactose increases absorption -

phytic acid - and oxalic acid - react with calcium -

preventing absorption

only about 5%-20% **iron** is absorbed -

depends on the body's need for iron -

depends on the form of iron - haem iron - ferrous -

more easily absorbed - non-haem iron - ferric -

can be reduced to ferrous - by vitamin C - so aids iron absorption -

phytic acid - oxalic acid - and tannin - reduce iron absorption

Vitamins

vitamin A and vitamin D are fat soluble - absorbed with fat

N.B. Avoid crediting some information more than once e.g.

villi - liver - hepatic portal vein etc.

26 points

2 points = 1 mark

[13]

Page 6	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2013	9336	01

3 (a) Characteristics of fats and oils and uses in preparation of dishes

Characteristics

fats are esters - result of reaction between acids and alcohol -
 3 hydroxyl groups of the glycerol molecule can each combine with a fatty acid molecule to give a triglyceride - a type of ester -
 each molecule could have more than one type of fatty acid - glycerol common to all
 difference between fats and oils due to different fatty acids in composition -
 saturated fatty acids - all carbon molecules saturated with hydrogen - hard -
 all c-c bonds within molecule are single -
 monounsaturated - one double bond between adjacent carbon atoms -
 can take up more hydrogen - soft fat -
 polyunsaturated - more than one double bond between adjacent carbon atoms
 can take up more hydrogen -
 saturated fats usually from animals - polyunsaturated usually from plants
 fish oils are unsaturated -
 double bonds are a point of weakness -
 can readily be broken by a range of substances -
 Max 4 for correct diagrams
cis fatty acids - hydrogen atoms on same side of double bond -
trans fatty acids -
hydrogen atoms diagonally opposite each other at double bond - DO NOT CREDIT in (a) and (b)

(could give some of this information in a diagram)

fats melt over a range of temperatures - depending on fatty acids -
 as temperature rises some fatty acids will melt - others remain solid -
 this allows solid triglycerides to move within the fat - fat is spreadable -
 softer fats spread more readily - shape changes with pressure -
 plasticity of fat affects creaming and shortening properties -
 pure fats such as lard have more shortening power than butter or margarine -
 because they contain water -
 creaming power of a fat is measured by its capacity to incorporate air bubbles
 a fat with a wide plastic range is best for creaming -
 hydrogenated vegetable fats are better than butter -
 butter has a narrow plastic range -

fats and oils have different smoke points - high smoke point for frying -
otherwise food will absorb fat or oil -
different fatty acids have different flavours - affect acceptability -
oils hydrogenated to form fats - process can be halted at varying stages -
to give desired hardness for intended use etc.
last points should not be credited in section (a) and (b)

Uses

spreading - flavouring - forming an emulsion - shortening - aeration - frying - adding colour -
 improving keeping quality - emulsifies with moisture - to retain moisture - e.g. in rich cakes
 1 use + 1 fat example = 1 mark
 increasing calorific value without adding bulk - e.g. fried foods etc.
 sealing - melted butter on pate sets on cooling - prevents surface drying -
 lubrication - prevents food sticking to cake tins etc. -
 makes food easier to eat - e.g. butter on bread compared with dry bread etc.

Max 5 for characteristics

Max 5 for uses of fat

20 points

2 points = 1 mark

[10]

Page 9	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2013	9336	01

(b) Globular and fibrous proteins

Globular

are bonded or compact shaped molecules - round ball -
albumins - e.g. ovalbumin (egg white), lactalbumin, blood plasma
soluble in water - and dilute salt solutions - coagulated by heat
globulins - e.g. lactoglobulin (milk, egg white, muscle cells, blood plasma)
insoluble in water - dissolve in dilute solutions of electrolytes -
coagulate on heating- correct ref. to tertiary and quaternary structure

Fibrous

fibrous molecules - many twisted in helices - can give elasticity or strength
others in form of pleated sheets
insoluble in water - and dilute solutions of salts
collagen - inelastic - high tensile strength -
connective tissue - e.g. meat, fish
converted to gelatine - by prolonged contact with boiling water -
gelatine soluble in water - more susceptible to enzymic digestion
elastin - present in elastic tissues - e.g. artery walls (meat), skin, tendons
unchanged by heat- correct ref. to tertiary structure

14 points to include named types of protein
named examples of types identified

structure of globular and fibrous protein
reference to solubility, coagulation etc.

2 points = 1 mark
Diagrams – max 2

[7]

(c) Effects of heat, agitation and acids

Heat - denaturation begins at 40 °C
coagulation - begins at approx. 60 °C -
setting - e.g. custards
coating - e.g. fish
binding - rissoles
setting of gluten - e.g. bread - to hold risen structure of baked goods
Lactalbumin forms skin on milk

Agitation - foam formation - unfolding of globular proteins -
to entangle air - causes partial coagulation of protein -
whisking of egg white - e.g. meringue
whisking of egg and sugar to trap air - e.g. sponge cakes

Acid - inactivation of enzymes / denaturation - above optimum pH
marinade meat in lemon juice / vinegar / wine - soften protein
food preservation - e.g. pickled herring
bacteria in milk ferment lactose into lactic acid - pH lowered -
milk protein coagulates - forms a curd / curdles
Must discuss all 3 for full marks
16 points - at least 3 points from each area
2 points = 1 mark

[8]

Page 11	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2013	9336	01

Advertising and packaging

- choice can be affected by how foods are shown to consumers -
- television / newspaper adverts - displays in stores - taste-testing -
- some advertisements appeal to children - sweets / McDonald's -
- children more easily persuaded than adults - sweets at till -
- tempted by free gifts / money off coupons / special offers -
- to introduce new foods - attracted by colours of packaging - endorsements - health claims
- peer pressure -
- advertising convinces people that they 'should' buy certain foods -
- packaging gives information - what is in package - some may be poor readers - serving suggestions -
- make food look appealing - may give nutritional information -
- manufacturers consider their role as educators -
- some people choose foods with the least packaging - environmental issues -
- waste of resources - others find packaged foods easier to store -
- boxes easy to stack - saves time - can be stored as soon as food is brought home protects food from damage - contamination - tampering

Physiological and psychological attributes

- depends how hungry the individual is - time available -
- appearance - flavour - aroma and texture of food -
- likes and dislikes - importance for appreciation of food -
- will repeat if experience is favourable - individual perceptions of food -
- occasion - entertaining - creating an impression -
- comfort food - relief from stress, boredom etc. -
- easy to eat too much in some circumstances - snack foods etc. -

Nutritional knowledge and skill

- food choice may be affected by its nutritional value -
- knowledge from school /home - may depend on level of education -
- differs between countries - nutrition may not be taught to everyone in school
- publicity campaigns in media - to increase awareness - and knowledge -
- need to know the functions - and dangers of food - sources of nutrients -
- in order to choose wisely - well-publicised dangers of excess fat / sugar / salt
- danger of CHD / obesity / hypertension - greater in affluent countries -
- cheaper sources of HBV protein are just as valuable in diet -
- milk / cheese / eggs -
- cheaper cuts of meat can be tenderised - complementary proteins used -
- cereals / pulses / nuts in same meal - improve quality of HBV protein -
- skill may depend on teaching in school - or at home -
- younger people may lack skill -
- less time spent at home - paid work outside home -
- more use of processed food - reduces preparation and cooking skills -
- may avoid certain meat / fish / fruit etc. -
- may buy convenience foods -
- e.g. puff pastry
- or may be skilled at limited dishes - choose foods to make those -
- lack variety - may cook as a pastime - cake decoration etc. -
- learn how to choose accordingly - expect success - to avoid waste -
- if dishes do not turn out well, will not repeat -
- more costly to buy ready-made food - but know the result to expect -
- to pay for reliability -
- e.g. choux pastry etc.

Page 12	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2013	9336	01

Cost

depends on money available to spend on food - need to budget -
 poorer people must spend a higher proportion of income on food -
 staple foods cheap - filling - can be basis of meals - local recipes etc. -
 may be a status symbol for rich people - spend more than they can afford -
 e.g. smoked salmon / caviar -
 cost is not related to nutritional value - may depend on supply available -
 or demand for the food - *cheaper foods can have high nutritive value* -
 e.g. *milk / cheese / eggs* - *can choose cheaper sources of HBV* - **DO NOT CREDIT IF CREDITED ELSEWHERE**
 pulses / cereals etc. - protein complementation - grow own produce -
 keep chickens for family consumption - use food in season - special offers -
 buy locally - reduce transport cost - foods in season are usually cheaper -
 make rather than buy ready-made food - careful use of convenience foods -
 nutritious diet may need not be expensive - reduce purchase of junk food -
 judge amount required - to avoid waste - use left-over foods -
 poor people may receive government help - free school meals / food aid
 poverty limits choice of food -
 those with unlimited income may make poor choices -
 convenience foods often contain a lot of fat -
 high sugar - high salt - risk of obesity - CHD - hypertension -
 low in NSP - etc.

Half marks for 1-2 sections. Must discuss all sections for full marks

30 points

2 points = 1 mark (credit same point only once)

[15]

(b) Recommended Daily Intake (RDI)

RDI is the estimated intake of nutrients to achieve healthy living. 1 whole mark for this statement

The RDI varies between individuals according to their age and activity level.

It represents an approximate amount of each nutrient required each day.

Can help individuals to assess whether the food eaten is adequate for their daily need. They can compare the nutritional value of the food eaten with the nutritional requirement. Make definition more explicit

Allows individuals to plan meals so that they do not exceed their RDI -

Especially important to avoid deficiency diseases -

or consuming more energy-giving food than the body can use.

RDI is only a guide - can see how diet could be improved to supply RDI s etc.

Definition of RDI

1 mark

+ 4 points

2 points = 1 mark

[3]

(c) Importance of nutritional information on packaging

allows individuals to monitor intake of nutrients - compare with RDI

may be on a special diet - diabetic - CHD - hypertension etc.

need to avoid / have a reduced intake of fat / saturated fat / salt / sugar etc.

can compare the nutritional value of different products and make choices -

can learn that different foods and be used together to compensate for high / low levels of particular nutrients to give an average intake -

may be controlling calorie intake - can calculate the total value of food eaten

given per 100g - for easy comparison between products - and per product -

so that calories consumed can be calculated etc.

14 points

2 points = 1 mark

[7]

Page 13	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2013	9336	01

6 (a) Reasons for including sauces in meals

to provide a contrasting flavour - cheese sauce with cauliflower etc.
to enhance the flavour - mint sauce with lamb etc.
to provide a contrasting texture - parsley sauce with fish etc.
to bind ingredients together - fish cakes / potato croquettes etc.
to add colour - jam sauce with steamed pudding etc.
to add nutrients - custard etc.
to reduce richness - orange sauce with duck / apple sauce with pork
to add interest or variety - chocolate sauce with ice cream etc.
to add moisture - apple pie and custard etc.
part of the dish - lasagne / macaroni cheese etc.

6 reasons + 6 examples

12 points

2 points = 1 mark

[6]

(b) (i) Gelatinisation

suspension of starch in water - separates grains -
when heated with water -
water penetrates outer layers of starch granule - softens - swells -
at 60°C to 80°C - to 5 x original size - mixture becomes viscous -
at 80°C - starch grains break up - granules dispersed throughout water - long chains
of molecules begin to unfold -
mixture thickens - forms a sol - water enclosed in meshwork - forms a gel - when
cooled - thickness depends on type of starch - and proportion used - and
temperature of liquid -
and the effect of other ingredients e.g. sugar etc.

e.g. boiled rice, roux sauce, arrowroot glaze, blancmange, custard etc.

10 points (to include at least one example)

2 points = 1 mark

[5]

(ii) Emulsification

mixture of oil and water - which does not separate -
can be oil-in-water -
e.g. milk, cream, mayonnaise, ice cream, gravy etc.
or water-in-oil -
e.g. butter, margarine, egg yolk etc. -
two liquids which do not normally mix - need an emulsifying agent -
e.g. lecithin in egg yolk - has hydrophobic group - attracted to oil -
and hydrophilic group - attracted to water -
oil molecules are suspended in water -
small amounts of oil are added at first - e.g. in mayonnaise -
to ensure thorough emulsification - more oil causes thickening -
e.g. creamed cake mixture - liquid egg added to creamed fat and sugar
do not separate / curdle if added gradually -
lecithin is emulsifying agent -
(may show in a diagram)

10 points (to include at least one example)

2 points = 1 mark

[5]

Page 14	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2013	9336	01

(iii) Coagulation

when heated - protein molecules change structure - denatured -
 primary structure unchanged - secondary structure altered -
 cross-linkages break - these maintain shape of molecule -
 irreversible - molecule unfolds - sequence of amino acids remains the same
 loses elasticity - hardens / sets - properties of protein alter - less soluble -
 more viscous - unfolded molecules bond with each other - form clumps –
 causing protein to harden - egg albumin coagulates at 60 °C - yolk at 70°C -
 muscle fibres (in meat) shrink - easier to digest by digestive enzymes -
 occurs more readily with addition of acid -
 e.g. vinegar in water when boiling eggs -
 milk protein coagulates with lactic acid when turning sour - forms a curd -
 salt added to cooking water when boiling eggs - sets albumen which leaks if shell is
 cracked -
 enzyme - rennin - coagulates milk - in young children -
 rennet used commercially - to make junket - and in cheese-making -
 mechanical action / whisking - partially coagulates protein -
 molecules unfold - form a network around air - stabilising foam -
 e.g. meringue, soufflé etc. –
 e.g. of coagulation include boiled egg, egg custard, egg for coating,
 skin on boiled milk etc.
 10 points (to include at least one example)
 2 points = 1 mark

[5]

(c) Syneresis

loss of water - form a gel - as it shrinks -
 scrambled egg / baked egg custard are gels - protein shrinks if overcooked - squeezes
 out liquid which has been enclosed / syneresis occurs -
 reverse of gelatinisation in a starch sauce - retrogradation -
 starches which are rich in amylopectin are resistant to retrogradation -
 water can slowly weep out of a pectin gel during storage -
 could be due to the presence of too much acid
 8 points
 2 points = 1 mark

[4]

Page 16	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2013	9336	01

(c) (i) Principles of deep freezing

water turns to ice - below 0 °C - cell contents impure -
lowers freezing temperature - frozen by -10 °C -
micro-organisms dormant - water unavailable for bacterial growth -
some bacteria destroyed - enzyme action slowed down -
store at -18 °C - enzymes which attack fat are still active - blanching inactivates
enzymes - reduces bacteria by 90% -
must be quick frozen - to ensure small ice crystals - at -25 °C -
food passes through zone of maximum ice crystal formation -
in shortest time possible - 30 minutes or less -
does not damage cell walls - large crystals rupture cell walls -
liquid is lost on thawing - drips out - flavour lost -
texture / appearance spoilt -
make sure temperature of freezer is at -25 °C before food is put in -
freeze small amounts at a time -
to avoid raising temperature inside freezer etc.

(ii) Advice on use of freezer

use air-tight packaging materials / waterproof / completely sealed -
to prevent evaporation of liquids - 'freezer burn' on meat etc. -
dries out surface - cannot be reversed - leave a head space for liquids
expand on freezing - remove air - suck out with a straw etc.
air circulation causes moisture loss - label - on dry surface / name / amount / date of
freezing - use by date / storage time - refer to textbooks for recommended time - e.g.
polythene bags / wax cartons / plastic tubs - separate layers with plastic film etc. - use
in rotation - pack in useable amounts - keep freezer full -
for maximum efficiency -
food packed together maintains low temperature more efficiently -
defrost every 6mths / do not allow the ice to build up
10 points - at least 4 points from each area
2 points = 1 mark
Even balance on **(c) (i)** and **(c) (ii)** for full marks

[5]

Page 17	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2013	9336	01

8 (a) Nutritive value of milk

HBV - protein - casein, lactalbumin, lactoglobulin - all IAAs -
 growth - repair - maintenance - enzymes - hormones etc.
 fat - finely divided - tiny droplets in oil in water emulsion -
 easily digested - warmth - concentrated source of energy etc. – forms cell membrane
 carbohydrate - lactose - energy
 calcium - growth of bones and teeth - clotting of blood -
 nerve function - muscle function
 phosphorus - growth of bones and teeth - found in cells -
 release of energy - DNA etc.
 vitamin A - depends on season -
 more in summer milk when cows are fed outside -
 formation of visual purple - perception of light - healthy skin mucous membranes etc.
 vitamin D - absorption of calcium
 riboflavin - release of energy from glucose -
 forms part of enzyme system
 nicotinic acid - oxidation of glucose - release of energy -
 forms part of enzyme system
 thiamine - oxidation of nutrients to release energy -
 acts as co-enzyme in process
 24 points
 2 points = 1 mark

[12]

(b) Problems associated with milk only diet

no vitamin C - required for absorption of iron - formation of collagen connective tissue -
 resistance to infection -
 formation of blood
 no iron - babies born with a 6 month supply -
 but need dietary supply - to prevent anaemia - fatigue etc.
 poor source of vitamin D - absorption of calcium - bones and teeth
 poor source of vitamin A - anti-infection -
 skin and mucous membranes
 little carbohydrate - only lactose - baby needs energy for growth -
 and movement
 high percentage of water -
 would need to consume large amount to gain sufficient nutrients
 no NSP - needed to encourage chewing -
 development of digestive system etc. as child grows
 marasmus - too little food for children under 1 year -
 thin bodies - weak - eventually death
 kwashiorkor - insufficient protein - lack of protein causes anaemia -
 protein needed for formation of blood -
 retarded growth - muscle wasting - enlarged liver -
 fluid retention etc.
 12 points
 2 points = 1 mark

[6]

(c) Advantages and disadvantages of breast feeding and bottle feeding

Breast feeding

Advantages

nutrients in correct proportion - readily absorbed - free -
 correct temperature - antibodies present in milk - immune system
 protect baby against disease - clean/safe/instant -

Page 18	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2013	9336	01

cannot be prepared incorrectly - does not cause allergies -
bond between mother and child – possible link to increased intelligence

Disadvantages

mother's diet may affect child - child may have intolerances -
tiring for mother - no-one else can help unless milk expressed and stored -
do not know how much baby has taken - some feel uncomfortable -
may be intolerance in public places etc. – cracked nipples

Bottle feeding

Advantages

can monitor the amount of milk taken by baby - others can help –
allows other members of family to spend time with child -
can feed in public without embarrassment

Disadvantages

equipment needs to be sterilised - danger of bacteria being transferred -
cost of bottles, formula milk etc. - people may not read instructions carefully - may add too
little powder / too much water /baby does not get correct amount of food - water may be
contaminated -
if left-over milk kept until later bacteria may thrive - perfect conditions -
some mothers may be influenced by advertising of formula milk -
unnecessary cost to family etc.

14 points - at least 2 points from each area (*ensure that same knowledge is not credited in both sections i.e. do not accept argument and converse as two points*)

2 points = 1 mark

[7]