

General Certificate of Education
January 2005
Advanced Subsidiary Examination



MATHEMATICS
Unit Mechanics 1A

MM1A/W

Monday 31 January 2005 Morning Session

In addition to this paper you will require:

- an 8-page answer book;
 - the **blue** AQA booklet of formulae and statistical tables.
- You may use a graphics calculator.

Time allowed: 1 hour 15 minutes

Instructions

- Use blue or black ink or ball-point pen. Pencil should only be used for drawing.
- Write the information required on the front of your answer book. The *Examining Body* for this paper is AQA. The *Paper Reference* is MM1A/W.
- Answer **all** questions.
- All necessary working should be shown; otherwise marks for method may be lost.
- The final answer to questions requiring the use of calculators should be given to three significant figures, unless stated otherwise.
- Take $g = 9.8 \text{ m s}^{-2}$, unless stated otherwise.

Information

- The maximum mark for this paper is 60.
- Mark allocations are shown in brackets.
- Unit Mechanics 1A has a **written paper and coursework**.

Advice

- Unless stated otherwise, formulae may be quoted, without proof, from the booklet.

Answer **all** questions.

1 A train travels along a straight horizontal track. It is travelling at a speed of 12 m s^{-1} when it begins to accelerate uniformly. It reaches a speed of 40 m s^{-1} after accelerating for 100 seconds.

(a) (i) Show that the acceleration of the train is 0.28 m s^{-2} . (2 marks)

(ii) Find the distance that the train travelled in the 100 seconds. (2 marks)


(b) The mass of the train is 200 tonnes and a resistance force of 40 000 N acts on the train. Find the magnitude of the driving force produced by the engine that acts on the train as it accelerates. (3 marks)

2 A particle, A , of mass 12 kg is moving on a smooth horizontal surface with velocity $\begin{bmatrix} 4 \\ 7 \end{bmatrix} \text{ m s}^{-1}$. It then collides and coalesces with a second particle, B , of mass 4 kg.

(a) If before the collision the velocity of B was $\begin{bmatrix} 2 \\ 3 \end{bmatrix} \text{ m s}^{-1}$, find the velocity of the combined particle after the collision. (4 marks)

(b) If after the collision the velocity of the combined particle is $\begin{bmatrix} 1 \\ 4 \end{bmatrix} \text{ m s}^{-1}$, find the velocity of B before the collision. (3 marks)

3 The diagram shows a rope that is attached to a box of mass 25 kg, which is being pulled along rough horizontal ground. The rope is at an angle of 30° to the ground. The tension in the rope is 40 N. The box accelerates at 0.1 m s^{-2} .




(a) Draw a diagram to show all of the forces acting on the box. (1 mark)

(b) Show that the magnitude of the friction force acting on the box is 32.1 N, correct to three significant figures. (3 marks)

(c) Show that the magnitude of the normal reaction force that the ground exerts on the box is 225 N. (3 marks)

(d) Find the coefficient of friction between the box and the ground. (2 marks)

- 4 The diagram shows two particles, A and B , that are connected by a taut light inextensible string. Particle A has mass 8 kg and is on a rough horizontal surface. Particle B has mass 7 kg and is attached to the other end of the string, as shown in the diagram. The string passes over a smooth light pulley. Particle A moves towards the pulley as particle B falls.




The coefficient of friction between A and the horizontal surface is 0.8 .

- (a) Find the magnitude of the friction force between A and the surface. *(2 marks)*
- (b) By forming an equation of motion for each particle, show that the magnitude of the acceleration of each particle is 0.392 m s^{-2} . *(5 marks)*
- 5 A golf ball is struck so that its initial velocity is 30 m s^{-1} at an angle of 60° above the horizontal. Assume that the ground is horizontal.
- (a) (i) Show that the time of flight of the particle is 5.30 seconds, correct to three significant figures. *(3 marks)*
- (ii) Find the distance of the ball from its initial position when it hits the ground for the first time. *(2 marks)*
- (b) In fact, the ball hits a tree at a point which is at a height of 5 metres above the ground. Given that the ball is descending when it hits the tree, calculate the distance of the tree from the point where the ball was struck. *(6 marks)*

TURN OVER FOR THE NEXT QUESTION

Turn over ►

- 6 A motor boat can travel at a speed of 6 m s^{-1} relative to the water. It is used to cross a river in which the current flows at 2 m s^{-1} . The resultant velocity of the boat makes an angle of 60° to the river bank, as shown in the diagram.



The angle between the direction in which the boat is travelling relative to the water and the resultant velocity is α .

- (a) Show that $\alpha = 16.8^\circ$, correct to three significant figures. (4 marks)
- (b) Find the magnitude of the resultant velocity. (3 marks)
- 7 The unit vectors \mathbf{i} and \mathbf{j} are directed east and north respectively. A yacht moves with a constant acceleration. At time t seconds the position vector of the yacht is \mathbf{r} metres. When $t = 0$ the velocity of the yacht is $(2\mathbf{i} - \mathbf{j}) \text{ m s}^{-1}$, and when $t = 10$ the velocity of the yacht is $(-\mathbf{i} + \mathbf{j}) \text{ m s}^{-1}$.
- (a) Find the acceleration of the yacht. (3 marks)
- (b) When $t = 0$ the yacht is 20 metres due east of the origin. Find an expression for \mathbf{r} in terms of t . (3 marks)
- (c) (i) Show that when $t = 20$ the yacht is due north of the origin. (2 marks)
- (ii) Find the speed of the yacht when $t = 20$. (4 marks)

END OF QUESTIONS