

General Certificate of Education  
June 2006  
Advanced Subsidiary Examination



**MATHEMATICS**  
**Unit Further Pure 1**

**MFP1**

Monday 12 June 2006 1.30 pm to 3.00 pm

**For this paper you must have:**

- an 8-page answer book
- the **blue** AQA booklet of formulae and statistical tables

You may use a graphics calculator.

Time allowed: 1 hour 30 minutes

**Instructions**

- Use blue or black ink or ball-point pen. Pencil should only be used for drawing.
- Write the information required on the front of your answer book. The *Examining Body* for this paper is AQA. The *Paper Reference* is MFP1.
- Answer **all** questions.
- Show all necessary working; otherwise marks for method may be lost.

**Information**

- The maximum mark for this paper is 75.
- The marks for questions are shown in brackets.

**Advice**

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.

---

Answer **all** questions.

---

1 The quadratic equation

$$3x^2 - 6x + 2 = 0$$

has roots  $\alpha$  and  $\beta$ .

(a) Write down the numerical values of  $\alpha + \beta$  and  $\alpha\beta$ . (2 marks)

(b) (i) Expand  $(\alpha + \beta)^3$ . (1 mark)

(ii) Show that  $\alpha^3 + \beta^3 = 4$ . (3 marks)

(c) Find a quadratic equation with roots  $\alpha^3$  and  $\beta^3$ , giving your answer in the form  $px^2 + qx + r = 0$ , where  $p$ ,  $q$  and  $r$  are integers. (3 marks)

2 A curve satisfies the differential equation

$$\frac{dy}{dx} = \log_{10} x$$

Starting at the point (2, 3) on the curve, use a step-by-step method with a step length of 0.2 to estimate the value of  $y$  at  $x = 2.4$ . Give your answer to three decimal places. (6 marks)

3 Show that

$$\sum_{r=1}^n (r^2 - r) = kn(n+1)(n-1)$$

where  $k$  is a rational number. (4 marks)

4 Find, in **radians**, the general solution of the equation

$$\cos 3x = \frac{\sqrt{3}}{2}$$

giving your answers in terms of  $\pi$ . (5 marks)

5 The matrix  $\mathbf{M}$  is defined by

$$\mathbf{M} = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix}$$

(a) Find the matrix:

(i)  $\mathbf{M}^2$ ; *(3 marks)*

(ii)  $\mathbf{M}^4$ . *(1 mark)*

(b) Describe fully the geometrical transformation represented by  $\mathbf{M}$ . *(2 marks)*

(c) Find the matrix  $\mathbf{M}^{2006}$ . *(3 marks)*

6 It is given that  $z = x + iy$ , where  $x$  and  $y$  are real numbers.

(a) Write down, in terms of  $x$  and  $y$ , an expression for

$$(z + i)^*$$

where  $(z + i)^*$  denotes the complex conjugate of  $(z + i)$ . *(2 marks)*

(b) Solve the equation

$$(z + i)^* = 2iz + 1$$

giving your answer in the form  $a + bi$ . *(5 marks)*

**Turn over for the next question**

**Turn over ►**

- 7 (a) Describe a geometrical transformation by which the hyperbola


$$x^2 - 4y^2 = 1$$

can be obtained from the hyperbola  $x^2 - y^2 = 1$ .

(2 marks)

- (b) The diagram shows the hyperbola  $H$  with equation

$$x^2 - y^2 - 4x + 3 = 0$$



By completing the square, describe a geometrical transformation by which the hyperbola  $H$  can be obtained from the hyperbola  $x^2 - y^2 = 1$ .

(4 marks)

- 8 (a) The function  $f$  is defined for all real values of  $x$  by

$$f(x) = x^3 + x^2 - 1$$

- (i) Express  $f(1+h) - f(1)$  in the form

$$ph + qh^2 + rh^3$$

where  $p$ ,  $q$  and  $r$  are integers.


(4 marks)

- (ii) Use your answer to part (a)(i) to find the value of  $f'(1)$ .

(2 marks)

- (b) The diagram shows the graphs of

$$y = \frac{1}{x^2} \quad \text{and} \quad y = x + 1 \quad \text{for} \quad x > 0$$



The graphs intersect at the point  $P$ .

- (i) Show that the  $x$ -coordinate of  $P$  satisfies the equation  $f(x) = 0$ , where  $f$  is the function defined in part (a). (1 mark)
- (ii) Taking  $x_1 = 1$  as a first approximation to the root of the equation  $f(x) = 0$ , use the Newton–Raphson method to find a second approximation  $x_2$  to the root. (3 marks)
- (c) The region enclosed by the curve  $y = \frac{1}{x^2}$ , the line  $x = 1$  and the  $x$ -axis is shaded on the diagram. By evaluating an improper integral, find the area of this region. (3 marks)

Turn over ►

9 A curve  $C$  has equation

$$y = \frac{(x+1)(x-3)}{x(x-2)}$$

- (a) (i) Write down the coordinates of the points where  $C$  intersects the  $x$ -axis. (2 marks)
- (ii) Write down the equations of all the asymptotes of  $C$ . (3 marks)

- (b) (i) Show that, if the line  $y = k$  intersects  $C$ , then

$$(k-1)(k-4) \geq 0 \quad (5 \text{ marks})$$

- (ii) Given that there is only one stationary point on  $C$ , find the coordinates of this stationary point.

(No credit will be given for solutions based on differentiation.) (3 marks)

- (c) Sketch the curve  $C$ . (3 marks)

**END OF QUESTIONS**

**There are no questions printed on this page**

**There are no questions printed on this page**