

General Certificate of Education  
January 2008  
Advanced Level Examination



**MATHEMATICS**  
**Unit Pure Core 3**

**MPC3**

Thursday 17 January 2008 1.30 pm to 3.00 pm

**For this paper you must have:**

- an 8-page answer book
- the blue AQA booklet of formulae and statistical tables.

You may use a graphics calculator.

Time allowed: 1 hour 30 minutes

**Instructions**

- Use blue or black ink or ball-point pen. Pencil should only be used for drawing.
- Write the information required on the front of your answer book. The *Examining Body* for this paper is AQA. The *Paper Reference* is MPC3.
- Answer **all** questions.
- Show all necessary working; otherwise marks for method may be lost.

**Information**

- The maximum mark for this paper is 75.
- The marks for questions are shown in brackets.

**Advice**

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.

---

Answer **all** questions.

---

1 (a) Find  $\frac{dy}{dx}$  when:

(i)  $y = (2x^2 - 5x + 1)^{20}$ ; *(2 marks)*

(ii)  $y = x \cos x$ . *(2 marks)*

(b) Given that

$$y = \frac{x^3}{x-2}$$

show that

$$\frac{dy}{dx} = \frac{kx^2(x-3)}{(x-2)^2}$$

where  $k$  is a positive integer. *(3 marks)*

2 (a) Solve the equation  $\cot x = 2$ , giving all values of  $x$  in the interval  $0 \leq x \leq 2\pi$  in radians to two decimal places. *(2 marks)*

(b) Show that the equation  $\operatorname{cosec}^2 x = \frac{3 \cot x + 4}{2}$  can be written as

$$2 \cot^2 x - 3 \cot x - 2 = 0$$
 *(2 marks)*

(c) Solve the equation  $\operatorname{cosec}^2 x = \frac{3 \cot x + 4}{2}$ , giving all values of  $x$  in the interval  $0 \leq x \leq 2\pi$  in radians to two decimal places. *(4 marks)*

## 3 The equation

$$x + (1 + 3x)^{\frac{1}{4}} = 0$$

has a single root,  $\alpha$ .

(a) Show that  $\alpha$  lies between  $-0.33$  and  $-0.32$ . (2 marks)

(b) Show that the equation  $x + (1 + 3x)^{\frac{1}{4}} = 0$  can be rearranged into the form

$$x = \frac{1}{3}(x^4 - 1) \quad (2 \text{ marks})$$

(c) Use the iteration  $x_{n+1} = \frac{(x_n^4 - 1)}{3}$  with  $x_1 = -0.3$  to find  $x_4$ , giving your answer to three significant figures. (3 marks)

4 The functions  $f$  and  $g$  are defined with their respective domains by

$$f(x) = x^3, \quad \text{for all real values of } x$$

$$g(x) = \frac{1}{x-3}, \quad \text{for real values of } x, x \neq 3$$

(a) State the range of  $f$ . (1 mark)

(b) (i) Find  $fg(x)$ . (1 mark)

(ii) Solve the equation  $fg(x) = 64$ . (3 marks)

(c) (i) The inverse of  $g$  is  $g^{-1}$ . Find  $g^{-1}(x)$ . (3 marks)

(ii) State the range of  $g^{-1}$ . (1 mark)

5 (a) (i) Given that  $y = 2x^2 - 8x + 3$ , find  $\frac{dy}{dx}$ . (1 mark)

(ii) Hence, or otherwise, find

$$\int_4^6 \frac{x-2}{2x^2-8x+3} dx$$

giving your answer in the form  $k \ln 3$ , where  $k$  is a rational number. (4 marks)

(b) Use the substitution  $u = 3x - 1$  to find  $\int x\sqrt{3x-1} dx$ , giving your answer in terms of  $x$ . (4 marks)

**Turn over for the next question**

**Turn over ►**

- 6 (a) Sketch the curve with equation  $y = \operatorname{cosec} x$  for  $0 < x < \pi$ . (2 marks)
- (b) Use the mid-ordinate rule with four strips to find an estimate for  $\int_{0.1}^{0.5} \operatorname{cosec} x \, dx$ , giving your answer to three significant figures. (4 marks)
- 7 (a) Describe a sequence of **two** geometrical transformations that maps the graph of  $y = x^2$  onto the graph of  $y = 4x^2 - 5$ . (4 marks)
- (b) Sketch the graph of  $y = |4x^2 - 5|$ , indicating the coordinates of the point where the curve crosses the  $y$ -axis. (3 marks)
- (c) (i) Solve the equation  $|4x^2 - 5| = 4$ . (3 marks)
- (ii) Hence, or otherwise, solve the inequality  $|4x^2 - 5| \geq 4$ . (2 marks)
- 8 (a) Given that  $e^{-2x} = 3$ , find the exact value of  $x$ . (2 marks)
- (b) Use integration by parts to find  $\int x e^{-2x} \, dx$ . (4 marks)
- (c) A curve has equation  $y = e^{-2x} + 6x$ .
- (i) Find the exact values of the coordinates of the stationary point of the curve. (4 marks)
- (ii) Determine the nature of the stationary point. (2 marks)
- (iii) The region  $R$  is bounded by the curve  $y = e^{-2x} + 6x$ , the  $x$ -axis and the lines  $x = 0$  and  $x = 1$ .
- Find the volume of the solid formed when  $R$  is rotated through  $2\pi$  radians about the  $x$ -axis, giving your answer to three significant figures. (5 marks)

**END OF QUESTIONS**