



ASSESSMENT and
QUALIFICATIONS
ALLIANCE

Mark scheme January 2004

GCE

Mathematics A

Unit MAM3

Copyright © 2004 AQA and its licensors. All rights reserved.

Key to mark scheme

M	mark is for	method
m	mark is dependent on one or more M marks and is for	method
A	mark is dependent on M or m mark and is for	accuracy
B	mark is independent of M or m marks and is for	method and accuracy
E	mark is for	explanation
✓ or ft or F		follow through from previous incorrect result
CAO		correct answer only
AWFW		anything which falls within
AWRT		anything which rounds to
AG		answer given
SC		special case
OE		or equivalent
A2,1		2 or 1 (or 0) accuracy marks
- x EE		Deduct x marks for each error
NMS		No method shown
PI		Perhaps implied
c		Candidate


Abbreviations used in marking


MC - x	deducted x marks for miscopy
MR - x	deducted x marks for misread
ISW	ignored subsequent working
BOD	gave benefit of doubt
WR	work replaced by candidate

Application of mark scheme



Correct answer without working	mark as in scheme
Incorrect answer without working	zero marks unless specified otherwise


Award method and accuracy marks as appropriate to an alternative solution using a correct method or partially correct method.

Q	Solution	Marks	Total	Comments
1	 <p>(a) $h = 2\pi r = 2\pi \times 0.2$ $\approx 1.26 \text{ m}$</p> <p>(b) Change in PE of mass $= -mgh$ $= -5 \times g \times 0.4\pi$ $= -2\pi g$</p> <p>Change in KE of mass $= \frac{1}{2}mv^2$ $= \frac{1}{2} \times 5 \times (a\omega)^2$ $= \frac{1}{2} \times 5 \times (0.2)^2 \omega^2$ $= 0.1\omega^2$</p> <p>Change in KE of wheel $= \frac{1}{2}I\omega^2$ $= \frac{1}{2} \times 10 \times \omega^2$ $= 5\omega^2$</p> <p>$\therefore 5\omega^2 + 0.1\omega^2 = 2\pi g$</p> <p>$\omega^2 = \frac{2\pi g}{5.1}$</p> <p>$\Rightarrow \omega = 3.47 \text{ rad s}^{-1}$</p>	<p>M1 A1</p> <p>M1 A1</p> <p>A1</p> <p>M1AF</p> <p>A1F</p>	<p>2</p> <p>6</p> <p>8</p>	<p>Allow 0.4π</p> <p>For PE + KE of mass For either of PE/KE results</p> <p>(1 error only)</p>
		Total	8	

Q	Solution	Marks	Total	Comments
3	<div style="text-align: center;">  </div> <p>(a) Mass of elementary ring = $2\pi\rho x\delta x$ M.I. of element = $2\pi\rho x \cdot x^2 \delta x$ $= 2\pi\rho x^3 \delta x$ $\therefore 2\pi\rho \int_a^{2a} x^3 dx = 2\pi\rho \left[\frac{x^4}{4} \right]_a^{2a}$ $= \frac{2\pi\rho}{4} [16a^4 - a^4]$ $= \frac{30\pi\rho a^4}{4}$ but $M = 3\pi\rho a^2$ $\Rightarrow I = \frac{10Ma^2}{4} = \frac{5Ma^2}{2}$</p> <p>(b) \perp axes $I_z = I_x + I_y$ $\Rightarrow \frac{5Ma^2}{2} = 2I_D$ $\Rightarrow I_D = \frac{5Ma^2}{4}$</p>	M1 M1 M1 A1 M1A1 A1 M1 A1 A1	 7 3	
		Total	10	

Q	Solution	Marks	Total	Comments
4				
(a)	$X = 7 - 6 + 5 \cos \theta$ $= 1 + 5 \times \frac{4}{5}$ $= 9$ $Y = 4 + 5 \sin \theta$ $= 4 + 5 \times \frac{3}{5}$ $= 9$	M1A1 M1 A1		Correct at this stage (both X and Y correct)
	$\therefore \text{Resultant} = \sqrt{9^2 + 9^2}$ $= 12.7$	A1	5	CAO
(b)(i)	$Xd = -4 \times 4 - 6 \times 3 + 19$ $5d = -15$ $d = -3$ $\therefore \text{line cuts axis at } (0, -3)$	M1A1 A1 A1 A1		(for Xd) 1 st 2 terms RHS (+ 19)
			5	CAO [candidate may use "anticlockwise + ve" convention for full marks]
(ii)	$\text{Gradient line of action } + \frac{Y}{X} = \frac{12}{5}$ $\therefore y = \frac{12}{5}x - 3$ <p>(or any acceptable equivalent e.g $5y = 12x - 15$ etc)</p>	M1A1F } A1F }		}ft from (a)
			3	
	Total		13	

Q	Solution	Marks	Total	Comments
5	 <p data-bbox="145 622 480 656">(a) Block sliding $\therefore P > F$</p> <p data-bbox="233 689 325 723">$R = W$</p> <p data-bbox="209 730 312 763">$F = \mu R$</p> <p data-bbox="209 770 376 804">hence $P > \mu W$</p> <p data-bbox="118 842 185 875">(b)(i)</p>  <p data-bbox="145 1223 408 1256">(ii) $P \times 3a \geq W \times 2a$</p> <p data-bbox="217 1263 568 1330">$\Rightarrow P \geq \frac{2W}{3}$ allow = for A1</p> <p data-bbox="145 1337 715 1413">(c) If $\mu = 0.6$, slides when $P > 0.6W$, if not toppled previously</p> <p data-bbox="209 1426 711 1503">If $\mu = 0.667$, topples when $P > 0.667W$ if not started to slide</p> <p data-bbox="209 1532 363 1565">\therefore slides first</p>	<p data-bbox="740 622 783 656">M1</p> <p data-bbox="740 689 783 723">B1</p> <p data-bbox="740 792 783 826">A1</p> <p data-bbox="740 842 783 875">A1</p> <p data-bbox="740 1223 783 1256">M1</p> <p data-bbox="740 1285 783 1319">A1</p> <p data-bbox="740 1375 783 1408">B1</p> <p data-bbox="740 1464 783 1498">B1</p> <p data-bbox="740 1532 783 1565">B1</p>	<p data-bbox="911 792 935 826">3</p> <p data-bbox="911 842 935 875">1</p> <p data-bbox="911 1285 935 1319">2</p> <p data-bbox="911 1487 935 1520">3</p>	<p data-bbox="991 842 1066 875">forces</p>
Total			9	

Q	Solution	Marks	Total	Comments
6	 <p>Before</p> <p>After</p>			
(a)	$I = \frac{4}{3} \times 3m \times l^2$ $= 4ml^2$	A1	1	AG
(b)(i)	Collision elastic, so $l\omega - v = -(0 - u)$ $\Rightarrow l\omega = u + v$	M1 A1	2	AG
(ii)	Angular momentum before: Rod = 0 Particle = $5m \times ul = 5mul$ \therefore Total = $5mul$ after: Rod = $I\omega = 4ml^2\omega$ Particle = $5mvl$ \therefore Total = $4ml^2\omega + 5mvl$ Momentum conserved $\therefore 5mul = 4ml^2\omega + 5mvl$ $\Rightarrow 5u = 4l\omega + 5v$ $\therefore 5u = 4(u + v) + 5v$ $5u = 4u + 9v$ $\Rightarrow v = \frac{u}{9}$	M1 A1 A1 A1 M1 A1		(Angular momentum attempted)
(iii)	Particle moving in same direction initially	A1	1	
(c)	$l\omega = u + \frac{u}{9}$ $= \frac{10u}{9}$ $\Rightarrow \omega = \frac{10u}{9l}$	A1	1	
		Total	11	
		Total	60	