

General Certificate of Education  
January 2004  
Advanced Level Examination



**MATHEMATICS (SPECIFICATION A)**  
**Unit Pure 5**

**MAP5**

Friday 16 January 2004 Afternoon Session

**In addition to this paper you will require:**

- an 8-page answer book;
- the AQA booklet of formulae and statistical tables.

You may use a standard scientific calculator **only**.

Time allowed: 1 hour 20 minutes

**Instructions**

- Use blue or black ink or ball-point pen. Pencil should only be used for drawing.
- Write the information required on the front of your answer book. The *Examining Body* for this paper is AQA. The *Paper Reference* is MAP5.
- Answer **all** questions.
- All necessary working should be shown; otherwise marks for method may be lost.
- The **final** answer to questions requiring the use of tables or calculators should normally be given to three significant figures.
- Tie loosely any additional sheets you have used to the back of your answer book before handing it to the invigilator.

**Information**

- The maximum mark for this paper is 60.
- Mark allocations are shown in brackets.

**Advice**

- Unless stated otherwise, formulae may be quoted, without proof, from the booklet.

---

Answer **all** questions.

---

- 1 The function  $y(x)$  satisfies the differential equation

$$\frac{d^2y}{dx^2} = x^2 + y^2.$$

Given that  $y = 2$  when  $x = 1$  and that  $y = 2.08$  when  $x = 1.1$ , use the formula

$$y_{r+1} \approx 2y_r - y_{r-1} + h^2 y_r''$$

with a step interval of 0.1 to estimate the value of  $y(1.2)$ . (4 marks)

- 2 (a) By means of a suitable substitution, or otherwise, evaluate

$$\int_0^a \frac{x}{\sqrt{1-x^2}} dx,$$

where  $0 \leq a < 1$ . (4 marks)

(b) Explain why the integral is improper when  $a = 1$ . (1 mark)

(c) Evaluate

$$\int_0^1 \frac{x}{\sqrt{1-x^2}} dx. (2 marks)$$

- 3 (a) A curve  $C$  has polar equation

$$r = e^{k\theta},$$


where  $k \neq 0$  and  $0 \leq \theta \leq \frac{1}{2}\pi$ .

The points  $P$  and  $Q$  on  $C$  have polar coordinates  $(r_1, \theta_1)$  and  $(r_2, \theta_2)$  respectively, where  $\theta_2 > \theta_1$ .

Show that the area  $A$  bounded by  $C$  and the lines  $OP$  and  $OQ$ , where  $O$  is the pole, is given by

$$A = \frac{1}{4k}(r_2^2 - r_1^2). \quad (4 \text{ marks})$$

- (b) The diagram shows a sketch of part of the curve  $r = e^\theta$  and part of the circle  $r = 2$ .



- (i) Find the polar coordinates of  $K$ , the point of intersection of the two curves. (2 marks)
- (ii) Find the area of the shaded region  $R$ , between the curves and the initial line, giving your answer in the form  $p \ln 2 + q$  where  $p$  and  $q$  are rational numbers. (5 marks)

**TURN OVER FOR THE NEXT QUESTION**

Turn over ►

- 4 (a) (i) Use the approximation

$$\cos x \approx 1 - \frac{x^2}{2!} + \frac{x^4}{4!}$$

to show that the first three non-zero terms in the expansion in ascending powers of  $x$  of  $\frac{1}{\cos x}$  are

$$1 + \frac{x^2}{2} + \frac{5x^4}{24}. \quad (5 \text{ marks})$$

- (ii) By writing  $\tan x = \frac{\sin x}{\cos x}$ , find the first three non-zero terms in the expansion in ascending powers of  $x$  of  $\tan x$ . (4 marks)

- (b) Find

$$\lim_{x \rightarrow 0} \left( \frac{\tan 2x - 2x}{\tan x - x} \right). \quad (4 \text{ marks})$$

- 5 (a) Given that  $y = ax^2 + bx$  is a particular integral of the differential equation

$$\frac{d^2y}{dx^2} + \frac{dy}{dx} = x,$$

find the values of  $a$  and  $b$ . (4 marks)

- (b) Hence, or otherwise, solve the differential equation

$$\frac{d^2y}{dx^2} + \frac{dy}{dx} = x,$$

given that  $y = 1$  and  $\frac{dy}{dx} = 3$  when  $x = 0$ . (7 marks)

- 6 (a) The function  $y(x)$  satisfies the differential equation

$$\frac{dy}{dx} = f(x, y),$$

where 
$$f(x, y) = \frac{x^3 + y^3}{xy^2}$$

and 
$$y(1) = 1.$$

Use the formula

$$y_{r+1} \approx y_r + \frac{1}{2}(k_1 + k_2)$$

where 
$$k_1 = h f(x_r, y_r)$$

and 
$$k_2 = h f(x_r + h, y_r + k_1),$$

with  $h = 0.1$  to calculate a value for  $y(1.1)$ , giving your answer to four decimal places. *(5 marks)*

- (b) (i) Use the substitutions  $y = ux$  and  $\frac{dy}{dx} = u + x \frac{du}{dx}$  to show that the differential equation

$$\frac{dy}{dx} = \frac{x^3 + y^3}{xy^2}$$

transforms to

$$x \frac{du}{dx} = \frac{1}{u^2}. \quad (3 \text{ marks})$$

- (ii) Solve the differential equation

$$x \frac{du}{dx} = \frac{1}{u^2}$$

and hence express  $y$  in terms of  $x$ , given that  $y = 1$  when  $x = 1$ . *(5 marks)*

- (iii) Use your solution to part (b)(ii) to find the value of  $y(1.1)$ , giving your answer to four decimal places. *(1 mark)*

**END OF QUESTIONS**