

GCE 2004

June Series



Mark Scheme

Mathematics A

Unit MAS3

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation meeting attended by all examiners and is the scheme which was used by them in this examination. The standardisation meeting ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for the standardisation meeting each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed at the meeting and legislated for. If, after this meeting, examiners encounter unusual answers which have not been discussed at the meeting they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available from:

Publications Department, Aldon House, 39, Heald Grove, Rusholme, Manchester, M14 4NA
Tel: 0161 953 1170

or

download from the AQA website: www.aqa.org.uk

Copyright © 2004 AQA and its licensors

COPYRIGHT

AQA retains the copyright on all its publications. However, registered centres for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to centres to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

The Assessment and Qualifications Alliance (AQA) is a company limited by guarantee registered in England and Wales 3644723 and a registered charity number 1073334. Registered address AQA, Devas Street, Manchester. M15 6EX.

Dr Michael Cresswell Director General

Key to Mark Scheme

M	mark is for	method
m	mark is dependent on one or more M marks and is for.....	method
A	mark is dependent on M or m marks and is for	accuracy
B	mark is independent of M or m marks and is for	method and accuracy
E	mark is for	explanation
✓ or ft or F	follow through from previous incorrect result	
CAO	correct answer only	
AWFW	anything which falls within	
AWRT	anything which rounds to	
AG	answer given	
SC	special case	
OE	or equivalent	
A2,1	2 or 1 (or 0) accuracy marks	
-x EE	deduct x marks for each error	
NMS	no method shown	
PI	possibly implied	
SCA	substantially correct approach	
c	candidate	
SF	significant figure(s)	
DP	decimal place(s)	

Abbreviations used in Marking

MC – x	deducted x marks for mis-copy
MR – x	deducted x marks for mis-read
ISW	ignored subsequent working
BOD	given benefit of doubt
WR	work replaced by candidate
FB	formulae booklet

Application of Mark Scheme

No method shown:

Correct answer without working.....	mark as in scheme
Incorrect answer without working	zero marks unless specified otherwise

More than one method/choice of solution:

2 or more complete attempts, neither/none crossed out	mark both/all fully and award the mean mark rounded down
1 complete and 1 partial attempt, neither crossed out	award credit for the complete solution only

Crossed out work

do not mark unless it has not been replaced

Alternative solution using a correct or partially correct method

award method and accuracy marks as appropriate

MAS3

Q	Solution	Marks	Total	Comments
1(a)	Likely to adjust amount as she goes along/ measures not independent.	E1	1	
(b)	$\hat{\mu} = \bar{x} = \frac{473}{9} = 52.6$ $\hat{\sigma}^2 = s^2 = \frac{24935}{8} - \frac{(473)^2}{8 \times 9} = 9.53$	B1 M1 A1	3	awrt awrt; if error in s^2 from rounding \bar{x} to 4 or 5 sf, lose 1 mark here, then full marks available.
(c)(i)	Assume that weights of flour are normally distributed. $\nu = 9 - 1 = 8$ Critical value of t is 1.86 Confidence limits are $52.6 \pm 1.860 \sqrt{\frac{9.53}{9}}$ giving (50.6 to 50.7, 54.4 to 54.5)	E1 B1 B1 M1 A1✓	6	cao; award here or in (ii) cao allow z; M1 if not divided by 9. ✓ on (b)
(ii)	$\nu = 8$ $\chi_{0.05}^2 = 2.733; \chi_{0.95}^2 = 15.507$ Confidence limits are $\frac{8 \times 9.53}{15.507} \text{ and } \frac{8 \times 9.53}{2.733}$ Confidence interval for σ^2 is (4.92, 27.9) Confidence interval for σ is (2.22, 5.28)	B1 M1 A1✓ A1 A1✓	5	cao; both ✓ on χ^2 values cao ✓ on CI for variance
(d)	The whole of the CI for μ is above 50; Standard deviation seems to be more than 2 grams. Not very useful as Emma overestimates and her measures are rather variable.	E1 E1	2	Reference to CIs required with some assessment.
Total			17	

MAS3 (Cont)

Q	Solution	Marks	Total	Comments
2(a)	H_0 : Median score = 50	B1		both; must refer to average.
	H_1 : Median score \neq 50	B1		
	Differences from 50 are: $+8 - 2 - 10 - 12 + 4 + 1 - 16 + 13 - 11 + 9$	M1		
	Signed ranks are: $+4 - 2 - 6 - 8 + 3 + 1 - 10 + 9 - 7 + 5$	A1		
	$T_+ = 22$; $T_- = 33$	A1✓		either; ✓ on ranks
	Critical value of T is 8	B1		cao
	Accept H_0 . Not enough evidence to say median is not 50.	A1✓	7	✓ on T_{crit} and T_{calc}
(b)(i)	First and last ranks become + 4.5	B1	1	
(ii)	Values of T_+ and T_- unchanged	B1	1	either
(c)	H_0 : Median of Jamie's – Samir's score = 0	B1		or equivalent; both
	H_1 : Median $>$ 0			
	Under H_0 , $X \sim B(15, 0.5)$	B1		cao
	$P(X \geq 12) = P(X \leq 3)$ $= 0.0176$	M1 A1		cao
	$0.0176 < 5\%$ so reject H_0 ; Evidence suggests that Jamie scores higher than Samir on average.	A1✓	5	✓ on probability
	Total		14	

MAS3 (Cont)

Q	Solution	Marks	Total	Comments
3(a)	Shape of histogram similar to pdf of exponential distribution. Mean and SD approximately equal.	E1 E1	2	
(b)(i)	$E(T) = \frac{1}{0.3} = 3.33$	B1	1	awrt
(ii)	$P(T \leq 1) = F(1)$ $= 1 - e^{-0.3}$ $= 0.259$	M1 A1	2	awrt
(iii)	$P(T > 1.75 T > 1)$ $= \frac{1 - F(1.75)}{1 - F(1)} \left(= \frac{1 - F(1.75)}{0.741} \right)$ $= \frac{e^{-0.525}}{e^{-0.3}} = 0.799$	M1 A1 A1✓	3	identifies correct probability. numerator correct ✓ on answer to (b)(ii) B1 for $P(T < 1.75 T > 1)$ correctly evaluated.
(iv)	Let median value be m $F(m) = 0.5$ $1 - e^{-0.3m} = 0.5$ $e^{-0.3m} = 0.5$ $-0.3m = \ln(0.5)$ $m = 2.31$ Median time interval = 2.31 minutes	M1 m1 A1	3	valid attempt to solve cao
Total			11	

MAS3 (Cont)

Q	Solution	Marks	Total	Comments
4(a)(i)	$\frac{s_x^2}{s_y^2} = \frac{1.60}{1.40} = 1.143$	M1 A1		
	$\nu_1 = 10 - 1 = 9; \nu_2 = 7 - 1 = 6$ 90% interval so $p = 0.95$ $F_6^9 = 4.099; F_9^6 = 3.374$ Confidence interval given by $\frac{1}{F_6^9} \leq \frac{\sigma_x^2 / \sigma_y^2}{1.143} \leq F_9^6$	B1 B1 M1		CAO both CAO; either use of
	$\frac{1}{4.099} \leq \frac{\sigma_x^2 / \sigma_y^2}{1.143} \leq 3.374$ giving (0.279, 3.86)	A1 A1✓ A1✓	8	correct values of F right way round; ✓ on Fs ✓ on ratio and F values. M1A1 if one CL correct.
(ii)	Confidence interval includes 1	E1	1	
(b)	$H_0: \mu_x = \mu_y$ $H_1: \mu_x > \mu_y$ Pooled estimate of variance is $\frac{(9 \times 1.6) + (6 \times 1.4)}{15} = 1.52$	B1 M1 A1		both
	$\bar{x} - \bar{y} = 1.16$ $\nu = 15$ Critical value of $t = 1.753$ Sample statistic = $\frac{1.16}{\sqrt{1.52 \left(\frac{1}{10} + \frac{1}{7} \right)}}$ $= 1.91$	B1 B1 B1 M1		CAO CAO
	Sample $t > t_{crit}$ so reject H_0 . Evidence supports Jayne's belief.	A1✓ A1✓	9	✓ on $\bar{x} - \bar{y}$ and variance ✓ on sample t and t_{crit}
	Total		18	
	Total		60	