

General Certificate of Education
June 2004
Advanced Level Examination



**MATHEMATICS AND STATISTICS
(SPECIFICATION B)
Unit Pure 6**

MBP6

Tuesday 29 June 2004 Afternoon Session

In addition to this paper you will require:

- a 12-page answer book;
- the AQA booklet of formulae and statistical tables.

You may use a graphics calculator.

Time allowed: 1 hour 45 minutes

Instructions

- Use blue or black ink or ball-point pen. Pencil should only be used for drawing.
- Write the information required on the front of your answer book. The *Examining Body* for this paper is AQA. The *Paper Reference* is MBP6.
- Answer **all** questions.
- All necessary working should be shown; otherwise marks for method may be lost.
- The **final** answer to questions requiring the use of tables or calculators should normally be given to three significant figures.

Information

- The maximum mark for this paper is 80.
- Mark allocations are shown in brackets.

Advice

- Unless stated otherwise, formulae may be quoted, without proof, from the booklet.

Answer **all** questions.

1 Show that the curve with equation

$$y = 3 \tanh x + 4 \operatorname{sech} x$$

has a stationary point when $\sinh x = \frac{3}{4}$. (5 marks)

2 The complex number $\omega = 30\sqrt{31} + 194i$.

(a) Express ω in the form $re^{i\theta}$, where $r > 0$ and θ is to be given to four decimal places in radians between $-\pi$ and π . (2 marks)

(b) The complex number z_1 is the complex number with least positive argument such that $z_1^8 = \omega$. Determine $|z_1|$ in an exact form, and find $\arg(z_1)$ to four decimal places. (2 marks)

(c) On a diagram of the complex plane, mark the position of z_1 , and describe the positions of the remaining seven complex roots of the equation $z^8 = \omega$. (3 marks)

3 (a) Find the general solution of the second-order differential equation

$$\frac{d^2y}{dx^2} + 2\frac{dy}{dx} + y = 8e^{3x} \quad (7 \text{ marks})$$

(b) Given that $y = 1$ and $\frac{dy}{dx} = 2$ when $x = 0$, solve this differential equation. (3 marks)

4 (a) Prove the identity

$$\sin x + \sin 2x + \sin 3x + \sin 4x \equiv 4 \cos \frac{x}{2} \cos x \sin \frac{5x}{2} \quad (5 \text{ marks})$$

(b) Hence find all solutions in the interval $0 \leq x \leq \pi$ of the equation

$$\sin x + \sin 2x + \sin 3x + \sin 4x = 0$$

giving your answers in terms of π . (4 marks)

- 5 (a) Use mathematical induction to prove that $(\cos \theta + i \sin \theta)^n = \cos n\theta + i \sin n\theta$ for all positive integers n . (4 marks)
- (b) (i) Express $(-\sqrt{3} + i)^n$ in the form $2^n(\cos n\theta + i \sin n\theta)$, giving the value of θ in terms of π . (3 marks)
- (ii) Hence find the least positive integer value of n for which $(-\sqrt{3} + i)^n$ is a positive real number. (2 marks)

- 6 (a) Determine the eigenvalues and corresponding eigenvectors of the matrix

$$\mathbf{M} = \begin{bmatrix} 8 & 6 \\ 6 & 17 \end{bmatrix} \quad (6 \text{ marks})$$

- (b) The transformation \mathbf{T} has matrix \mathbf{M} .

(i) Show that the invariant lines of \mathbf{T} are perpendicular. (2 marks)

(ii) Give a full geometrical interpretation of \mathbf{T} . (3 marks)

- 7 (a) Differentiate $\sqrt{1+t^2}$ with respect to t . (1 mark)

(b) (i) Given that $I_n = \int_0^1 \frac{t^n}{\sqrt{1+t^2}} dt$, where $n \geq 0$, show that

$$nI_n = \sqrt{2} - (n-1)I_{n-2}, \quad n \geq 2 \quad (5 \text{ marks})$$

(ii) Evaluate I_3 in surd form. (3 marks)

- (c) Use the substitution $t = \tan \frac{1}{2}x$ to evaluate

$$\int_0^{\frac{\pi}{2}} \sec \frac{1}{2}x \tan^3 \frac{1}{2}x \, dx \quad (3 \text{ marks})$$

TURN OVER FOR THE NEXT QUESTION

Turn over ►

8 (a) Using the definitions $\sinh \theta = \frac{1}{2}(e^\theta - e^{-\theta})$ and $\cosh \theta = \frac{1}{2}(e^\theta + e^{-\theta})$, prove that:

(i) $2 \sinh \theta \cosh \theta \equiv \sinh 2\theta$;

(ii) $2 \sinh^2 \theta \equiv \cosh 2\theta - 1$. *(4 marks)*

(b) Use the substitution $\cosh \theta = 2x + 1$ to show that:

(i) $\int \sqrt{4x^2 + 4x} \, dx = k \int \sinh^2 \theta \, d\theta$, stating the value of k ; *(4 marks)*

(ii) $\int \sqrt{4x^2 + 4x} \, dx = \frac{1}{4}(2x + 1)\sqrt{4x^2 + 4x} - \frac{1}{4}\cosh^{-1}(2x + 1) + C$, where C is an arbitrary constant. *(4 marks)*

(c) Find the length of the arc of the curve with equation

$$y = \frac{1}{4}(2x + 1)\sqrt{4x^2 + 4x} - \frac{1}{4}\cosh^{-1}(2x + 1)$$

between the points where $x = 77$ and $x = 89$. *(5 marks)*

END OF QUESTIONS