General Certificate of Education January 2005 Advanced Level Examination

MBP6

MATHEMATICS AND STATISTICS (SPECIFICATION B) Unit Pure 6

Tuesday 1 February 2005 Morning Session

In addition to this paper you will require:

- a 12-page answer book;
- the AQA booklet of formulae and statistical tables.

You may use a graphics calculator.

Time allowed: 1 hour 45 minutes

Instructions

- Use blue or black ink or ball-point pen. Pencil should only be used for drawing.
- Write the information required on the front of your answer book. The *Examining Body* for this paper is AQA. The *Paper Reference* is MBP6.
- Answer all questions.
- All necessary working should be shown; otherwise marks for method may be lost.
- The **final** answer to questions requiring the use of tables or calculators should normally be given to three significant figures.

Information

- The maximum mark for this paper is 80.
- Mark allocations are shown in brackets.

Advice

• Unless stated otherwise, formulae may be quoted, without proof, from the booklet.


P75479/0205/MBP6 6/6/6/ MBP6

Answer all questions.

1 Find the general solution of the differential equation

$$4\frac{d^2y}{dx^2} - 8\frac{dy}{dx} + 5y = 0$$
 (5 marks)

- 2 Find the area of the region bounded by the curve $y = \cosh x + \mathrm{sech}^2 x$, the x-axis and the lines x = 0 and $x = \ln 2$.
- 3 The Argand diagram of the complex plane shows the three points U, V and O which represent the complex numbers u, v and 0 + 0i respectively.



- (a) Write down expressions involving u and v which represent:
 - (i) the length of the line segment UV;
 - (ii) the angle UOV. (2 marks)
- (b) On a copy of this Argand diagram, draw the point W which represents the complex number w = u + v. (1 mark)
- 4 A curve has parametric equations $x = 2 \ln t$, $y = t + \frac{1}{t}$, t > 0.
 - (a) Show that, for this curve, $\left(\frac{\mathrm{d}x}{\mathrm{d}t}\right)^2 + \left(\frac{\mathrm{d}y}{\mathrm{d}t}\right)^2 = \left(\frac{t^2+1}{t^2}\right)^2$. (4 marks)
 - (b) The arc of this curve between the points where t = 1 and t = 2 is rotated through 2π radians about the x-axis to form a surface of revolution with area S. Find the exact value of S. (6 marks)

- 5 (a) Using the definitions of $\sinh x$ and $\cosh x$ in terms of exponential functions, prove the identity $2 \sinh x \cosh x \equiv \sinh 2x$. (2 marks)
 - (b) Solve the differential equation

$$\frac{\mathrm{d}y}{\mathrm{d}x} + y \tanh x = \sinh x$$

given that y = 1 when x = 0.

(7 marks)

6 (a) Given that $t = \tan \frac{1}{2}x$, show that

$$\sin x \tan \frac{1}{2}x + 2\sec x = \frac{2 + 6t^2}{(1 - t^2)(1 + t^2)}$$
 (2 marks)

(b) Solve the equation

$$2\sin x \tan \frac{1}{2}x + 4\sec x + 5 = 0$$

giving all answers for x in radians in the interval $0 < x < 2\pi$.

(6 marks)

(c) (i) Use the substitution $t = \tan \frac{1}{2}x$ to show that

$$\int_0^{\frac{\pi}{3}} \frac{3}{\sin x \tan \frac{1}{2}x + 2\sec x} dx = \int_0^{\alpha} \frac{3 - 3t^2}{1 + 3t^2} dt$$

stating the exact value of α .

(4 marks)

(ii) Hence evaluate exactly
$$\int_0^{\frac{\pi}{3}} \frac{3}{\sin x \tan \frac{1}{2}x + 2 \sec x} dx.$$
 (4 marks)

7 (a) Determine the real part of $(1 + i \tan \theta)^3$.

(2 marks)

(b) Deduce the identity

$$1 - 3\tan^2\theta \equiv \frac{\cos 3\theta}{\cos^3\theta} \tag{3 marks}$$

8 (a) Determine the eigenvalues and corresponding eigenvectors of the matrix

$$\begin{bmatrix} 3 & 5 \\ 2 & 6 \end{bmatrix}$$
 (6 marks)

- (b) The transformation **T** is given by x' = 3x + 5y + 1, y' = 2x + 6y + 1.
 - (i) Show that (2, -1) is a fixed point of **T**. (2 marks)
 - (ii) Hence express T in the form

$$\begin{bmatrix} x' - \alpha \\ y' - \beta \end{bmatrix} = \begin{bmatrix} 3 & 5 \\ 2 & 6 \end{bmatrix} \begin{bmatrix} x - \alpha \\ y - \beta \end{bmatrix}$$

for suitable values of the constants α and β .

(1 mark)

- (iii) Hence, using the answer to part (a), find the equation of the line of fixed points of **T**. (2 marks)
- (c) Show that all lines parallel to y = x are fixed lines of **T**. (3 marks)
- 9 For integers $n \ge 0$, let $I_n = \int_0^{\frac{\pi}{3}} e^{3x} \tan^n x \, dx$.
 - (a) Show that, for $n \ge 1$,

$$nI_{n+1} + 3I_n + nI_{n-1} = \left(\sqrt{3}\right)^n e^{\pi}$$
 (5 marks)

(b) (i) By considering the cases when n = 1 and n = 3, or otherwise, show that

$$I_4 + I_3 - 3I_1 = I_0$$
 (5 marks)

(ii) Hence evaluate exactly

$$\int_0^{\frac{\pi}{3}} e^{3x} \tan x \left(\tan^3 x + \sec^2 x - 4 \right) dx$$
 (4 marks)

END OF QUESTIONS