



**ADVANCED**  
**General Certificate of Education**  
**2011**

---

**Mathematics**  
**Assessment Unit C4**  
*assessing*  
**Module C4: Core Mathematics 4**  
**[AMC41]**  
**WEDNESDAY 1 JUNE, MORNING**

---

**MARK  
SCHEME**

		AVAILABLE MARKS
1	(i) $\vec{AB} = \vec{AO} + \vec{OB} = -3\mathbf{i} + \mathbf{j} + 2\mathbf{i} + 6\mathbf{j}$ $= -\mathbf{i} + 7\mathbf{j}$	M1W1
	(ii) $ \vec{AB}  = \sqrt{(-1)^2 + (7)^2} = \sqrt{50} = 5\sqrt{2}$	M1W1
	(iii) $\vec{OA} \cdot \vec{OB} = (3\mathbf{i} - \mathbf{j}) \cdot (2\mathbf{i} + 6\mathbf{j})$ $= 6 - 6$	M1
		W1
	(iv) $90^\circ$	MW1
		7
2	(i) $x + xy - 12$ $1 + y + \frac{dy}{dx} x$	M2W2
	(ii) at $(2, 5)$ $1 + 5 + 2 \frac{dy}{dx} = 0$ $\frac{dy}{dx} = -3$	MW1
	Eqn: tangent $y - 5 = -3(x - 2)$ $y = -3x + 11$ $y + 3x = 11$	M1 W1 7

3 (i)  $2 \cos x + 4 \sin x = R \cos(x - \alpha)$

$$\cos \alpha = 2/R$$

$$\sin \alpha = 4/R$$

$$\tan \alpha = 2 \Rightarrow \alpha = 63.43^\circ$$

M1W1

$$= 63.4^\circ \text{ (3.s.f.)}$$

$$R = \sqrt{2^2 + 4^2} = \sqrt{20} = 2\sqrt{5}$$

M1W1

$$2\sqrt{5} \cos(x - 63.4^\circ)$$

(ii)  $2 \cos x + 4 \sin x = 3$

$$\sqrt{20} \cos(x - 63.4^\circ) = 3$$

M1

$$\cos(x - 63.4^\circ) = \frac{3}{\sqrt{20}}$$

$$x - 63.4^\circ = 47.8695^\circ \text{ or } 312.13^\circ$$

MW2

$$x = 111^\circ \text{ or } 15.6^\circ$$

W2

9

4 (i) Volume =  $\int_{-12}^3 \pi y^2 dx$

M2

$$= \int_{-12}^3 \pi (144 - x^2) dx$$

MW1

$$= \pi \left[ 144x - \frac{x^3}{3} \right]_{-12}^3$$

MW1

$$= \pi \left[ (432 - 9) - (-1728 + 576) \right]$$

M1

$$= 1575 \pi \text{ cm}^3$$

W2


(ii) The bowl needs a flat bottom

MW1

8

		AVAILABLE MARKS
5	(i) $\sin(A+B) = \sin A \cos B + \cos A \sin B$	M1
	$\sin(A+A) = \sin A \cos A + \cos A \sin A$	M1
	$= 2 \sin A \cos A$	W1
(ii)	LHS $\tan A + \cot A$	
	$\frac{\sin A}{\cos A} + \frac{\cos A}{\sin A}$	M1W1
	$\frac{\sin^2 A + \cos^2 A}{\sin A \cos A}$	MW1
	$\frac{1}{\frac{1}{2} \sin 2A}$	M2
	$\frac{2}{\sin 2A} = \text{R.H.S.}$	W1
		9
6	$\frac{dx}{dt} = k(3-x)(4-x)$	
	$\int_0^2 \frac{dx}{(3-x)(4-x)} = \int_0^{10} k dt$	M2W2
	$\frac{1}{(3-x)(4-x)} = \frac{A}{3-x} + \frac{B}{4-x}$	MW1
	$1 = A(4-x) + B(3-x)$	MW1
Let	$x = 4 \quad 1 = -B \Rightarrow B = -1$	MW2
	$x = 3 \quad 1 = A$	
	$\frac{1}{3-x} - \frac{1}{4-x}$	
	$\int_0^2 \left( \frac{1}{3-x} - \frac{1}{4-x} \right) dx = \int_0^{10} k dt$	MW1

		AVAILABLE MARKS
$\left[ -\ln  3-x  + \ln  4-x  \right]_0^2 = [k t]_0^{10}$	W3	
$\left[ \ln \left  \frac{4-x}{3-x} \right  \right]_0^2 = 10 k$		
$\ln 2 - \ln \frac{4}{3} = 10 k$		
$k = \frac{1}{10} \ln \frac{3}{2}$	W2	
$[k = 0.0405]$		14
7 (i) $x = 1$ $y = 2$	MW2	
(ii) $f : x \rightarrow \frac{2x+3}{x-1}$		
Let $y = \frac{2x+3}{x-1}$	M1	
$y(x-1) = 2x + 3$	M1	
$yx - 2x = y + 3$		
$x = \frac{y+3}{y-2}$	M1W1	
$f^{-1} : x \rightarrow \frac{x+3}{x-2}$	W1	
domain $x > 2$	MW1	8

		AVAILABLE MARKS
8	(i) $\int x \ln x \, dx$  $= \frac{x^2}{2} \ln x - \int \frac{x^2}{2} \frac{1}{x} \, dx$ $= \frac{x^2}{2} \ln x - \frac{x^2}{4} + c$	M1 W3 W2
	(ii) (a) $\int x \ln x^2 \, dx$ $= \int 2x \ln x \, dx$ $= x^2 \ln x - \frac{x^2}{2} + d$	M1W1 MW1
	(b) $\int x \ln 3x \, dx$ $= \int x (\ln 3 + \ln x) \, dx$ $= \int x \ln 3 + x \ln x \, dx$ $= \frac{x^2}{2} \ln 3 + \frac{x^2}{2} \ln x - \frac{x^2}{4} + g$	M1W1 MW2
		13
	<b>Total</b>	<b>75</b>