

# EDEXCEL - LONDON EXAMINATIONS

Stewart House 32 Russell Square London WC1B 5DN

June 2001

Advanced Supplementary/Advanced Level

General Certificate of Education

Subject MECHANICS 6677

Paper No. M1

Question number	Scheme	Marks
1.	<p> <math>3 \rightarrow</math>      <math>\leftarrow 2</math> Before  <math>0.5 \circlearrowleft</math>    <math>\circlearrowright 0.2</math>  <math>\rightarrow</math>      <math>\rightarrow v</math> After  <math>1.5</math> </p> <p>(a) <math>0.5 \times 3 - 0.2 \times 2 = 0.5 \times 1.5 + 0.2 \times v</math> <span style="float: right;">(Mom<sup>m</sup> eqn. with 4 terms)</span></p> <p style="text-align: center;"><math>\Rightarrow v = \underline{1.75 \text{ ms}^{-1}}</math></p> <p>(b) <math>I = 0.2(2 + 1.75)</math></p> <p style="text-align: center;"><math>= \underline{0.75 \text{ N s}}</math></p>	<p>M1 A1</p> <p>A1 (3)</p> <p>M1 A1 ✓</p> <p>A1 (3)</p> <p style="text-align: center;">(6)</p>

# EDEXCEL - LONDON EXAMINATIONS

Stewart House 32 Russell Square London WC1B 5DN



June 2001

Advanced Supplementary/Advanced Level

General Certificate of Education

Subject MECHANICS 6677

Paper No. M1

Question number	Scheme	Marks
<u>2.</u>	<u>EITHER</u>	
(a)	 <p>Vector <math>\Delta</math> attempt Correct</p> $F^2 = 5^2 + 3^2 - 2 \cdot 5 \cdot 3 \cos 140 \quad (\text{cos rule})$ $\rightarrow F \approx \underline{7.55 \text{ N}}$	M1 A1 M1 A1 A1 (5)
(b)	$\frac{F}{\sin 140} = \frac{3}{\sin \theta} \Rightarrow \theta \approx \underline{14.8^\circ}$	M1 A1, A1 (3) (8)
or.		
(a)	 <p>Vector <math>\Delta</math> attempt correct</p> $F^2 = (5 + 3 \cos 40)^2 + (3 \sin 40)^2$ $F \approx \underline{7.55 \text{ N}}$	M1 A1 M1 A1 ✓ A1 (5)
(b)	$\tan \theta = \frac{3 \sin 40}{5 + 3 \cos 40}, \quad \theta \approx \underline{14.8^\circ}$	M1 A1, ✓ A1 (3) (8)
or		
(a)	$\underline{P} = \begin{pmatrix} 5 \\ 0 \end{pmatrix} \text{ or } 5\underline{i} \quad \underline{Q} = \begin{pmatrix} 3 \cos 40 \\ 3 \sin 40 \end{pmatrix} \text{ or } 3 \cos 40 \underline{i} + 3 \sin 40 \underline{j}$ $\Rightarrow \underline{F} = \begin{pmatrix} 5 + 3 \cos 40 \\ 3 \sin 40 \end{pmatrix}$	M1 A1
	$ \underline{F}  = \sqrt{(5 + 3 \cos 40)^2 + (3 \sin 40)^2}$ $\approx \underline{7.55 \text{ N}}$	M1 A1 ✓ A1 (5)
(b)	$\tan \theta = \frac{3 \sin 40}{5 + 3 \cos 40}$ $\approx \underline{14.8^\circ}$	M1 A1 ✓ A1 (3) (8)

# EDEXCEL - LONDON EXAMINATIONS

Stewart House 32 Russell Square London WC1B 5DN


June 2001

Advanced Supplementary/Advanced Level

General Certificate of Education

Subject MECHANICS 6677

Paper No. M1

Question number	Scheme	Marks
3.	<p>(a) Distance = <math>\frac{1}{2} \times (30+17) \times 3, + 4 \times 17</math>  <math>= 138.5 \text{ m.}</math></p> <p>[OR <math>\frac{1}{2} \times 3 \times (30-17) + 3 \times 17 + 4 \times 17</math>  <math>= 138.5 \text{ m}</math>]</p> <p>(b) Str. line graph <math>\Rightarrow</math> const. decel<sup>2</sup>  <math>"F=ma" \Rightarrow \underline{F \text{ const}}</math></p> <p>(c) Decel<sup>2</sup> = <math>\frac{30-17}{3}</math>  Force = <math>1200 \times \left(\frac{30-17}{3}\right) = \underline{5200 \text{ N}}</math></p>	<p>m1 A1, m1  A1 (4)</p> <p>m1 A1, m1  A1</p> <p>m1  A1 c.s.o.  (2)</p> <p>m1  m1 A1  (3)  (9)</p>
4.	<p>(a)    Diag. with 4 forces marked  (Allow F &amp; R combined if clear)</p> <p>(b) R(<math>\uparrow</math>) <math>R = 3g \cos 30^\circ + 30 \sin 30^\circ</math> (3 terms)  <math>= 40.46 \dots \approx 40.5 \text{ or } 40 \text{ N.}</math></p> <p>(c) R(<math>\leftarrow</math>) <math>F = 30 \cos 30^\circ - 3g \sin 30^\circ</math> (3 terms)  <math>F = \mu R, \Rightarrow \mu = \frac{F}{R} = \frac{11.28}{40.46}</math>  <math>\approx 0.28 \text{ (or } 0.279)</math></p>	<p>B2  -1 e.e. (2)</p> <p>m1 A2  -1 e.e.  A1 (4)</p> <p>m1 A1</p> <p>m1, m1  A1 (5)</p> <p>(11)</p>

# EDEXCEL - LONDON EXAMINATIONS

Stewart House 32 Russell Square London WC1B 5DN



June 2001

Advanced Supplementary/Advanced Level

General Certificate of Education

Subject MECHANICS 6677

Paper No. M1

Question number	Scheme	Marks
5.	<p>(a) <math>\circ</math></p> <p>(b)  <math>M(D): 2W = 1500 \cdot 5</math>  <math>\Rightarrow W = \underline{3750 \text{ N}}</math></p> <p>[If moments about another pt: M1 for a complete method to get W, A1 for a moments eqn<sup>2</sup> correct.]</p> <p>(c)  <math>M(D) 1500 \cdot 5 = W'(4-x)</math>  <math>M(C) 1000 \cdot 5 = W'x</math>                      Solve <math>\rightarrow W' = \underline{3125 \text{ N}}</math></p> <p>(d) <math>x = 1.6 \text{ m}</math></p> <p>(e) AB remains straight line (o.e.)</p>	<p>B1 (1)</p> <p>M1 A1</p> <p>A1 (3)</p> <p>M1 A1</p> <p>M1 A1</p> <p>M1 A1 (6)</p> <p>M1 A1 (2)</p> <p>B1 (1)</p> <p>(13)</p>

# EDEXCEL - LONDON EXAMINATIONS

Stewart House 32 Russell Square London WC1B 5DN




June 2001

Advanced Supplementary/Advanced Level

General Certificate of Education

Subject MECHANICS 6677

Paper No. M1

Question number	Scheme	Marks
<p><u>b.</u></p>	<p>(a) Car + Van: <math>3200 a = 2320 - 800 - 240</math>  <math>a = \underline{0.4 \text{ ms}^{-2}}</math></p> <p>(b) Car:  <math>1200 a = T - 240</math>  <math>\rightarrow T = 720 \text{ N}</math></p> <p>[OR Van:  <math>2000 a = 2320 - 800 - T</math>  <math>\rightarrow T = 720</math>]</p> <p><u>NB</u> If use eqn<sup>s</sup> for car &amp; van alone, allow M1 A2 for one eqn<sup>2</sup> involving T, then M1 A1 for a second eqn<sup>2</sup> <u>provided</u> it is part of a complete method to find a/T.  Then A1 A1 for a &amp; T.</p> <p>(c)  <math>3200 a' = 2320 - 1040 - 3200g \cdot \frac{1}{20}</math> (4 terms)  <math>a' = -0.09 \text{ ms}^{-2}</math>  <math>\Rightarrow</math> magn. <math>0.09 \text{ ms}^{-2}</math>  speed decreasing</p>	<p>M1 A1  A1 (3)</p> <p>M1 A2 ✓  -1 e.e.  A1 (4)</p> <p>M1 A2  -1 e.e.  M1  A1  A1 ✓ (6)</p> <p>(13)</p>

# EDEXCEL - LONDON EXAMINATIONS

Stewart House 32 Russell Square London WC1B 5DN


June 2001

Advanced Supplementary/Advanced Level

General Certificate of Education

Subject MECHANICS 6677

Paper No. M1

Question number	Scheme	Marks
7.	 <p>(a) <math>\underline{w}_1 = 2\underline{j} + 6\underline{i} + 8\underline{j}</math>  <math>= 6\underline{i} + 8\underline{j}</math></p> <p>(b) <math>OW_1 = \sqrt{(6^2 + 8^2)} = 10 \text{ km}</math>  <math>\text{Est. time} = \frac{10}{5} = 2 \text{ hrs}</math></p> <p>(c) <math>\underline{w}_2 = 2\underline{j} + 6\underline{i} - 6\underline{j}</math>  <math>= 6\underline{i} - 4\underline{j}</math></p> <p>(d) P.v. of rescue party after 1 hour =  <math>\underline{R} = 3\underline{i} + 4\underline{j}</math>  <math>\underline{R}\underline{w}_2 = 3\underline{i} - 8\underline{j}</math>  <math>\tan \theta = \frac{3}{8} = 20.6^\circ</math>  <math>\Rightarrow</math> Required bearing = <math>180^\circ - 20.6^\circ</math>  <math>= 159.4^\circ</math></p>	<p>BI BI (2)</p> <p>M1 M1 A1 (3)</p> <p>BI, M1 A1 (3)</p> <p>M1 A1</p> <p>M1 A1</p> <p>M1</p> <p>M1</p> <p>A1 (7)</p> <p>(15)</p>