


Question Number	Scheme	Marks
1. (a)	$a = 0.25$ $\frac{2\pi}{\omega} = 2 \Rightarrow \omega = \pi$ $-0.125 = 0.25 \cos \omega t$	B1 B1 M1A1
(b)	$t = \frac{1}{\pi} \cos^{-1}(-0.5)$ $= \frac{2}{3}$	M1 A1 (6) (6 marks)
2. (a)	$(\uparrow) 3mg \cos \alpha^\circ = mg$ $\alpha = \cos^{-1}(\frac{1}{3})$ $= 70.5$	M1 A1 M1 A1 (4)
(b)	$(\leftarrow) 3mg \sin \alpha = mr \times 2gk$ $l \sin \alpha = r$ $l = \frac{3}{2} k$	M1 A1 B1 M1 A1 (5) (9 marks)
3. (a)	$2e^{-0.1x} = 2.5a$ $\frac{4}{5} e^{-0.1x} = v \frac{dv}{dx}$ $-8e^{-0.1x} = \frac{1}{2} v^2 (+c)$ $x = 0, v = 2 \Rightarrow c = -10$ $v^2 = 20 - 16e^{-0.1x}$	M1 A1 M1 A1 M1 A1 (6)
(b)	$16 = 20 - 16e^{-0.1x} \Rightarrow e^{-0.1x} = \frac{1}{4}$ $0.1x = \ln 4$ $x = 13.9$	M1 M1 A1 (3)
(c)	Appropriate comment.	B1 (1) (10 marks)

Question Number	Scheme	Marks																		
4. (a)	$\frac{1}{2} \times 0.2 \times 5^2 - \frac{1}{2} \times 0.2 \times u^2 = \frac{1}{2} \times \frac{20(0.5)^2}{1.5}$ $u^2 = \frac{25}{3}$ $u = 2.89 \text{ ms}^{-1}$	M1 A1 A1 M1 A1 (5)																		
(b)	$\frac{1}{2} \times 0.2 \times 5^2 - \frac{1}{2} \times 0.2 \times 1.5^2 = \frac{1}{2} \times \frac{20x^2}{1.5}$ $x^2 = 0.34125$ $T = \frac{20x}{1.5} = 7.8 \text{ N}$	M1 A1 M1 M1 A1 (5) (10 marks)																		
5. (a)	<table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 33%;">Cone</td> <td style="width: 33%;">Cylinder</td> <td style="width: 33%;">Whole</td> </tr> <tr> <td>$\frac{1}{3} \pi (2r)^2 h$</td> <td>$\pi r^2 h$</td> <td>$\frac{1}{3} \pi (2r)^2 h + \pi r^2 h$</td> </tr> <tr> <td>(4)</td> <td>(3)</td> <td>(7)</td> </tr> <tr> <td>$\frac{1}{4} h$</td> <td>$\frac{1}{2} h$</td> <td>\bar{x}</td> </tr> <tr> <td>$- 4 \times \frac{1}{4} h$</td> <td>$+ 3 \times \frac{1}{2} h$</td> <td>$= 7\bar{x}$</td> </tr> <tr> <td colspan="3">$\bar{x} = \frac{1}{14} h$</td> </tr> </table>	Cone	Cylinder	Whole	$\frac{1}{3} \pi (2r)^2 h$	$\pi r^2 h$	$\frac{1}{3} \pi (2r)^2 h + \pi r^2 h$	(4)	(3)	(7)	$\frac{1}{4} h$	$\frac{1}{2} h$	\bar{x}	$- 4 \times \frac{1}{4} h$	$+ 3 \times \frac{1}{2} h$	$= 7\bar{x}$	$\bar{x} = \frac{1}{14} h$			M1 A1 B1 B1 M1 A1 M1 A1 cso (8)
Cone	Cylinder	Whole																		
$\frac{1}{3} \pi (2r)^2 h$	$\pi r^2 h$	$\frac{1}{3} \pi (2r)^2 h + \pi r^2 h$																		
(4)	(3)	(7)																		
$\frac{1}{4} h$	$\frac{1}{2} h$	\bar{x}																		
$- 4 \times \frac{1}{4} h$	$+ 3 \times \frac{1}{2} h$	$= 7\bar{x}$																		
$\bar{x} = \frac{1}{14} h$																				
(b)	 Use of G above N $\tan \alpha = \frac{r}{h - \frac{1}{14} h} = \frac{7}{26}$ $r = \frac{1}{4} h$	M1 M1 A1 A1 (4) (12 marks)																		

Question Number	Scheme	Marks
6. (a)	$mg = \frac{8mge}{4a}$ $\frac{9}{2}a = AO$	M1 A1 (2)
(b)	$mg - \frac{8mg}{4a}(e + x) = m\ddot{x}$ $\ddot{x} = -\frac{2g}{a}x$ $T = 2\pi\sqrt{\frac{a}{2g}} = \pi\sqrt{\frac{2a}{g}}$ (*) $v = d\omega$ $\frac{1}{2}\sqrt{ga} = d\sqrt{\frac{2g}{a}}$ $d = \frac{a}{2\sqrt{2}} = a\frac{\sqrt{2}}{4} = 0.35a$ (awrt)	M1 M1 A1 M1 A1 M1 A1 (7) M1 A1 ft on ω A1 (3)
(d)	Partly under gravity, partly SHM	B1 B1 (2) (14 marks)
7. (a)	$\frac{1}{2}mu^2 = mgl(1 - \cos \theta)$ $u = \sqrt{\frac{2}{3}}gl$	M1 A1 A1 A1 (4)
(b)	$T - mg \cos \theta = \frac{mv^2}{l}$ $\frac{1}{2}mu^2 - \frac{1}{2}mv^2 = mgl(1 - \cos \theta)$ eliminating v^2 , $T = \frac{mg}{3}(9 \cos \theta - 4)$ (*)	M1 A1 M1 A1 cso (6)
(c)	$\max T, \theta = 0, T_{MAX} = \frac{5mg}{3}$ $\min T, \cos \theta = \frac{2}{3}, T_{MIN} = \frac{2mg}{3}$ $\frac{2mg}{3} \leq T \leq \frac{5mg}{3}$	M1 M1 A1 A1 (4) (14 marks)