

EDEXCEL PURE MATHEMATICS P4 (6671) – JANUARY 2003 PROVISIONAL MARK SCHEME

Question Number	Scheme	Marks
1.	$zw =$ $12 \left( \cos \frac{\pi}{4} \cos \frac{2\pi}{3} - \sin \frac{\pi}{4} \sin \frac{2\pi}{3} \right) + 12i \left( \sin \frac{\pi}{4} \cos \frac{2\pi}{3} + \cos \frac{\pi}{4} \sin \frac{2\pi}{3} \right)$ $= 12 \left[ \cos \frac{11\pi}{12} + i \sin \frac{11\pi}{12} \right]$	B1 for 12  M1 A1  (3 marks)
2. (a)	<p>shape</p> <p>points on axes</p>	B1  B1 (2)
(b)	$-2x + 3 = 5x - 1$ $x = \frac{4}{7}$ $x > \frac{4}{7}$	M1  A1  A1 ft  (3)  (5 marks)

**EDEXCEL PURE MATHEMATICS P4 (6671) – JANUARY 2003 PROVISIONAL MARK SCHEME**

Question Number	Scheme	Marks
3. (a)	$\frac{1}{r+1} - \frac{1}{r+3}$	B1 B1 (2)
(b)	$\sum_1^n \frac{1}{r+1} - \frac{1}{r+3} = \frac{1}{2} - \cancel{\frac{1}{4}}$ $+ \frac{1}{3} - \frac{1}{5}$ $+ \cancel{\frac{1}{4}} - \cancel{\frac{1}{6}}$ $\vdots$ $+ \cancel{\frac{1}{n}} - \frac{1}{n+2}$ $+ \cancel{\frac{1}{n+1}} - \frac{1}{n+3}$ $= \left( \frac{1}{2} + \frac{1}{3} \right) + \left( -\frac{1}{n+2} - \frac{1}{n+3} \right)$ $= \frac{5}{6} - \left( \frac{5n^2 + 25n + 30 - 12n - 30}{6(n+2)(n+3)} \right)$ $= \frac{n(5n+13)}{6(n+2)(n+3)} *$	M1 A1 A1 M1 A1 cs (5) <b>(7 marks)</b>


EDEXCEL PURE MATHEMATICS P4 (6671) – JANUARY 2003 PROVISIONAL MARK SCHEME

Question Number	Scheme	Marks
4. (a)	$f(2) = -1.514$ $f(\pi) = 1.142$  $\frac{\pi - \alpha}{\alpha - 2} = \frac{1.142}{1.514}$ $\pi \times 1.514 + 2 \times 1.142 = (1.142 + 1.514)\alpha$ $\alpha = 2.65$	B1 B1 M1 A1 (4)
(b)	$f(x) = 4 \cos 2x + 1$ $f(2.8) = -0.4625$ $f'(2.8) = 4.1023$ $x_2 = 2.8 - \frac{(-0.4625)}{4.1023}$ $= 2.91 \quad \text{only}$	$k \cos 2x + c$ M1 A1 M1 A1
		(9 marks)


**EDEXCEL PURE MATHEMATICS P4 (6671) – JANUARY 2003 PROVISIONAL MARK SCHEME**

Question Number	Scheme	Marks
5. (a)	$v + x \frac{dv}{dx} = (4 + v)(1 + v)$ $x \frac{dv}{dx} = v^2 + 5v + 4 - v$ $x \frac{dv}{dx} = (v + 2)^2 \quad *$	M1, M1 A1 A1 (4)
(b)	$\int \frac{1}{(v+2)^2} dv = \int \frac{1}{x} dx$ $-\frac{1}{2+v} = \ln x + c$ $2+v = -\frac{1}{\ln x + c}$ $v = -\frac{1}{\ln x + c} - 2$	must have $+ c$ M1 A1 M1 A1 (5)
(c)	$y = -2x - \frac{x}{\ln x + c}$	B1 (1)  <b>(10 marks)</b>

**EDEXCEL PURE MATHEMATICS P4 (6671) – JANUARY 2003 PROVISIONAL MARK SCHEME**

Question Number	Scheme	Marks
6. (a)	$z^2 = (3 - 3i)(3 - 3i) = -18i$	M1 A1 (2)
(b)	$\frac{1}{z} = \frac{(3 + 3i)}{(3 - 3i)(3 + 3i)} = \frac{3 + 3i}{18} = \frac{1+i}{6}$	M1 A1 (2)
(c)	$ z  = \sqrt{(9+9)} = \sqrt{18} = 3\sqrt{2}$ $ z  = 18$	two correct M1 all three correct A1 (2)
(d)		two correct B1 four correct B1 (2)
(e)	$\frac{OB}{OD} = 18, \quad \frac{OA}{OC} = \frac{3\sqrt{2}}{\sqrt{2}/6} = 18$ $\angle AOB = \angle COD = 45^\circ \therefore \text{similar}$	M1 A1 B1 (3) <b>(11 marks)</b>

EDEXCEL PURE MATHEMATICS P4 (6671) – JANUARY 2003 PROVISIONAL MARK SCHEME

Question Number	Scheme	Marks
7. (a)	$y = \lambda x \cos 3x$ $\frac{dy}{dx} = \lambda \cos 3x - 3\lambda x \sin 3x$ $\frac{d^2y}{dx^2} = -3\lambda \sin 3x - 3\lambda \sin 3x - 9\lambda x \cos 3x$ $\therefore -6\lambda \sin 3x - 9\lambda x \cos 3x + 9\lambda x \cos 3x = -12 \sin 3x$ $\lambda = 2$	M1 A1 A1 A1 cso A1 (4)
(b)	$\lambda^2 - 9 = 0$ $\lambda = (\pm)3i$ $\therefore y = A \sin 3x + B \cos 3x$ $\therefore y = A \sin 3x + B \cos 3x + 2x \cos 3x$	M1 A1 form M1 A1 ft on $\lambda$ 's (4)
(c)	$y = 1, x = 0 \Rightarrow B = 1$ $\frac{dy}{dx} = 3A \cos 3x - 3B \sin 3x + 2 \cos 3x - 6x \sin 3x$ $2 = 3A + 2 \Rightarrow A = 0$ $\therefore y = \cos 3x + 2x \cos 3x$	B1 M1 A1 ft on $\lambda$ 's A1 (4)
(d)		axes shape B1 B1 (2)  <b>(14 marks)</b>

**EDEXCEL PURE MATHEMATICS P4 (6671) – JANUARY 2003 PROVISIONAL MARK SCHEME**

Question Number	Scheme	Marks
8. (a)	$\frac{1}{2}a^2 \int 1 + \cos^2 \theta + 2 \cos \theta \ d\theta$ $= \frac{1}{2}a^2 \int 1 + \frac{\cos 2\theta + 1}{2} + 2 \cos \theta \ d\theta$ $= 2 \times \frac{1}{2}a^2 \left[ \theta + \frac{\sin 2\theta}{4} + \frac{\theta}{2} + 2 \sin \theta \right]_0^\pi$ $= a^2 \left[ \frac{3\pi}{2} \right] = \frac{3\pi a^2}{2}$	M1 A1 correct with limits M1 A1 A1 A1 (6)
(b)	$x = a \cos \theta + a \cos^2 \theta$ $\frac{dx}{d\theta} = -a \sin \theta - 2a \cos \theta \sin \theta$ $\frac{dx}{d\theta} = 0 \Rightarrow \cos \theta = -\frac{1}{2}$ $\theta = \frac{2\pi}{3} \text{ or } \theta = \frac{4\pi}{3}$ $r = \frac{a}{2} \text{ or } r = \frac{a}{2}$ $A: r = \frac{a}{2}, \theta = \frac{2\pi}{3}$ $B: r = \frac{a}{2}, \theta = \frac{-2\pi}{3}$	$r \cos \theta$ M1 A1 finding $\theta$ M1 finding $r$ M1 both $A$ and $B$ A1 (5)
(c)	$x = -\frac{1}{4}a \quad \therefore WX = 2a + \frac{1}{4}a = 2\frac{1}{4}a$	M1 A1
(d)	$WXYZ = \frac{27\sqrt{3}a^2}{8}$	B1 ft (1)
(e)	$\text{Area} = \frac{27\sqrt{3}}{8} \times 100 - \frac{3\pi \times 100}{2} = 113.3 \text{ cm}^2$	M1 A1 (2)  <b>(16 marks)</b>