

Paper Reference(s)

**6673**

**Edexcel GCE**

**Pure Mathematics P3**

**Advanced/Advanced Subsidiary**

**Thursday 9 January 2003 – Afternoon**

**Time: 1 hour 30 minutes**

**Materials required for examination**

Answer Book (AB16)  
Mathematical Formulae (Lilac)  
Graph Paper (ASG2)

**Items included with question papers**

Nil

**Candidates may only use one of the basic scientific calculators approved by the Qualifications and Curriculum Authority.**

**Instructions to Candidates**

---

In the boxes on the answer book, write the name of the examining body (Edexcel), your centre number, candidate number, the unit title (Pure Mathematics P3), the paper reference (6673), your surname, other name and signature.

When a calculator is used, the answer should be given to an appropriate degree of accuracy.

**Information for Candidates**

---

A booklet ‘Mathematical Formulae and Statistical Tables’ is provided.

Full marks may be obtained for answers to ALL questions.

This paper has eight questions. Pages 6, 7 and 8 are blank.

**Advice to Candidates**

---

You must ensure that your answers to parts of questions are clearly labelled.

You must show sufficient working to make your methods clear to the Examiner. Answers without working may gain no credit.

1. The function  $f$  is given by

$$f(x) = \frac{3(x+1)}{(x+2)(x-1)}, \quad x \in \mathbb{R}, \quad x \neq -2, \quad x \neq 1.$$

(a) Express  $f(x)$  in partial fractions.


**(3)**

(b) Hence, or otherwise, prove that  $f'(x) < 0$  for all values of  $x$  in the domain.

**(3)**

2.

**Figure 1**



The circle  $C$ , with centre  $(a, b)$  and radius 5, touches the  $x$ -axis at  $(4, 0)$ , as shown in Fig. 1.

(a) Write down the value of  $a$  and the value of  $b$ .

**(1)**

(b) Find a cartesian equation of  $C$ .

**(2)**

A tangent to the circle, drawn from the point  $P(8, 17)$ , touches the circle at  $T$ .

(c) Find, to 3 significant figures, the length of  $PT$ .

**(3)**

3.  $f(n) = n^3 + pn^2 + 11n + 9$ , where  $p$  is a constant.

(a) Given that  $f(n)$  has a remainder of 3 when it is divided by  $(n + 2)$ , prove that  $p = 6$ . (2)

(b) Show that  $f(n)$  can be written in the form  $(n + 2)(n + q)(n + r) + 3$ , where  $q$  and  $r$  are integers to be found. (3)

(c) Hence show that  $f(n)$  is divisible by 3 for all positive integer values of  $n$ . (2)

---

4. (a) Expand  $(1 + 3x)^{-2}$ ,  $|x| < \frac{1}{3}$ , in ascending powers of  $x$  up to and including the term in  $x^3$ , simplifying each term. (4)

(b) Hence, or otherwise, find the first three terms in the expansion of  $\frac{x + 4}{(1 + 3x)^2}$  as a series in ascending powers of  $x$ . (4)

---

5. Liquid is poured into a container at a constant rate of  $30 \text{ cm}^3 \text{ s}^{-1}$ . At time  $t$  seconds liquid is leaking from the container at a rate of  $\frac{2}{15} V \text{ cm}^3 \text{ s}^{-1}$ , where  $V \text{ cm}^3$  is the volume of liquid in the container at that time.

(a) Show that

$$-15 \frac{dV}{dt} = 2V - 450. \quad (3)$$

Given that  $V = 1000$  when  $t = 0$ ,

(b) find the solution of the differential equation, in the form  $V = f(t)$ . (7)

(c) Find the limiting value of  $V$  as  $t \rightarrow \infty$ . (1)

---

6. Referred to a fixed origin  $O$ , the points  $A$  and  $B$  have position vectors  $(\mathbf{i} + 2\mathbf{j} - 3\mathbf{k})$  and  $(5\mathbf{i} - 3\mathbf{j})$  respectively.

(a) Find, in vector form, an equation of the line  $l_1$  which passes through  $A$  and  $B$ . (2)

The line  $l_2$  has equation  $\mathbf{r} = (4\mathbf{i} - 4\mathbf{j} + 3\mathbf{k}) + \mu(\mathbf{i} - 2\mathbf{j} + 2\mathbf{k})$ , where  $\mu$  is a scalar parameter.

(b) Show that  $A$  lies on  $l_2$ . (1)

(c) Find, in degrees, the acute angle between the lines  $l_1$  and  $l_2$ . (4)

The point  $C$  with position vector  $(2\mathbf{i} - \mathbf{k})$  lies on  $l_2$ .

(d) Find the shortest distance from  $C$  to the line  $l_1$ . (4)

7.

**Figure 2**




Figure 2 shows the curve with equation  $y = x^{\frac{1}{2}} e^{-2x}$ .

(a) Find the  $x$ -coordinate of  $M$ , the maximum point of the curve. (5)

The finite region enclosed by the curve, the  $x$ -axis and the line  $x = 1$  is rotated through  $2\pi$  about the  $x$ -axis.

(b) Find, in terms of  $\pi$  and  $e$ , the volume of the solid generated. (7)

8. (a) Use the identity for  $\cos(A + B)$  to prove that  $\cos 2A = 2 \cos^2 A - 1$ . (2)

(b) Use the substitution  $x = 2\sqrt{2} \sin \theta$  to prove that

$$\int_2^{\sqrt{6}} \sqrt{8 - x^2} \, dx = \frac{1}{3}(\pi + 3\sqrt{3} - 6).$$
(7)

A curve is given by the parametric equations

$$x = \sec \theta, \quad y = \ln(1 + \cos 2\theta), \quad 0 \leq \theta < \frac{\pi}{2}.$$

- (c) Find an equation of the tangent to the curve at the point where  $\theta = \frac{\pi}{3}$ . (5)

---

**END**