







Question Number	Scheme	Marks
1. (a)	 $v_A = v_{AW} + v_W$ $\Rightarrow \begin{pmatrix} k \\ 0 \end{pmatrix} = \begin{pmatrix} v \cos \phi \\ -v \sin \phi \end{pmatrix} + \begin{pmatrix} w \cos \theta \\ w \sin \theta \end{pmatrix}$ $\Rightarrow v \sin \phi = w \sin \theta *$	M1
(b)	$k = v \sin \phi + w \sin \theta$ $= \frac{v \sqrt{v^2 - w^2 \sin^2 \theta}}{v} + w \cos \theta$ $= \sqrt{v^2 - w^2 \sin^2 \theta} + w \cos \theta$	A1 (2) M1 A1 M1 A1 (4)
		(6 marks)
2.	 $\sin \theta = \frac{a}{2a}$ $\Rightarrow \theta = 30^\circ$ $v_1 + v_2 = u \cos \theta$ $-v_1 + v_2 = eu \cos \theta$ $\frac{u \sin \theta}{v_1} = \tan (\theta + 30^\circ) \text{ (or equivalent)}$ <p>Producing an equation in <math>e</math> only</p> $e = \frac{1}{3}$	M1 A1 M1 A1 M1 A1 M1 A1 M1 A1 c.s.o  (11 marks)

Question Number	Scheme	Marks
3. (a)	 <p>Vector <math>\Delta</math></p> $\cos \theta = \frac{12}{13}$ $(\theta = 22.6^\circ)$ <p>Course is <math>360^\circ - 22.6^\circ</math></p> $= 337^\circ \text{ (AWRT)}$	M1 A1 M1 A1 (4)
(b)	$v = \sqrt{13^2 - 12^2} = 5$ $t = \frac{6 \cos \theta}{5} = 1.107$ <p>Time is 1.06 p.m.</p>	
(c)	$d = 6 \sin \theta = 6 \times \frac{5}{13} = 2.31 \text{ km}$ <p>(AWRT 2.3 km)</p>	M1 A1 (2) <b>(11 marks)</b>

Question Number	Scheme	Marks
4. (a)	 $OQ = 4a \sin \theta$ $V = (-)mga \sin 2\theta + \frac{mg}{2\sqrt{3}2a}(4a \sin \theta - a)^2 + C$ $= -mga \sin 2\theta + \frac{mga^2}{4a\sqrt{3}}(16 \sin^2 \theta - 8 \sin \theta + 1) + C$ $= -mga \sin 2\theta + \frac{mga}{4\sqrt{3}}(8(1 - \cos 2\theta) - 8 \sin \theta) + C$	B1 B1; M1 A1 M1 M1
	i.e. $V = -\frac{mga}{\sqrt{3}}(2 \cos 2\theta + \sqrt{3} \sin 2\theta + 2 \sin \theta) + C$ *	A1 c.s.o (7)
(b)	$V'(\theta) = -\frac{mga}{\sqrt{3}}(-4 \sin 2\theta + 2\sqrt{3} \cos 2\theta + 2 \cos \theta)$	M1 A1
	$V'\left(\frac{\pi}{6}\right) = -\frac{mga}{\sqrt{3}}(-2\sqrt{3} + 2\sqrt{3}\frac{1}{2} + 2\frac{\sqrt{3}}{2}) = 0$	M1 A1 (4)
(c)	$V''(\theta) = \frac{mga}{\sqrt{3}}(+8 \cos 2\theta + 4\sqrt{3} \sin 2\theta + 2 \sin \theta)$	M1 A1
	Hence, $V''\left(\frac{\pi}{6}\right) = \frac{11mga}{\sqrt{3}} > 0 \therefore$ stable	M1 A1 c.s.o (4)
		<b>(15 marks)</b>

Question Number	Scheme	Marks
5. (a)	 <p> <math>\mathbf{R}(\downarrow), mg - 2k\dot{x}m - T = m\ddot{x}</math>  <math>g - 2k\dot{x} - \frac{2ak^2x}{a} = \ddot{x}</math>  <math>\Rightarrow \ddot{x} + 2k\dot{x} + 2k^2x = g \quad *</math> </p>	M1 A1 c.s.o (4)
(b)	$t = 0, x = a: \quad a = D + \frac{g}{2k^2} \Rightarrow D = a - \frac{g}{2k^2}$ $\dot{x} = -ke^{-kt}(C \sin kt + D \cos kt) + -ke^{-kt}(C \cos kt - D \sin kt)$ $t = 0, \dot{x} = 0: \quad 0 = -kD + kC \Rightarrow C = D$ $\Rightarrow C = a - \frac{g}{2k^2}$	M1 A1 M1 A1 A1 (5)
(c)	$\dot{x} = 0 \Rightarrow C(\sin kt + \cos kt) + C(\cos kt - \sin kt)$ $\Rightarrow \sin kt = 0$ $\Rightarrow kt = \pi$ $\Rightarrow t = \frac{\pi}{k}$	M1 A1 ft A1 (3)
(d)	When $t = \frac{\pi}{k}, x = -De^{-\pi} + \frac{g}{2k^2}$ $\frac{g}{2k^2} - e^{-\pi}(a - \frac{g}{2k^2})$ $\Rightarrow xe^\pi = \frac{g}{2k^2}(e^\pi + 1) - a$ $> 0$ (given) $\Rightarrow g(e^\pi + 1) > 2k^2a \quad *$	M1 A1 ft M1 A1 c.s.o (4) <b>(16 marks)</b>

Question Number	Scheme	Marks
6. (a)	 $F = \frac{kmg}{v}$ $\text{R}(\uparrow), F - mg = ma$ $\frac{kmg}{v} - mg = mv \frac{dv}{dx}$ $g(k - v) = v^2 \frac{dv}{dx} \quad *$	B1 M1      M1 A1      (4)
(b)	$g dx = \frac{v^2}{k-v} dv$ $\int g dx = \int -v - k + \frac{k^2}{k-v} dv$ $gx = -\frac{v^2}{2} - kv - k^2 \ln(k-v) + (c)$	M1 M1 A1 M1 A1
	$x = 0, v = 0$ $0 = 0 - 0 - k^2 \ln k + c$ $c = k^2 \ln k$ $gx = -\frac{v^2}{2} - kv - k^2 \ln \left( \frac{k}{k-v} \right) \quad *$	M1 M1 A1      (7)
(c)	<p>Work done by engine      =      Energy gain</p> $kmgt = \frac{1}{2} mv^2 + mgx$ $kmgt = mk^2 \ln \left( \frac{k}{k-v} \right) - mkv$ $\Rightarrow t = \frac{k}{g} \ln \left( \frac{k}{k-v} \right) - \frac{v}{g}$	M1 A1 M1 A1      (5)
		<b>(16 marks)</b>