

6665 Core C3
Mark Scheme (Post standardisation)

Question Number	Scheme	Marks
1	(a) Dividing by $\cos^2 \theta$: $\frac{\sin^2 \theta}{\cos^2 \theta} + \frac{\cos^2 \theta}{\cos^2 \theta} \equiv \frac{1}{\cos^2 \theta}$ Completion : $1 + \tan^2 \theta \equiv \sec^2 \theta$ (no errors seen)	M1 A1 (2)
	(b) Use of $1 + \tan^2 \theta = \sec^2 \theta$: $2(\sec^2 \theta - 1) + \sec \theta = 1$ $[2\sec^2 \theta + \sec \theta - 3 = 0]$ Factorising or solving: $(2\sec \theta + 3)(\sec \theta - 1) = 0$ $[\sec \theta = -\frac{3}{2} \text{ or } \sec \theta = 1]$ $\theta = 0$ $\cos \theta = -\frac{2}{3}$; $\theta_1 = 131.8^\circ$ $\theta_2 = 228.2^\circ$ $[A1ft \text{ for } \theta_2 = 360^\circ - \theta_1]$	M1 M1 B1 M1 A1 A1√ (6)
		[8]

6665 Core C3
Mark Scheme (Post standardisation)

Question Number	Scheme	Marks
2	<p>(a) (i) $6 \sin x \cos x + 2 \sec 2x \tan 2x$</p> <p>or $3 \sin 2x + 2 \sec 2x \tan 2x$ [M1 for $6 \sin x$]</p> <p>(ii) $3(x + \ln 2x)^2 \left(1 + \frac{1}{x}\right)$ [B1 for $3(x + \ln 2x)^2$]</p> <p>(b) Differentiating numerator to obtain $10x - 10$</p> <p style="padding-left: 40px;">Differentiating denominator to obtain $2(x-1)$</p> <p>Using quotient rule formula correctly:</p> <p>To obtain $\frac{dy}{dx} = \frac{(x-1)^2(10x-10) - (5x^2-10x+9)2(x-1)}{(x-1)^4}$</p> <p>Simplifying to form $\frac{2(x-1)[5(x-1)^2 - (5x^2-10x+9)]}{(x-1)^4}$</p> <p>$= -\frac{8}{(x-1)^3}$ * (c.s.o.)</p>	<p>M1A1A1 (3)</p> <p>B1M1A1 (3)</p>
3	<p>(a) $\frac{5x+1}{(x+2)(x-1)} - \frac{3}{x+2}$</p> <p style="padding-left: 40px;">$= \frac{5x+1-3(x-1)}{(x+2)(x-1)}$</p> <p style="padding-left: 40px;">M1 for combining fractions even if the denominator is not lowest common</p> <p>$= \frac{2x+4}{(x+2)(x-1)} = \frac{2(x+2)}{(x+2)(x-1)} = \frac{2}{x-1}$ *</p> <p style="padding-left: 100px;">M1 must have linear numerator</p> <p>(b) $y = \frac{2}{x-1} \Rightarrow xy - y = 2 \Rightarrow xy = 2 + y$</p> <p>$f^{-1}(x) = \frac{2+x}{x}$ o.e.</p> <p>$fg(x) = \frac{2}{x^2+4}$ (attempt) [$\frac{2}{"g"-1}$]</p> <p>Setting $\frac{2}{x^2+4} = \frac{1}{4}$ and finding $x^2 = \dots$; $x = \pm 2$</p>	<p>B1</p> <p>M1</p> <p>M1 A1 cso (4)</p> <p>M1A1</p> <p>A1 (3)</p> <p>M1</p> <p>M1; A1 (3)</p> <p style="text-align: right;">[10]</p>



6665 Core C3
Mark Scheme (Post standardisation)

Question Number	Scheme	Marks
4	<p>(a) $f'(x) = 3e^x - \frac{1}{2x}$</p> <p>$3e^x - \frac{1}{2x} = 0$</p> <p>$\Rightarrow 6\alpha e^\alpha = 1 \quad \Rightarrow \alpha = \frac{1}{6} e^{-\alpha} \quad (*)$</p>	<p>M1A1A1 (3)</p> <p>M1</p> <p>A1 cso (2)</p>
	<p>(c) $x_1 = 0.0613\dots, x_2 = 0.1568\dots, x_3 = 0.1425\dots, x_4 = 0.1445\dots$</p> <p>[M1 at least x_1 correct, A1 all correct to 4 d.p.]</p>	<p>M1 A1 (2)</p>
	<p>(d) Using $f'(x) = 3e^x - \frac{1}{2x}$ with suitable interval</p> <p>e.g. $f'(0.14425) = -0.0007$</p> <p>$f'(0.14435) = +0.002(1)$</p> <p>Accuracy (change of sign and correct values)</p>	<p>M1</p> <p>A1 (2)</p>
		[9]

6665 Core C3
Mark Scheme (Post standardisation)

Question Number	Scheme	Marks
5	(a) $\cos 2A = \cos^2 A - \sin^2 A$ (+ use of $\cos^2 A + \sin^2 A \equiv 1$)	M1
	$= (1 - \sin^2 A) - \sin^2 A = 1 - 2 \sin^2 A$ (*)	A1 (2)
	(b) $2 \sin 2\theta - 3 \cos 2\theta - 3 \sin \theta + 3 \equiv 4 \sin \theta \cos \theta; -3(1 - 2 \sin^2 \theta) - 3 \sin \theta + 3$	B1; M1
	$\equiv 4 \sin \theta \cos \theta + 6 \sin^2 \theta - 3 \sin \theta$	M1
	$\equiv \sin \theta (4 \cos \theta + 6 \sin \theta - 3)$ (*)	A1 (4)
	(c) $4 \cos \theta + 6 \sin \theta \equiv R \sin \theta \cos \alpha + R \cos \theta \sin \alpha$	
	Complete method for R (may be implied by correct answer)	
	$[R^2 = 4^2 + 6^2, R \sin \alpha = 4, R \cos \alpha = 6]$	M1
	$R = \sqrt{52}$ or 7.21	A1
	Complete method for α ; $\alpha = 0.588$ (allow 33.7°)	M1 A1 (4)
	(d) $\sin \theta (4 \cos \theta + 6 \sin \theta - 3) = 0$	M1
	$\theta = 0$	B1
$\sin(\theta + 0.588) = \frac{3}{\sqrt{52}} = 0.4160..$ (24.6°)	M1	
$\theta + 0.588 = (0.4291), 2.7125$ [or $\theta + 33.7^\circ = (24.6^\circ), 155.4^\circ$]	dM1	
$\theta = 2.12$ cao	A1 (5)	
	[15]	

6665 Core C3
Mark Scheme (Post standardisation)

Question Number	Scheme	Marks
6. (a)	 <p style="text-align: center;">Translation ← by 1</p> <p style="text-align: center;">Intercepts correct</p>	M1 A1 (2)
(b)	 <p style="text-align: center;">$x \geq 0$, correct “shape”</p> <p style="text-align: center;">[provided not just original]</p> <p style="text-align: center;">Reflection in y-axis</p> <p style="text-align: center;">Intercepts correct</p>	B1 B1√ B1 (3)
(c)	$a = -2, b = -1$	B1 B1 (2)
(d)	<p>Intersection of $y = 5x$ with $y = -x - 1$</p> <p style="text-align: center;">Solving to give $x = -\frac{1}{6}$</p>	M1A1 M1A1 (4)
	<p>[Notes:</p> <p>(i) If both values found for $5x = -x - 1$ and $5x = x - 3$, or solved algebraically, can score 3 out of 4 for $x = -\frac{1}{6}$ and $x = -\frac{3}{4}$; required to eliminate $x = -\frac{3}{4}$ for final mark.</p> <p>(ii) Squaring approach: M1 correct method, $24x^2 + 22x + 3 = 0$ (correct 3 term quadratic, any form) A1 Solving M1, Final correct answer A1.]</p>	[11]

6665 Core C3
Mark Scheme (Post standardisation)

Question Number	Scheme	Marks
7	(a) Setting $p = 300$ at $t = 0 \Rightarrow 300 = \frac{2800a}{1+a}$ $(300 = 2500a); \quad a = 0.12$ (c.s.o) *	M1 dM1A1 (3)
	(b) $1850 = \frac{2800(0.12)e^{0.2t}}{1 + 0.12e^{0.2t}} ; \quad e^{0.2t} = 16.2\dots$ Correctly taking logs to $0.2 t = \ln k$	M1A1 M1
	$t = 14$ (13.9..)	A1 (4)
	(c) Correct derivation: (Showing division of num. and den. by $e^{0.2t}$; using a)	B1 (1)
	(d) Using $t \rightarrow \infty, e^{-0.2t} \rightarrow 0,$ $p \rightarrow \frac{336}{0.12} = 2800$	M1 A1 (2)
		[10]