

GCE

Edexcel GCE

Core Mathematics C2 (6664)

Summer 2005

advancing learning, changing lives

Mark Scheme (Results)

Edexcel GCE
Core Mathematics C2 (6664)

June 2005
6664 Core Mathematics C2
Mark Scheme

Question number	Scheme	Marks
1.	$\frac{dy}{dx} = 4x - 12$ $4x - 12 = 0 \quad x = 3$ $y = -18$	B1 M1 A1ft A1 (4) 4
	<p>M1: Equate $\frac{dy}{dx}$ (not just y) to zero and proceed to $x = \dots$ A1ft: Follow through only from a linear equation in x.</p> <p><u>Alternative:</u> $y = 2x(x - 6) \Rightarrow$ Curve crosses x-axis at 0 and 6 B1 (By symmetry) $x = 3$ M1 A1ft $y = -18$ A1</p> <p><u>Alternative:</u> $(x - 3)^2$ B1 for $(x - 3)^2$ seen somewhere $y = 2(x^2 - 6x) = 2\{(x - 3)^2 - 9\}$ $x = 3$ M1 for attempt to complete square and deduce $x = \dots$ A1ft [$(x - a)^2 \Rightarrow x = a$] $y = -18$ A1</p>	

Question number	Scheme	Marks
2.	<p>(a) $x \log 5 = \log 8, \quad x = \frac{\log 8}{\log 5}, \quad = 1.29$</p> <p>(b) $\log_2 \frac{x+1}{x} \quad (\text{or } \log_2 7x)$</p> <p>$\frac{x+1}{x} = 7 \quad x = \dots, \quad \frac{1}{6} \quad (\text{Allow } 0.167 \text{ or better})$</p>	<p>M1, A1, A1 (3)</p> <p>B1</p> <p>M1, A1 (3)</p> <p>6</p>
	<p>(a) Answer only 1.29 : Full marks.</p> <p>Answer only, which rounds to 1.29 (e.g. 1.292): M1 A1 A0</p> <p>Answer only, which rounds to 1.3 : M1 A0 A0</p> <p>Trial and improvement: Award marks as for “answer only”.</p> <p>(b) M1: Form (by legitimate log work) and solve an equation in x.</p> <p>Answer only: No marks unless verified (then full marks are available).</p>	

Question number	Scheme	Marks
3.	<p>(a) Attempt to evaluate $f(-4)$ or $f(4)$</p> $f(-4) = 2(-4)^3 + (-4)^2 - 25(-4) + 12 \quad (= 128 + 16 + 100 + 12) = 0,$ <p style="text-align: center;">so is a factor.</p> <p>(b) $(x + 4)(2x^2 - 7x + 3)$</p> <p style="text-align: center;">.....$(2x - 1)(x - 3)$</p>	<p>M1</p> <p>A1 (2)</p> <p>M1 A1</p> <p>M1 A1 (4)</p> <p>6</p>
	<p>(b) First M requires $(2x^2 + ax + b)$, $a \neq 0, b \neq 0$.</p> <p>Second M for the attempt to factorise the quadratic.</p> <p><u>Alternative:</u></p> $(x + 4)(2x^2 + ax + b) = 2x^3 + (8 + a)x^2 + (4a + b)x + 4b = 0,$ <p>then compare coefficients to find <u>values</u> of a and b. [M1]</p> $a = -7, b = 3 \quad [A1]$ <p><u>Alternative:</u></p> <p>Factor theorem: Finding that $f\left(\frac{1}{2}\right) = 0, \therefore (2x - 1)$ is a factor [M1, A1]</p> <p>n.b. Finding that $f\left(\frac{1}{2}\right) = 0, \therefore (x - \frac{1}{2})$ is a factor scores M1, A0, unless the factor 2 subsequently appears.</p> <p style="text-align: center;">Finding that $f(3) = 0, \therefore (x - 3)$ is a factor [M1, A1]</p>	

Question number	Scheme	Marks
4.	<p>(a) $1 + 12px, + \frac{12 \times 11}{2}(px)^2$</p> <p>(b) $12p(x) = -q(x) \quad 66p^2(x^2) = 11q(x^2) \quad (\text{Equate terms, or coefficients})$</p> <p>$\Rightarrow 66p^2 = -132p \quad (\text{Eqn. in } p \text{ or } q \text{ only})$</p> <p>$p = -2, \quad q = 24$</p>	<p>B1, B1 (2)</p> <p>M1</p> <p>M1</p> <p>A1, A1 (4)</p> <p>6</p>
	<p>(a) Terms can be listed rather than added. First B1: Simplified form must be seen, but may be in (b).</p> <p>(b) First M: May still have $\binom{12}{2}$ or ${}^{12}C_2$</p> <p>Second M: <u>Not</u> with $\binom{12}{2}$ or ${}^{12}C_2$. Dependent upon having p's in each term.</p> <p>Zero solutions must be rejected for the final A mark.</p>	

Question number	Scheme	Marks
5.	<p>(a) $(x + 10 =) \quad 60 \quad \alpha$ 120 (M: $180 - \alpha$ or $\pi - \alpha$) $x = 50 \quad x = 110$ (or 50.0 and 110.0) (M: Subtract 10)</p> <p>(b) $(2x =) \quad 154.2 \quad \beta$ Allow a.w.r.t. 154 or a.w.r.t. 2.69 (radians) 205.8 (M: $360 - \beta$ or $2\pi - \beta$) $x = 77.1 \quad x = 102.9$ (M: Divide by 2)</p>	<p>B1 M1 M1 A1 (4)</p> <p>B1 M1 M1 A1 (4)</p> <p>8</p>
	<p>(a) First M: Must be subtracting from 180 <u>before</u> subtracting 10. (b) First M: Must be subtracting from 360 <u>before</u> dividing by 2, <u>or</u> dividing by 2 then subtracting from 180.</p> <p>In each part: Extra solutions outside 0 to 180 : Ignore. Extra solutions between 0 and 180 : A0.</p> <p><u>Alternative for (b): (double angle formula)</u></p> $1 - 2\sin^2 x = -0.9 \qquad 2\sin^2 x = 1.9 \qquad \text{B1}$ $\sin x = \sqrt{0.95} \qquad \text{M1}$ $x = 77.1$ $x = 180 - 77.1 = 102.9 \qquad \text{M1 A1}$	

Question number	Scheme	Marks
7.	<p>(a) $\frac{\sin x}{8} = \frac{\sin 0.5}{7}$ or $\frac{8}{\sin x} = \frac{7}{\sin 0.5}$, $\sin x = \frac{8 \sin 0.5}{7}$ $\sin x = 0.548$</p> <p>(b) $x = 0.58$ (α) (This mark may be earned in (a)). $\pi - \alpha = 2.56$</p>	<p>M1 A1ft A1 (3) B1 M1 A1ft (3) 6</p>
	<p>(a) M: Sine rule attempt (sides/angles possibly the “wrong way round”). A1ft: follow through from sides/angles are the “wrong way round”.</p> <p><u>Too many d.p. given:</u> Maximum 1 mark penalty in the complete question. (Deduct on first occurrence).</p>	

Question number	Scheme	Marks
8.	<p>(a) Centre (5, 0) (or $x = 5, y = 0$)</p> <p>(b) $(x \pm a)^2 \pm b \pm 9 + (y \pm c)^2 = 0 \Rightarrow r^2 = \dots$ or $r = \dots$, Radius = 4</p> <p>(c) (1, 0), (9, 0) Allow just $x = 1, x = 9$</p> <p>(d) Gradient of $AT = -\frac{2}{7}$</p> $y = -\frac{2}{7}(x - 5)$	<p>B1 B1 (2)</p> <p>M1, A1 (2)</p> <p>B1ft, B1ft (2)</p> <p>B1</p> <p>M1 A1ft (3)</p> <p style="text-align: right;">9</p>
	<p>(a) (0, 5) scores B1 B0.</p> <p>(d) M1: Equation of straight line through centre, <u>any</u> gradient (except 0 or ∞) (The equation can be in any form).</p> <p>A1ft: Follow through from centre, but gradient must be $-\frac{2}{7}$.</p>	

Question number	Scheme	Marks
9.	<p>(a) ($S =$) $a + ar + \dots + ar^{n-1}$ “$S =$” not required. Addition required.</p> <p>($rS =$) $ar + ar^2 + \dots + ar^n$ “$rS =$” not required (M: Multiply by r)</p> <p>$S(1-r) = a(1-r^n)$ $S = \frac{a(1-r^n)}{1-r}$ (M: Subtract and factorise) (*)</p> <p>(b) $ar^{n-1} = 35000 \times 1.04^3 = 39400$ (M: Correct a and r, with $n = 3, 4$ or 5).</p> <p>(c) $n = 20$ (Seen or implied)</p> <p>$S_{20} = \frac{35000(1-1.04^{20})}{(1-1.04)}$</p> <p>(M1: Needs <u>any</u> r value, $a = 35000$, $n = 19, 20$ or 21).</p> <p>(A1ft: ft from $n = 19$ or $n = 21$, but r must be 1.04).</p> <p>$= 1\,042\,000$</p>	<p>B1</p> <p>M1</p> <p>M1 A1cso (4)</p> <p>M1 A1 (2)</p> <p>B1</p> <p>M1 A1ft</p> <p>A1 (4)</p> <p>10</p>
	<p>(a) B1: At least the 3 terms shown above, and no extra terms. A1: Requires a completely correct solution. <u>Alternative for the 2 M marks:</u> M1: Multiply numerator and denominator by $1 - r$. M1: Multiply out numerator convincingly, and factorise.</p> <p>(b) M1 can also be scored by a “year by year” method. <u>Answer only:</u> 39 400 scores full marks, 39 370 scores M1 A0.</p> <p>(c) M1 can also be scored by a “year by year” method, <u>with terms added</u>. In this case the B1 will be scored if the correct number of years is considered. <u>Answer only:</u> Special case: 1 042 000 scores 2 B marks, scored as 1, 0, 0, 1 (Other answers score no marks).</p> <p><u>Failure to round correctly in (b) and (c):</u> Penalise once only (first occurrence).</p>	

Question number	Scheme	Marks
10.	<p>(a) $\int (2x + 8x^{-2} - 5)dx = x^2 + \frac{8x^{-1}}{-1} - 5x$</p> $\left[x^2 + \frac{8x^{-1}}{-1} - 5x \right]_1^4 = (16 - 2 - 20) - (1 - 8 - 5) \quad (= 6)$ <p>$x = 1: y = 5$ and $x = 4: y = 3.5$</p> <p>Area of trapezium = $\frac{1}{2}(5 + 3.5)(4 - 1) \quad (= 12.75)$</p> <p>Shaded area = $12.75 - 6 = 6.75 \quad (\text{M: Subtract either way round})$</p> <p>(b) $\frac{dy}{dx} = 2 - 16x^{-3}$</p> <p>(Increasing where) $\frac{dy}{dx} > 0; \quad \text{For } x > 2, \frac{16}{x^3} < 2, \therefore \frac{dy}{dx} > 0 \quad (\text{Allow } \geq)$</p>	<p>M1 A1 A1</p> <p>M1</p> <p>B1</p> <p>M1</p> <p>M1 A1 (8)</p> <p>M1 A1</p> <p>dM1; A1 (4)</p> <p>12</p>
	<p>(a) Integration: One term wrong M1 A1 A0; two terms wrong M1 A0 A0. Limits: M1 for substituting limits 4 and 1 into a changed function, and subtracting the right way round.</p> <p><u>Alternative:</u></p> <p>$x = 1: y = 5$ and $x = 4: y = 3.5$</p> <p>Equation of line: $y - 5 = -\frac{1}{2}(x - 1) \quad y = \frac{11}{2} - \frac{1}{2}x$, subsequently used in integration with limits.</p> $\left(\frac{11}{2} - \frac{1}{2}x \right) - \left(2x + \frac{8}{x^2} - 5 \right) \quad (\text{M: Subtract either way round})$ $\int \left(\frac{21}{2} - \frac{5x}{2} - 8x^{-2} \right) dx = \frac{21x}{2} - \frac{5x^2}{4} - \frac{8x^{-1}}{-1}$ <p>(Penalise integration mistakes, not algebra for the ft marks)</p> $\left[\frac{21x}{2} - \frac{5x^2}{4} - \frac{8x^{-1}}{-1} \right]_1^4 = (42 - 20 + 2) - \left(\frac{21}{2} - \frac{5}{4} + 8 \right) \quad (\text{M: Right way round})$ <p>Shaded area = 6.75</p> <p>(The follow through marks are for the subtracted version, and again deduct an accuracy mark for a wrong term: One wrong M1 A1 A0; two wrong M1 A0 A0.)</p> <p><u>Alternative for the last 2 marks in (b):</u></p> <p>M1: Show that $x = 2$ is a minimum, using, e.g., 2nd derivative.</p> <p>A1: Conclusion showing understanding of “increasing”, with accurate working.</p>	<p>B1</p> <p>3rd M1</p> <p>4th M1</p> <p>1st M1 A1ft A1ft</p> <p>2nd M1</p> <p>A1</p>