

**6665 Core C3**  
**Mark Scheme (Post standardisation)**

Question Number	Scheme	Marks
1 (a)	Dividing by $\cos^2 \theta$ : $\frac{\sin^2 \theta}{\cos^2 \theta} + \frac{\cos^2 \theta}{\cos^2 \theta} \equiv \frac{1}{\cos^2 \theta}$ Completion : $1 + \tan^2 \theta \equiv \sec^2 \theta$ (no errors seen)	M1  A1 (2)
(b)	Use of $1 + \tan^2 \theta = \sec^2 \theta$ : $2(\sec^2 \theta - 1) + \sec \theta = 1$ $[2\sec^2 \theta + \sec \theta - 3 = 0]$ Factorising or solving: $(2\sec \theta + 3)(\sec \theta - 1) = 0$ $[\sec \theta = -\frac{3}{2} \text{ or } \sec \theta = 1]$ $\theta = 0$ $\cos \theta = -\frac{2}{3}; \theta_1 = 131.8^\circ$ $\theta_2 = 228.2^\circ$ [A1ft for $\theta_2 = 360^\circ - \theta_1$ ]	M1  M1  B1  M1 A1  A1✓ (6)  [8]

**6665 Core C3**  
**Mark Scheme (Post standardisation)**

Question Number	Scheme	Marks
2 (a)	(i) $6\sin x \cos x + 2\sec 2x \tan 2x$ <b>or</b> $3 \sin 2x + 2\sec 2x \tan 2x$ [M1 for $6 \sin x$ ] (ii) $3(x + \ln 2x)^2(1 + \frac{1}{x})$ [B1 for $3(x + \ln 2x)^2$ ]	M1A1A1 (3)  B1M1A1 (3)
(b)	Differentiating numerator to obtain $10x - 10$ Differentiating denominator to obtain $2(x-1)$ Using quotient rule formula correctly: To obtain $\frac{dy}{dx} = \frac{(x-1)^2(10x-10) - (5x^2 - 10x + 9)2(x-1)}{(x-1)^4}$ Simplifying to form $\frac{2(x-1)[5(x-1)^2 - (5x^2 - 10x + 9)]}{(x-1)^4}$ $= -\frac{8}{(x-1)^3}$ * (c.s.o.)	
3 (a)	$\begin{aligned} & \frac{5x+1}{(x+2)(x-1)} - \frac{3}{x+2} \\ &= \frac{5x+1-3(x-1)}{(x+2)(x-1)} \end{aligned}$ M1 for combining fractions even if the denominator is not lowest common $= \frac{2x+4}{(x+2)(x-1)} = \frac{2(x+2)}{(x+2)(x-1)} = \frac{2}{x-1}$ * M1 must have linear numerator	B1  M1  M1 A1 cso (4)
(b)	$y = \frac{2}{x-1} \Rightarrow xy - y = 2 \Rightarrow xy = 2 + y$ $f^{-1}(x) = \frac{2+x}{x}$ o.e. $fg(x) = \frac{2}{x^2+4}$ (attempt) [ $\frac{2}{"g"-1}$ ] Setting $\frac{2}{x^2+4} = \frac{1}{4}$ and finding $x^2 = \dots$ ; $x = \pm 2$	M1A1  A1 (3)  M1  M1; A1 (3)  [10]

**6665 Core C3**  
**Mark Scheme (Post standardisation)**

Question Number	Scheme	Marks
4 (a)	$f'(x) = 3e^x - \frac{1}{2x}$ $3e^x - \frac{1}{2x} = 0$ $\Rightarrow 6\alpha e^\alpha = 1 \quad \Rightarrow \alpha = \frac{1}{6} e^{-\alpha} \quad (*)$	M1A1A1 (3)
(c)	$x_1 = 0.0613\dots, x_2 = 0.1568\dots, x_3 = 0.1425\dots, x_4 = 0.1445\dots$ [ M1 at least $x_1$ correct , A1 all correct to 4 d.p. ]	M1 A1 (2)
(d)	Using $f'(x) = 3e^x - \frac{1}{2x}$ with suitable interval e.g. $f'(0.14425) = -0.0007$ $f'(0.14435) = +0.002(1)$ Accuracy (change of sign and correct values)	M1 A1 (2)  [9]

**6665 Core C3**  
**Mark Scheme (Post standardisation)**

Question Number	Scheme	Marks
5 (a)	$\cos 2A = \cos^2 A - \sin^2 A$ (+ use of $\cos^2 A + \sin^2 A \equiv 1$ ) $= (1 - \sin^2 A); -\sin^2 A = 1 - 2\sin^2 A$ (*)	M1  A1 (2)
(b)	$2\sin 2\theta - 3\cos 2\theta - 3\sin \theta + 3 \equiv 4\sin \theta \cos \theta; -3(1 - 2\sin^2 \theta) - 3\sin \theta + 3$ $\equiv 4\sin \theta \cos \theta + 6\sin^2 \theta - 3\sin \theta$ $\equiv \sin \theta(4\cos \theta + 6\sin \theta - 3)$ (*)	B1; M1  M1  A1 (4)
(c)	$4\cos \theta + 6\sin \theta \equiv R \sin \theta \cos \alpha + R \cos \theta \sin \alpha$  Complete method for $R$ (may be implied by correct answer)  $[R^2 = 4^2 + 6^2, R \sin \alpha = 4, R \cos \alpha = 6]$  $R = \sqrt{52}$ or 7.21	M1  A1  M1 A1 (4)
(d)	$\sin \theta (4\cos \theta + 6\sin \theta - 3) = 0$  $\theta = 0$  $\sin(\theta + 0.588) = \frac{3}{\sqrt{52}} = 0.4160..$ ( $24.6^\circ$ ) $\theta + 0.588 = (0.4291), 2.7125$ [or $\theta + 33.7^\circ = (24.6^\circ), 155.4^\circ$ ]  $\theta = 2.12$ cao	M1  B1  M1  dM1  A1 (5)

**6665 Core C3**  
**Mark Scheme (Post standardisation)**

Question Number	Scheme	Marks
6. (a)	 Translation $\leftarrow$ by 1 Intercepts correct	M1 A1 (2)
(b)	 $x \geq 0$ , correct “shape” [provided not just original] Reflection in $y$ -axis Intercepts correct	B1 B1 B1 (3)
(c)	$a = -2, b = -1$	B1 B1 (2)
(d)	Intersection of $y = 5x$ with $y = -x - 1$  Solving to give $x = -\frac{1}{6}$	M1A1 M1A1 (4)
		[11]
[Notes: (i) If both values found for $5x = -x - 1$ and $5x = x - 3$ , or solved algebraically, can score 3 out of 4 for $x = -\frac{1}{6}$ and $x = -\frac{3}{4}$ ; required to eliminate $x = -\frac{3}{4}$ for final mark. (ii) Squaring approach: M1 correct method, $24x^2 + 22x + 3 = 0$ (correct 3 term quadratic, any form) A1 Solving M1, Final correct answer A1.]		

**6665 Core C3**  
**Mark Scheme (Post standardisation)**

Question Number	Scheme	Marks
7 (a)	Setting $p = 300$ at $t = 0 \Rightarrow 300 = \frac{2800a}{1 + a}$ $(300 = 2500a); a = 0.12$ (c.s.o) *	M1 dM1A1 (3)
(b)	$1850 = \frac{2800(0.12)e^{0.2t}}{1 + 0.12e^{0.2t}}$ ; $e^{0.2t} = 16.2\dots$ Correctly taking logs to 0.2 $t = \ln k$	M1A1 M1
	$t = 14$ (13.9..)	A1 (4)
(c)	Correct derivation: (Showing division of num. and den. by $e^{0.2t}$ ; using $a$ )	B1 (1)
(d)	Using $t \rightarrow \infty$ , $e^{-0.2t} \rightarrow 0$ , $p \rightarrow \frac{336}{0.12} = 2800$	M1 A1 (2)
		[10]