

GCE

Edexcel GCE

Core Mathematics C3 (6665)




January 2006

advancing learning, changing lives

Mark Scheme (Results)

Edexcel GCE
Core Mathematics C3 (6665)

January 2006
6665 Core Mathematics C3
Mark Scheme


Question Number	Scheme	Marks
1.	<p>(a)</p>  <p>(b)</p>  <p>(c)</p> 	<p style="text-align: center;">Shape unchanged Point</p> <p style="text-align: center;">B1 B1</p> <p style="text-align: center;">(2)</p> <p style="text-align: center;">Shape Point</p> <p style="text-align: center;">B1 B1</p> <p style="text-align: center;">(2)</p> <p style="text-align: center;">Shape (2, 4) (-2, 4)</p> <p style="text-align: center;">B1 B1 B1</p> <p style="text-align: center;">(3) [7]</p>

Question Number	Scheme	Marks
2.	$x^2 - x - 2 = (x - 2)(x + 1)$ <p style="text-align: right;">At any stage</p> $\frac{2x^2 + 3x}{(2x + 3)(x - 2)} = \frac{x(2x + 3)}{(2x + 3)(x - 2)} = \frac{x}{x - 2}$ $\frac{2x^2 + 3x}{(2x + 3)(x - 2)} - \frac{6}{x^2 - x - 2} = \frac{x(x + 1) - 6}{(x - 2)(x + 1)}$ $= \frac{x^2 + x - 6}{(x - 2)(x + 1)}$ $= \frac{(x + 3)(x - 2)}{(x - 2)(x + 1)}$ $= \frac{x + 3}{x + 1}$	<p>B1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>M1 A1</p> <p>A1 (7)</p> <p>[7]</p>
	<p>Alternative method</p> $x^2 - x - 2 = (x - 2)(x + 1)$ <p style="text-align: right;">At any stage</p> <p>(2x + 3) appearing as a factor of the numerator at any stage</p> $\frac{2x^2 + 3x}{(2x + 3)(x - 2)} - \frac{6}{(x - 2)(x + 1)} = \frac{(2x^2 + 3x)(x + 1) - 6(2x + 3)}{(2x + 3)(x - 2)(x + 1)}$ $= \frac{2x^3 + 5x^2 - 9x - 18}{(2x + 3)(x - 2)(x + 1)}$ <p style="text-align: right;">can be implied</p> $= \frac{(x - 2)(2x^2 + 9x + 9)}{(2x + 3)(x - 2)(x + 1)} \text{ or } \frac{(2x + 3)(x^2 + x - 6)}{(2x + 3)(x - 2)(x + 1)} \text{ or } \frac{(x + 3)(2x^2 - x - 6)}{(2x + 3)(x - 2)(x + 1)}$ <p style="text-align: center;">Any one linear factor × quadratic</p> $= \frac{(2x + 3)(x - 2)(x + 3)}{(2x + 3)(x - 2)(x + 1)}$ <p style="text-align: right;">Complete factors</p> $= \frac{x + 3}{x + 1}$	<p>B1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>A1 (7)</p>

Question Number	Scheme	Marks
3.	$\frac{dy}{dx} = \frac{1}{x}$ <p style="text-align: right;">accept $\frac{3}{3x}$</p> $\text{At } x = 3, \frac{dy}{dx} = \frac{1}{3} \Rightarrow m' = -3$ <p style="text-align: right;">Use of $mm' = -1$</p> $y - \ln 1 = -3(x - 3)$ $y = -3x + 9$ <p style="text-align: right;">Accept $y = 9 - 3x$</p> <p>$\frac{dy}{dx} = \frac{1}{3x}$ leading to $y = -9x + 27$ is a maximum of M1 A0 M1 M1 A0 = 3/5</p>	M1 A1 M1 M1 A1 (5) [5]
4.	<p>(a) (i) $\frac{d}{dx}(e^{3x+2}) = 3e^{3x+2}$ (or $3e^2e^{3x}$)</p> <p style="text-align: right;">At any stage</p> <p>$\frac{dy}{dx} = 3x^2 e^{3x+2} + 2xe^{3x+2}$</p> <p style="text-align: right;">Or equivalent</p> <p>(ii) $\frac{d}{dx}(\cos(2x^3)) = -6x^2 \sin(2x^3)$</p> <p style="text-align: right;">At any stage</p> $\frac{dy}{dx} = \frac{-18x^3 \sin(2x^3) - 3\cos(2x^3)}{9x^2}$ <p style="text-align: right;">M1 A1 (4)</p> <p>Alternatively using the product rule for second M1 A1</p> $y = (3x)^{-1} \cos(2x^3)$ $\frac{dy}{dx} = -3(3x)^{-2} \cos(2x^3) - 6x^2 (3x)^{-1} \sin(2x^3)$ <p style="text-align: right;">Accept equivalent unsimplified forms</p> <p>(b) $1 = 8 \cos(2y + 6) \frac{dy}{dx}$ or $\frac{dx}{dy} = 8 \cos(2y + 6)$</p> <p style="text-align: right;">M1</p> $\frac{dy}{dx} = \frac{1}{8 \cos(2y + 6)}$ <p style="text-align: right;">M1 A1</p> $\frac{dy}{dx} = \frac{1}{8 \cos\left(\arcsin\left(\frac{x}{4}\right)\right)} \left(= (\pm) \frac{1}{2\sqrt{(16-x^2)}} \right)$ <p style="text-align: right;">M1 A1 (5)</p> <p style="text-align: right;">[13]</p>	B1 M1 A1+A1 (4) M1 A1 M1 A1 (4) M1 M1 A1 M1 A1 (5) [13]

Question Number	Scheme	Marks
5.	<p>(a) $2x^2 - 1 - \frac{4}{x} = 0$ Dividing equation by x M1 $x^2 = \frac{1}{2} + \frac{4}{2x}$ Obtaining $x^2 = \dots$ M1 $x = \sqrt{\left(\frac{2}{x} + \frac{1}{2}\right)}$ * cso A1 (3)</p> <p>(b) $x_1 = 1.41, x_2 = 1.39, x_3 = 1.39$ B1, B1, B1 If answers given to more than 2 dp, penalise first time then accept awrt above. (3)</p> <p>(c) Choosing (1.3915, 1.3925) or a tighter interval M1 $f(1.3915) \approx -3 \times 10^{-3}, f(1.3925) \approx 7 \times 10^{-3}$ Both, awrt A1 Change of sign (and continuity) $\Rightarrow \alpha \in (1.3915, 1.3925)$ $\Rightarrow \alpha = 1.392$ to 3 decimal places * cso A1 (3)</p>	<p>(3)</p> <p>(3)</p> <p>(3)</p> <p>[9]</p>
6.	<p>(a) $R \cos \alpha = 12, R \sin \alpha = 4$ $R = \sqrt{(12^2 + 4^2)} = \sqrt{160}$ Accept if just written down, awrt 12.6 M1 A1 $\tan \alpha = \frac{4}{12}, \Rightarrow \alpha \approx 18.43^\circ$ awrt 18.4° M1, A1(4)</p> <p>(b) $\cos(x + \text{their } \alpha) = \frac{7}{\text{their } R}$ (≈ 0.5534) M1 $x + \text{their } \alpha = 56.4^\circ$ awrt 56° A1 $= \dots, 303.6^\circ$ $360^\circ - \text{their principal value}$ M1 $x = 38.0^\circ, 285.2^\circ$ Ignore solutions out of range A1, A1 (5) If answers given to more than 1 dp, penalise first time then accept awrt above.</p> <p>(c)(i) minimum value is $-\sqrt{160}$ ft their R B1ft</p> <p>(ii) $\cos(x + \text{their } \alpha) = -1$ M1 $x \approx 161.57^\circ$ cao A1 (3)</p>	<p>(4)</p> <p>(5)</p> <p>(3)</p> <p>[12]</p>

Question Number	Scheme	Marks
7.	(a) (i) Use of $\cos 2x = \cos^2 x - \sin^2 x$ in an attempt to prove the identity. $\frac{\cos 2x}{\cos x + \sin x} = \frac{\cos^2 x - \sin^2 x}{\cos x + \sin x} = \frac{(\cos x - \sin x)(\cos x + \sin x)}{\cos x + \sin x} = \cos x - \sin x \quad *$	M1 A1 (2)
	(ii) Use of $\cos 2x = 2\cos^2 x - 1$ in an attempt to prove the identity. Use of $\sin 2x = 2\sin x \cos x$ in an attempt to prove the identity. $\frac{1}{2}(\cos 2x - \sin 2x) = \frac{1}{2}(2\cos^2 x - 1 - 2\sin x \cos x) = \cos^2 x - \cos x \sin x - \frac{1}{2} \quad *$	M1 M1 A1 (3)
	(b) $\cos \theta (\cos \theta - \sin \theta) = \frac{1}{2}$ $\cos^2 \theta - \cos \theta \sin \theta - \frac{1}{2} = 0$	Using (a)(i) M1
	$\frac{1}{2}(\cos 2\theta - \sin 2\theta) = 0$ $\cos 2\theta = \sin 2\theta \quad *$	Using (a)(ii) M1 A1 (3)
	(c) $\tan 2\theta = 1$ $2\theta = \frac{\pi}{4}, \left(\frac{5\pi}{4}, \frac{9\pi}{4}, \frac{13\pi}{4} \right)$	M1 any one correct value of 2θ A1
	$\theta = \frac{\pi}{8}, \frac{5\pi}{8}, \frac{9\pi}{8}, \frac{13\pi}{8}$	Obtaining at least 2 solutions in range M1
	$\text{The 4 correct solutions}$	A1 (4)
	If decimals (0.393, 1.963, 3.534, 5.105) or degrees (22.5°, 112.5°, 202.5°, 292.5°) are given, but all 4 solutions are found, penalise one A mark only. Ignore solutions out of range.	[12]

Question Number	Scheme	Marks
8.	<p>(a)</p> $gf(x) = e^{2(2x+\ln 2)}$ $= e^{4x} e^{2\ln 2}$ $= e^{4x} e^{\ln 4}$ $= 4e^{4x}$ <p>(Hence $gf : x \mapsto 4e^{4x}, x \in \mathbb{R}$)</p>	<p>M1 M1 M1 A1 (4)</p> <p>Give mark at this point, cso</p>
	<p>(b)</p>  <p>Shape and point</p>	<p>B1 (1)</p>
	<p>(c) Range is \mathbb{R}_+</p>	<p>Accept $gf(x) > 0, y > 0$</p> <p>B1 (1)</p>
	<p>(d)</p> $\frac{d}{dx}[gf(x)] = 16e^{4x}$ $e^{4x} = \frac{3}{16}$ $4x = \ln \frac{3}{16}$ $x \approx -0.418$	<p>M1 A1 M1 A1 (4) [10]</p>