

Question Number	Scheme	Marks												
1.	<p>Differentiates</p> <p style="text-align: center;">$6x + 8y \frac{dy}{dx} - 2,$</p> <p>to obtain : $\dots + (6x \frac{dy}{dx} + 6y) = 0$</p> <p style="text-align: center;">$\left[\frac{dy}{dx} = \frac{2 - 6x - 6y}{6x + 8y} \right]$</p> <p>Substitutes $x = 1, y = -2$ into expression involving $\frac{dy}{dx}$, to give $\frac{dy}{dx} = -\frac{8}{10}$</p> <p>Uses line equation with numerical ‘gradient’ $y - (-2) = (\text{their gradient})(x - 1)$ or finds c and uses $y = (\text{their gradient})x + "c"$</p> <p>To give $5y + 4x + 6 = 0$ (or equivalent = 0)</p>	M1 A1, +(B1)												
2. (a)	<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="padding: 5px; text-align: center;">x</td> <td style="padding: 5px; text-align: center;">0</td> <td style="padding: 5px; text-align: center;">$\frac{\pi}{16}$</td> <td style="padding: 5px; text-align: center;">$\frac{\pi}{8}$</td> <td style="padding: 5px; text-align: center;">$\frac{3\pi}{16}$</td> <td style="padding: 5px; text-align: center;">$\frac{\pi}{4}$</td> </tr> <tr> <td style="padding: 5px; text-align: center;">y</td> <td style="padding: 5px; text-align: center;">1</td> <td style="padding: 5px; text-align: center;">1.01959</td> <td style="padding: 5px; text-align: center;">1.08239</td> <td style="padding: 5px; text-align: center;">1.20269</td> <td style="padding: 5px; text-align: center;">1.41421</td> </tr> </table> <p>M1 for one correct, A1 for all correct</p>	x	0	$\frac{\pi}{16}$	$\frac{\pi}{8}$	$\frac{3\pi}{16}$	$\frac{\pi}{4}$	y	1	1.01959	1.08239	1.20269	1.41421	M1, A1 (2)
x	0	$\frac{\pi}{16}$	$\frac{\pi}{8}$	$\frac{3\pi}{16}$	$\frac{\pi}{4}$									
y	1	1.01959	1.08239	1.20269	1.41421									
2. (b)	<p>Integral = $\frac{1}{2} \times \frac{\pi}{16} \times \{1 + 1.4142 + 2(1.01959 + \dots + 1.20269)\}$</p> $\left(= \frac{\pi}{32} \times 9.02355 \right) = 0.8859$	M1 A1/ A1 cao (3)												
2. (c)	<p>Percentage error = $\frac{\text{approx} - 0.88137}{0.88137} \times 100 = 0.51\%$ (allow 0.5% to 0.54% for A1)</p> <p>M1 gained for $(\pm) \frac{\text{approx} - \ln(1 + \sqrt{2})}{\ln(1 + \sqrt{2})}$</p>	M1 A1 (2) (7)												

Question Number	Scheme	Marks
3.	Uses substitution to obtain $x = f(u) \left[\frac{u^2 + 1}{2} \right]$, M1 and to obtain $u \frac{du}{dx} = \text{const. or equiv.}$ M1 Reaches $\int \frac{3(u^2 + 1)}{2u} u du$ or equivalent A1 Simplifies integrand to $\int \left(3u^2 + \frac{3}{2} \right) du$ or equiv. M1 Integrates to $\frac{1}{2}u^3 + \frac{3}{2}u$ M1 A1√ A1√ dependent on all previous Ms Uses new limits 3 and 1 substituting and subtracting (or returning to function of x with old limits) M1 To give 16 cso A1	[8]

“By Parts”

Attempt at “right direction” by parts

$$[\ 3x \left(2x - 1 \right)^{\frac{1}{2}}] - \left\{ \int 3 \left(2x - 1 \right)^{\frac{1}{2}} dx \right\} \quad \text{M1}\{\text{M1A1}\}$$

$$\dots \dots \dots - (2x - 1)^{\frac{3}{2}} \quad \text{M1A1√}$$

Uses limits 5 and 1 correctly; [42 – 26] 16

M1A1

4.	<p>Attempts $V = \pi \int x^2 e^{2x} dx$</p> $= \pi \left[\frac{x^2 e^{2x}}{2} - \int x e^{2x} dx \right] \quad (\text{M1 needs parts in the correct direction})$ $= \pi \left[\frac{x^2 e^{2x}}{2} - \left(\frac{x e^{2x}}{2} - \int \frac{e^{2x}}{2} dx \right) \right] \quad (\text{M1 needs second application of parts})$ <p>M1A1\checkmark refers to candidates $\int x e^{2x} dx$, but dependent on prev. M1</p> $= \pi \left[\frac{x^2 e^{2x}}{2} - \left(\frac{x e^{2x}}{2} - \frac{e^{2x}}{4} \right) \right]$ <p>Substitutes limits 3 and 1 and subtracts to give... [dep. on second and third Ms]</p> $= \pi \left[\frac{13}{4} e^6 - \frac{1}{4} e^2 \right] \text{ or any correct exact equivalent.}$ <p>[Omission of } \pi \text{ loses first and last marks only]</p>	M1 M1 A1 M1 A1 \checkmark A1 cao dM1 A1 [8]
----	--	--

Question Number	Scheme	Marks
5. (a)	<p>Considers $3x^2 + 16 = A(2+x)^2 + B(1-3x)(2+x) + C(1-3x)$ and substitutes $x = -2$, or $x = 1/3$, or compares coefficients and solves simultaneous equations To obtain $A = 3$, and $C = 4$ Compares coefficients or uses simultaneous equation to show $B = 0$.</p>	M1 A1, A1 B1 (4)
(b)	<p>Writes $3(1-3x)^{-1} + 4(2+x)^{-2}$ $= 3(1+3x, +9x^2 + 27x^3 + \dots) +$ $\frac{4}{4}(1 + \frac{(-2)}{1}(\frac{x}{2}) + \frac{(-2)(-3)}{1.2}(\frac{x}{2})^2 + \frac{(-2)(-3)(-4)}{1.2.3}(\frac{x}{2})^3 + \dots)$ $= 4 + 8x, + 27\frac{3}{4}x^2 + 80\frac{1}{2}x^3 + \dots$ Or uses $(3x^2 + 16)(1-3x)^{-1}(2+x)^{-2}$ $(3x^2 + 16)(1+3x, +9x^2 + 27x^3 +) \times$ $\frac{1}{4}(1 + \frac{(-2)}{1}(\frac{x}{2}) + \frac{(-2)(-3)}{1.2}(\frac{x}{2})^2 + \frac{(-2)(-3)(-4)}{1.2.3}(\frac{x}{2})^3)$ $= 4 + 8x, + 27\frac{3}{4}x^2 + 80\frac{1}{2}x^3 + \dots$</p>	M1 (M1, A1) (M1 A1) A1, A1 (7) M1 (M1A1)× (M1A1) A1, A1 (7) [11]

	$\lambda = -4 \rightarrow a = 18, \quad \mu = 1 \rightarrow b = 9$	M1 A1, A1 (3)
6. (a)		
(b)	$\begin{pmatrix} 8+\lambda \\ 12+\lambda \\ 14-\lambda \end{pmatrix} \bullet \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix} = 0$	M1
	$\therefore 8 + \lambda + 12 + \lambda - 14 + \lambda = 0$	A1
	Solves to obtain λ ($\lambda = -2$)	dM1
	Then substitutes value for λ to give P at the point (6, 10, 16) (any form)	M1, A1 (5)
(c)	$OP = \sqrt{36+100+256}$ $(= \sqrt{392}) = 14\sqrt{2}$	M1 A1 cao (2) [10]
7. (a)	$\frac{dV}{dr} = 4\pi r^2$	B1 (1)
(b)	Uses $\frac{dr}{dt} = \frac{dV}{dt} \cdot \frac{dt}{dV}$ in any form, $= \frac{1000}{4\pi r^2 (2t+1)^2}$	M1, A1 (2)
(c)	$V = \int 1000(2t+1)^{-2} dt$ and integrate to $p(2t+1)^{-1}, = -500(2t+1)^{-1}(+c)$	M1, A1
	Using $V=0$ when $t=0$ to find c , ($c = 500$, or equivalent)	M1
	$\therefore V = 500(1 - \frac{1}{2t+1})$ (any form)	A1 (4)
(d)	(i) Substitute $t = 5$ to give V , then use $r = \sqrt[3]{\frac{3V}{4\pi}}$ to give $r, = 4.77$	M1, M1, A1 (3)
	(ii) Substitutes $t = 5$ and $r = \text{'their value'}$ into 'their' part (b)	M1
	$\frac{dr}{dt} = 0.0289 \quad (\approx 2.90 \times 10^{-2}) \text{ (cm/s)} * \text{ AG}$	A1 (2) [12]

8. (a) Solves $y = 0 \Rightarrow \cos t = \frac{1}{2}$ to obtain $t = \frac{\pi}{3}$ or $\frac{5\pi}{3}$ (need both for A1)

Or substitutes **both** values of t and shows that $y = 0$

$$(b) \quad \frac{dx}{dt} = 1 - 2 \cos t$$

$$\text{Area} = \int y dx = \int_{\pi/3}^{5\pi/3} (1 - 2 \cos t)(1 - 2 \cos t) dt = \int_{\pi/3}^{5\pi/3} (1 - 2 \cos t)^2 dt * \text{AG}$$

$$(c) \quad \text{Area} = \int 1 - 4 \cos t + 4 \cos^2 t dt \quad 3 \text{ terms}$$

$$= \int 1 - 4 \cos t + 2(\cos 2t + 1) dt \quad (\text{use of correct double angle formula})$$

$$= \int 3 - 4 \cos t + 2 \cos 2t dt$$

$$= [3t - 4 \sin t + \sin 2t]$$

Substitutes the two correct limits $t = \frac{5\pi}{3}$ and $\frac{\pi}{3}$ and subtracts.

$$= 4\pi + 3\sqrt{3}$$