


Question Number	Scheme	Marks
1.	$\cos \frac{\pi}{2} + i \sin \frac{\pi}{2}$ $\cos \frac{\pi}{10} + i \sin \frac{\pi}{10}$ $\cos\left(\frac{(4k+1)\pi}{10}\right) + i \sin\left(\frac{(4k+1)\pi}{10}\right), k = 2, 3, 4 (\text{or equiv.})$ $[\cos\left(\frac{9\pi}{10}\right) + i \sin\left(\frac{9\pi}{10}\right), \cos\left(\frac{13\pi}{10}\right) + i \sin\left(\frac{13\pi}{10}\right), \cos\left(\frac{17\pi}{10}\right) + i \sin\left(\frac{17\pi}{10}\right)]$ [Degrees : 18, 90, 162, 234, 306]	B1 B1 M1A2,1,0 (5)
		Total 5 marks
2.	$\left(\frac{dy}{dx}\right)_0 \approx \frac{y_1 - y_{-1}}{2h} \Rightarrow 2 \approx \frac{y_1 - y_{-1}}{0.2} \Rightarrow y_1 - y_{-1} \approx 0.4$ $\left(\frac{d^2y}{dx^2}\right)_0 \approx \frac{y_1 - 2y_0 + y_{-1}}{h^2} \Rightarrow 8 \approx \frac{y_1 - 2y_0 + y_{-1}}{0.01}$ [For M1, an attempt at evaluating $\left(\frac{d^2y}{dx^2}\right)_0$ is required.] $\Rightarrow y_1 + y_{-1} \approx 2.08$ Subtracting to give $y_{-1} \approx 0.84$	M1A1 M1 A1 M1A1 (6)
		Total 6 marks
3.	(a) Complete method for finding image: e.g. $\begin{pmatrix} -4 & 2 \\ 2 & -1 \end{pmatrix} \begin{pmatrix} x \\ 2x+1 \end{pmatrix} = \begin{pmatrix} 2 \\ -1 \end{pmatrix}$ The image is the point $(2, -1)$ (b) $\begin{vmatrix} k-\lambda & 2 \\ 2 & -1-\lambda \end{vmatrix} = 0$ Characteristic equation: $\lambda^2 - \lambda - 6 = 0$ Solving: $(\lambda - 3)(\lambda + 2) = 0 \Rightarrow \lambda = \dots$ $\lambda = -2, \lambda = 3$ (both) (c) Method for finding an eigenvector $\begin{pmatrix} 2 & 2 \\ 2 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = 3 \begin{pmatrix} x \\ y \end{pmatrix} \Rightarrow \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 2 \\ 1 \end{pmatrix}$ and	M1 A1 (2) M1 A1 M1 A1 (4) M1

	$\begin{pmatrix} 2 & 2 \\ 2 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = -2 \begin{pmatrix} x \\ y \end{pmatrix} \Rightarrow \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1 \\ -2 \end{pmatrix}$ <p>Equations are: $y = \frac{1}{2}x$ and $y = -2x$.</p> <p>Alt: $\begin{pmatrix} 2 & 2 \\ 2 & -1 \end{pmatrix} \begin{pmatrix} x \\ mx \end{pmatrix} = \begin{pmatrix} y \\ my \end{pmatrix} \Rightarrow 2m^2 + 3m - 2 = 0$ M1A1 $[m = \frac{1}{2}, -2]$ Correct equations A1</p>	A1 \checkmark A1 (3)
4.	$\mathbf{A} = \begin{pmatrix} k & 1 & -2 \\ 0 & -1 & k \\ 9 & 1 & 0 \end{pmatrix}$ <p>(a) Det. $\mathbf{A} = -k^2 + 9k - 18$</p> <p>Setting to zero and solving for k $[(k-6)(k-3) = 0]$ $\Rightarrow k = 3, k = 6$</p> <p>(b) Cofactors $\begin{pmatrix} -k & 9k & 9 \\ -2 & 18 & 9-k \\ k-2 & -k^2 & -k \end{pmatrix}$ [B1 for each row (or column)]</p> $\mathbf{A}^{-1} = \frac{1}{\det(\mathbf{A})} \begin{pmatrix} -k & -2 & k-2 \\ 9k & 18 & -k^2 \\ 9 & 9-k & -k \end{pmatrix}$ <p>[A1 f.t. is on determinant or cofactors]</p>	M1A1 M1 A1 (4) B3 M1A1 \checkmark (5) Total 9 marks

5.	<p>(i) When $n = 1$, LHS = $1(2)^1 = 2$; RHS = $2\{1 + 0\} = 2 \Rightarrow$ true for $n = 1$</p> <p>Suppose true for $n = k$, then</p> $\begin{aligned}\sum_{r=1}^{k+1} r 2^r &= 2\{1 + (k-1)2^k + (k+1)2^{k+1}\} \\&= 2 + k 2^{k+1} + k 2^{k+1} \\&= 2(1 + k 2^{k+1}) \\&= 2[1 + \{(k+1)-1\}2^{k+1}]\end{aligned}$ <p>So, if true for $n = k$ then true for $n = k+1$, but true for $n = 1$, \therefore true, by induction, for all values of $n \in \mathbb{Z}^+$.</p> <p>(ii) Showing true for $n = 1$,</p> $\begin{aligned}\frac{dy}{dx} &= \frac{3}{2+3x} \\ \frac{d^{n+1}y}{dx^{n+1}} &= (-1)^{n+1} \frac{3^n(n-1)!(-3n)}{(2+3x)^{n+1}} \\&= (-1)^{n+2} \frac{3^{n+1}(n)!}{(2+3x)^{n+1}}\end{aligned}$ <p>Conclusion</p>	<p>B1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1(cso)</p> <p>B1</p> <p>M1A1</p> <p>A1</p> <p>A1(cso)</p> <p>Total 10 marks</p>
6.	<p>(a) Correct method for producing 2nd order differential equation</p> <p>e.g. $\frac{d}{dx} \left\{ (1+2x) \frac{dy}{dx} \right\} = \frac{d}{dx} \left\{ x + 4y^2 \right\}$ attempted</p> $(1+2x) \frac{d^2y}{dx^2} + 2 \frac{dy}{dx} = 1 + 8y \frac{dy}{dx}$ seen + conclusion AG <p>(b) Differentiating again w.r.t. x:</p> $(1+2x) \frac{d^3y}{dx^3} + 2 \frac{d^2y}{dx^2} = 8y \frac{d^2y}{dx^2} + 8 \left(\frac{dy}{dx} \right)^2 - 2 \frac{d^2y}{dx^2}$ or equiv. <p>[e.g. $(1+2x) \frac{d^3y}{dx^3} = 8 \left(\frac{dy}{dx} \right)^2 + 4(2y-1) \frac{d^2y}{dx^2}$</p> <p>(c) $\frac{dy}{dx}$ (at $x = 0$) = 1</p>	<p>M1</p> <p>A1*</p> <p>(2)</p> <p>M1A2,1,0</p> <p>(3)</p>

	<p>Finding $\frac{d^2y}{dx^2}$ (at $x = 0$) ($= 3$)</p> <p>Finding $\frac{d^3y}{dx^3}$, at $x = 0$; $= 8$ [A1 f.t. is on part (c) values only]</p> $y = \frac{1}{2} + x + \frac{3}{2}x^2 + \frac{4}{3}x^3 + \dots$ <p>[Alternative (c): Polynomial for y: $y = \frac{1}{2} + ax + bx^2 + cx^3 + \dots$ In given d.e.: $(1 + 2x)(a + 2bx + 3cx^2 + \dots) \equiv x + 4(\frac{1}{2} + ax + bx^2 + cx^3 + \dots)^2$ M1A1 $a = 1$ B1, Complete method for other coefficients M1, answer A1]</p>	B1 M1 M1A1✓ M1A1 M1A1 Total 11 marks (6)
7.	<p>(a) $R\vec{Q} = \begin{pmatrix} 1 \\ -1 \\ -1-c \end{pmatrix}$, $R\vec{P} = \begin{pmatrix} -4 \\ 3 \\ -2-c \end{pmatrix}$ (both)</p> $R\vec{P} \times R\vec{Q} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -4 & 3 & -2-c \\ 1 & -1 & -1-c \end{vmatrix}$ $= (-5 - 4c)\mathbf{i} - (6 + 5c)\mathbf{j} + \mathbf{k}$	B1
	(b) $c = -2$ $d = -6 - 5c = 4$ AG	M1A1✓ (3)
	(c) $\mathbf{r} \cdot \begin{pmatrix} 3 \\ 4 \\ 1 \end{pmatrix} = p$ Substituting point in plane to give p , $\mathbf{r} \cdot \begin{pmatrix} 3 \\ 4 \\ 1 \end{pmatrix} = 7$.	A1✓ A1*(cso) (2)
	(d) Equation of normal to plane through S : $\mathbf{r} = \begin{pmatrix} 1 \\ 5 \\ 10 \end{pmatrix} + t \begin{pmatrix} 3 \\ 4 \\ 1 \end{pmatrix}$ Meets plane where $\begin{pmatrix} 1+3t \\ 5+4t \\ 10+t \end{pmatrix} \cdot \begin{pmatrix} 3 \\ 4 \\ 1 \end{pmatrix} = 7 \Rightarrow t = -1$ S' has position vector $\begin{pmatrix} 1 \\ 5 \\ 10 \end{pmatrix} + 2t \begin{pmatrix} 3 \\ 4 \\ 1 \end{pmatrix} = \begin{pmatrix} -5 \\ -3 \\ 8 \end{pmatrix}$	M1A1 B1 M1A1✓ (3)

	(a) Relating lines and angle (generous) [angle between $\pm 2i$ to P and ± 2 to P]	M1A1 Total 13 marks
8.	<p>Angle between correct lines is $\frac{\pi}{2}$ Circle Selecting correct ("top half") semi-circle .</p> <p>[If algebraic approach: Method for finding Cartesian equation M1 Correct equation, any form, $\Rightarrow x(x + 2) + y(y - 2) = 0$ A1</p> <p>Sketch: showing circle M1 Correct circle { centre $(-1, 1)$ }, choosing only "top half" A1]</p> <p>(b) $z + 1 - i$ is radius; $= \sqrt{2}$ M1 A1</p> <p>(c) $z = \frac{2(1+i) - 2\omega}{\omega} \quad \left(= \frac{2(1+i)}{\omega} - 2 \right)$ M1A1 (4) $\frac{z - 2i}{z + 2} = \frac{2(1+i) - 2(1+i)\omega}{2(1+i)} \quad (= 1 - \omega)$ M1 (2)</p> <p>$\text{Arg}(1 - \omega) = \frac{\pi}{2}$ is line segment, passing through $(1, 0)$ M1A1</p>  <p>u A1,A1</p> <p>A1 (6)</p> <p>Alt C: $u + iv = \frac{2 + 2i}{(x + 2) + iy} = \frac{(2x + 2y + 4) + i(x + 2 - y)}{(x + 2)^2 + y^2}$ M1 $x = -1 + \sqrt{2} \cos \theta, y = 1 + \sqrt{2} \sin \theta$ M1 $\Rightarrow w = \frac{(2\sqrt{2} \cos \theta + 2\sqrt{2} \sin \theta + 4) + i \dots}{(2\sqrt{2} \cos \theta + 2\sqrt{2} \sin \theta + 4)} \quad \{= 1 + i f(\theta)\}$ A1, \Rightarrow part of line $u = 1$, show lower "half" of line A1,A1</p>	