


January 2007  
6674 Further Pure Mathematics FP1  
Mark Scheme

Question Number	Scheme	Marks
1.	<p>(a) Method for finding <math>z</math> : <math>z = \frac{-2 \pm \sqrt{4 - 68}}{2}</math> , <math>= \frac{-2 \pm \sqrt{64} i}{2}</math></p> <p>[Completing the square: <math>(z + 1)^2 + 16 = 0</math> , <math>z = -1 \pm \sqrt{16} i</math> M1,A1</p> <p style="text-align: center;"><math>z = -1 \pm 4i</math>      (<math>a = -1</math>, <math>b = \pm 4</math>)</p> <p>(b)</p> 	<p>M1, A1</p> <p>A1 (3)</p> <p>B1 ✓ (1)</p> <p>[4]</p>
	<p>Notes</p> <p>(a) First A1 is unsimplified but requires <math>i</math>  <math>-1 \pm 8i</math> only scores M1 unless intermediate step seen when M1A1 possible                      Correct answer with no working is full marks</p> <p>SC: If M0 awarded, <math>k \pm 4i</math>, <math>k + 4i</math>, <math>k - 4i</math> scores B1 (Epen M0A0A1)</p> <p>Use of <math>z = a + ib</math></p> <p>(i) <math>z^2 - 2az + a^2 + b^2 = z^2 + 2z + 17 = 0</math> and compare coefficients M1</p> <p style="text-align: center;"><math>a^2 + b^2 = 17</math> and <math>a = -1</math>;      <math>z = -1 \pm 4i</math> A1, A1</p> <p>(ii) <math>(a + ib)^2 + 2(a + ib) + 17 = 0</math> and compare coefficients M1</p> <p style="text-align: center;"><math>2b(a + 1) = 0</math> and <math>a^2 - b^2 + 2a = -17</math>, <math>a = -1</math> and <math>b = \pm 4</math> A1, A1</p> <p>(b) Must be a conjugate pair.</p> <p>Allow: Coords marked at points or “correct” numbers on axes.(allow “graduations”)</p> <p style="text-align: center;">(Ignore any lines drawn)</p>	

<p>2.</p>	<p>Attempt to arrange in correct form <math>\frac{dy}{dx} + \frac{2}{x}y = \frac{\cos x}{x}</math></p> <p>Integrating Factor: <math>= e^{\int \frac{2}{x} dx}</math>, <math>[ (= e^{2 \ln x} = e^{\ln x^2}) = x^2</math></p> <p><math>[ x^2 \frac{dy}{dx} + 2xy = x \cos x</math> implies M1M1A1]</p> <p><math>\therefore x^2 y = \int x^2 \cdot \frac{\cos x}{x} dx</math> or equiv.</p> <p>[ I.F. <math>y = \int I.F. (candidate's RHS) dx</math> ]</p> <p>By Parts: <math>(x^2 y) = x \sin x - \int \sin x dx</math></p> <p>i.e. <math>(x^2 y) = x \sin x, + \cos x (+ c)</math></p> $y = \frac{\sin x}{x} + \frac{\cos x}{x^2} + \frac{c}{x^2}$	<p>M1</p> <p>M1,A1</p> <p>M1√</p> <p>M1</p> <p>A1, A1cao</p> <p>A1√</p> <p>[8]</p>
	<p>Notes:</p> <p>First M: At least two terms divided by <math>x</math>.</p> <p>“By parts” M: Must be complete method, e.g <math>\int x^2 \cos x dx</math> requires <b>two</b> applications</p> <p>Because of functions involved, <b>be generous with sign</b>, but</p> <p><math>x \sin x \pm \int \cos x dx</math> is M0</p> <p>(S.C. “Loop” integral like <math>\int e^x \cos x dx</math>, allow M1 if two applications of “by parts”, despite incomplete method)</p> <p>Final A f.t. for dividing all terms by candidates I.F., providing “c” used.</p>	

<p>3.</p>	<p>(a) <math>\frac{z_2}{z_1} = \frac{1 + pi}{5 + 3i} \cdot \frac{(5 - 3i)}{(5 - 3i)}</math></p> <p><math>= \frac{5 + 5pi - 3i + 3p}{(34)}</math> [Multiply out and attempt use of <math>i^2 = -1</math>]</p> <p><math>= \frac{5 + 3p}{34} + \frac{5p - 3}{34}i</math> or <math>\frac{5 + 3p}{34} - \frac{3 - 3p}{34}i</math></p> <p>(b) For <math>\frac{z_2}{z_1} = c + id</math> using <math>\frac{d}{c} = \tan \frac{\pi}{4}</math>:</p> <p>[ <math>5p - 3 = 5 + 3p</math> ] <math>\Rightarrow p = 4</math></p>	<p>M1</p> <p>M1</p> <p>A1 (3)</p> <p>M1</p> <p>A1 (2)</p> <p>[5]</p>
	<p>Notes:</p> <p>In (a) if <math>\frac{z_1}{z_2}</math> used treat as MR. Can score (a)M1M1A0 (b)M1A0</p> <p><math>\left[ (a) \frac{5+3p}{1+p^2} + \frac{3-5p}{1+p^2}i \quad (b) -\frac{1}{4} \right]</math></p> <p>Allow A1 if answer “all over” 34, real and imag. collected up)</p> <p><math>1 + pi = (a + ib)(5 + 3i)</math>: M1 compare real and imag. is first M mark</p> <p>If denominator in (a) incorrect, both marks in (b) still available</p> <p>In (b), if use <math>\arg z_2 - \arg z_1 = \frac{\pi}{4}</math>:</p> <p>M1 for <math>\arctan p - \arctan \frac{3}{5} = \frac{\pi}{4}</math> [ <math>\arctan p = \frac{\pi}{4} + 0.5404\dots = 1.3258</math> ]</p> <p>Allow A1 for <math>p = 4</math> without further work or for that shown in brackets, i.e. assume values retained on calculator (no penalty because it looks as though not exact)</p>	

<p>4.</p>	<p>Working from RHS:</p> <p>(a) Combining <math>\frac{1}{r} - \frac{1}{r+1}</math> <math>\left[ \frac{1}{r(r+1)} \right]</math></p> <p>Forming single fraction : <math>\frac{r(r-1)(r+1) + (r+1) - r}{r(r+1)}</math></p> $= \frac{r(r^2 - 1) + 1}{r(r+1)} = \frac{r^3 - r + 1}{r(r+1)} \quad \text{AG}$ <p>Note: For A1, must be intermediate step, as shown</p> <p>Working from LHS:</p> <p>(a) <math>\frac{r(r^2 - 1) + 1}{r(r+1)} = \frac{r(r+1)(r-1) + 1}{r(r+1)} = r - 1 + \frac{1}{r(r+1)}</math></p> <p>Splitting <math>\frac{1}{r(r+1)}</math> into partial fractions</p> <p>Showing <math>= \frac{r(r^2 - 1) + 1}{r(r+1)} = r - 1 + \frac{1}{r} - \frac{1}{r+1}</math> no incorrect working seen</p>	<p>M1</p> <p>M1</p> <p>A1cso (3)</p> <p>M1</p> <p>M1</p> <p>A1</p>
	<p>Notes:</p> <p>In first method, second M needs all necessary terms, allowing for sign errors</p> <p>In second method first M is for division:</p> <p>Second method mark is for method shown (allow “cover up” rule stated)</p> <p>If long division, allow reasonable attempt which has remainder constant or linear function of r.</p> <p>Setting <math>\frac{r(r^2 - 1) + 1}{r(r+1)} = \frac{A}{r} + \frac{B}{r+1}</math> is M0</p> <p>If 3 or 4 constants used in a correct initial statement,</p> <p>M1 for finding 2 constants; M1 for complete method to find remaining constant(s)</p>	

	<p>(b) <math display="block">\sum_1^n r - \sum_1^n 1 + \sum_1^n \left( \frac{1}{r} - \frac{1}{r+1} \right)</math></p> <p><math display="block">= \frac{n(n+1)}{2}, (-) n, + \dots +</math></p> <p><math display="block">\left[ \left( 1 - \frac{1}{2} \right) + \left( \frac{1}{2} - \frac{1}{3} \right) + \dots + \left( \frac{1}{n} - \frac{1}{n+1} \right) \right] =</math></p> <p>Simplification of method of differences: <math>1 - \frac{1}{n+1}</math></p> <p><math display="block">\left\{ = \frac{n(n-1)}{2} + \left[ 1 - \frac{1}{(n+1)} \right] \right\}</math></p> <p>Attempt single fraction: <math>= \frac{n(n+1)(n-1) + 2n}{2(n+1)}</math> (dep. prev. M1)</p> <p><math display="block">= \frac{n(n^2+1)}{2(n+1)} \quad \text{or} \quad \frac{n^3+n}{2(n+1)}</math></p> <p><i>Alternative:</i> Using Difference method on whole expression:</p> <p><math display="block">\left[ 0 + 1 - \frac{1}{2} \right] + \left[ 1 + \frac{1}{2} - \frac{1}{3} \right] + \left[ 2 + \frac{1}{3} - \frac{1}{4} \right] \dots \left[ n-1 + \frac{1}{n} - \frac{1}{n+1} \right]</math></p> <p><math display="block">= (1+2+3 \dots + n-1), + \left[ \left( 1 - \frac{1}{n+1} \right) \right] \text{ any form}</math></p> <p><math display="block">= \frac{n(n-1)}{2}, \quad \left\{ + \frac{n}{n+1} \right\}</math></p> <p><math display="block">= \frac{n(n+1)(n-1) + 2n}{2(n+1)} \quad \text{[Attempt single fraction]}</math></p> <p><math display="block">= \frac{n(n^2+1)}{2(n+1)} \quad \text{or} \quad \frac{n^3+n}{2(n+1)}</math></p> <p>Notes:</p> <p>First M mark is for use of method of differences and attempt at some simplification</p> <p>First A mark is for simplified result of this method (no more than 2 terms)</p> <p>Second M mark for attempt at forming single fraction, dependent on first M mark</p> <p>In alternative first B1 need not be added but need to see <math>1 \quad 2 \quad \dots \quad (n-1)</math></p>	<p>B1, B1</p> <p>M1</p> <p>A1</p> <p>depM1</p> <p>A1 <b>(6)</b></p> <p><b>[9]</b></p> <p>M1</p> <p>B1, + [A1]</p> <p>B1,</p> <p>depM1</p> <p>A1</p>
--	--	---



<p>6.</p>	<p>(a) <math>f(2.0) = -0.30685\dots\dots = -0.3069</math>                      AWR T    3 d.p.  <math>f(2.5) = 0.41629\dots\dots\dots = 0.4163</math>                      both correct 4 d.p.          States change of sign, so root (between 2 and 2.5)</p> <p>Note:          B1 gained if candidate's 2 values do show a change of sign and statement made</p> <p>(b) <math>\alpha = (2) + \frac{ f(2) }{ f(2)  +  f(2.5) } \times 0.5</math>    or    <math>\frac{\alpha - 2}{2.5 - \alpha} = \frac{ f(2.0) }{ f(2.5) }</math> or equivalent          Or <math>\frac{x}{ f(2) } = \frac{0.5 - x}{ f(2.5) }</math> and <math>x</math> found  <math>= 2.212</math>    AWR T</p> <p>(c) <math>f(2.25) = 0.06093\dots\dots (\geq 3 \text{ d.p.})</math> [ Allow ln.2.25 + 2.25 - 3]  <math>f'(x) = \frac{1}{x} + 1,</math>                      <math>f'(2.25) = 1.4</math> or <math>1\frac{4}{9}</math> or <math>\frac{13}{9}</math> (allow 1.444)  <math>\alpha = 2.25 - \frac{f(2.25)}{f'(2.25)},</math>    = 2.20781.... = 2.208    AWR T</p> <p>(d) <math>f(2.2075) =,</math>                      <math>\{-6.3\dots\dots \times 10^{-4}\}</math>  <math>f(2.2085) =,</math>                      <math>\{8.1\dots\dots \times 10^{-4}\}</math></p> <p><math>\therefore</math> Correct values (<math>\geq 1</math> s.f.), (root in interval) so root is 2.208 to 3 d.p.</p>	<p>M1          A1          B1    <b>(3)</b></p> <p>M1          A1    <b>(2)</b></p> <p>B1          M1,A1          M1A1 <b>(5)</b></p> <p>M1          A1 <b>(2) [12]</b></p>
	<p>Notes:</p> <p>c) First M in (c) is just for <math>\frac{1}{x} + 1</math>          If no intermediate values seen B1M1A1M1A0 is possible for 2.209 or 2.21,          otherwise as scheme (B1 eased to award this if not evaluated)</p> <p>(d) A1 requires values correct (<math>\geq 1</math> s.f.) and statement (need not say change of sign)          M can be given for candidate's <math>f(2.2075)</math> and <math>f(2.2085)</math></p> <p>Allow N-R applied at least twice more, but A1 requires 2.20794 or better and statement</p> <p>MR in (c) 2.5 instead of 2.25 (Answer 2.203) award on ePen B0M1A0M1A1</p>	

<p>7.</p>	<p>(a) <math>y = x^{-2} \Rightarrow \frac{dy}{dt} = -2x^{-3} \frac{dx}{dt}</math> [Use of chain rule; need <math>\frac{dx}{dt}</math>]</p> $\Rightarrow \frac{d^2y}{dt^2} = -2x^{-3} \frac{d^2x}{dt^2}, + 6x^{-4} \left(\frac{dx}{dt}\right)^2$ <p>(<math>\div</math> given d.e. by <math>x^4</math>) <math>\frac{2}{x^3} \frac{d^2x}{dt^2} - \frac{6}{x^4} \left(\frac{dx}{dt}\right)^2 = \frac{1}{x^2} - 3</math></p> <p>becomes <math>(-\frac{d^2y}{dt^2} = y - 3) \quad \frac{d^2y}{dt^2} + y = 3</math> <b>AG</b></p> <p>(b) Auxiliary equation: <math>m^2 + 1 = 0</math> and produce Complementary Function <math>y = \dots</math></p> $(y) = A \cos t + B \sin t$ <p>Particular integral: <math>y = 3</math></p> <p><math>\therefore</math> General solution: <math>(y) = A \cos t + B \sin t + 3</math></p> <p>(c) <math>\frac{1}{x^2} = A \cos t + B \sin t + 3</math></p> $x = \frac{1}{2}, t = 0 \Rightarrow (4 = A + 3) \quad A = 1$ <p>Differentiating (to include <math>\frac{dx}{dt}</math>): <math>-2x^{-3} \frac{dx}{dt} = -A \sin t + B \cos t</math></p> $\frac{dx}{dt} = 0, t = 0 \Rightarrow (0 = 0 + B) \quad B = 0$ $\therefore \frac{1}{x^2} = 3 + \cos t \quad \text{so} \quad x = \frac{1}{\sqrt{3 + \cos t}}$ <p>(d) (Max. value of <math>x</math> when <math>\cos t = -1</math>) so <math>\max x = \frac{1}{\sqrt{2}}</math> or AWRT 0.707</p>	<p>M1</p> <p>A1√, M1A1</p> <p>A1 cso <b>(5)</b></p> <p>M1</p> <p>A1cao</p> <p>B1</p> <p>A1√ <b>(4)</b></p> <p>B1</p> <p>M1</p> <p>M1</p> <p>A1cao <b>(4)</b></p> <p>B1 <b>(1)</b> <b>[14]</b></p>
	<p>Notes: (See separate sheet for several variations)</p> <p>(a) Second M1 is for attempt at product rule. (be generous) Final A1 requires all working correct and sufficient “substitution” work</p> <p>(b) Answer can be stated; M1 is implied by correct C.F. stated (allow <math>\theta</math> for <math>t</math>) A1 f.t. for candidates CF + PI Allow <math>m^2 + m = 0</math> and <math>m^2 - 1 = 0</math> for M1. Marks for (b) can be gained in</p> <p>(c)</p> <p>(b) Second M : complete method to find other constant (This may involve solving two equations in A and B)</p>	



<p>8.</p>	<p>(a) <math>x = r \cos \theta = 4 \sin \theta \cos^3 \theta</math>  <math>\frac{dx}{d\theta} = 4 \cos^4 \theta - 12 \cos^2 \theta \sin^2 \theta</math> any correct expression  Solving <math>\frac{dx}{d\theta} = 0</math> <math>[\frac{dx}{d\theta} = 0 \Rightarrow 4 \cos^2 \theta (\cos^2 \theta - 3 \sin^2 \theta) = 0]</math>  <math>\sin \theta = \frac{1}{2}</math> or <math>\cos \theta = \frac{\sqrt{3}}{2}</math> or <math>\tan \theta = \frac{1}{\sqrt{3}} \Rightarrow \theta = \frac{\pi}{6}</math> AG  <math>r = 4 \sin \frac{\pi}{6} \cos^2 \frac{\pi}{6} = \frac{3}{2}</math> AG  (b) <math>A = \frac{1}{2} \int_{\frac{\pi}{6}}^{\frac{\pi}{4}} r^2 d\theta = \frac{1}{2} \cdot 16 \int_{\frac{\pi}{6}}^{\frac{\pi}{4}} \sin^2 \theta \cos^4 \theta d\theta</math>  <math>8 \sin^2 \theta \cos^4 \theta = 2 \cos^2 \theta (4 \sin^2 \theta \cos^2 \theta) = 2 \cos^2 \theta \sin^2 2\theta</math>  <math>= (\cos 2\theta + 1) \sin^2 2\theta</math>  <math>= \cos 2\theta \sin^2 2\theta + \frac{1 - \cos 4\theta}{2} = \text{Answer}</math> AG  (c) Area = <math>\left[ \frac{1}{6} \sin^3 2\theta + \frac{\theta}{2} - \frac{\sin 4\theta}{8} \right]_{\left(\frac{\pi}{6}\right)}^{\left(\frac{\pi}{4}\right)}</math> (ignore limits)  <math>= \left( \frac{1}{6} \sin^3 \frac{\pi}{2} + \frac{\pi}{8} - \frac{\sin \pi}{8} \right) - \left( \frac{1}{6} \sin^3 \frac{\pi}{3} + \frac{\pi}{12} - \frac{\sin \frac{2\pi}{3}}{8} \right)</math> (sub. limits)  <math>= \left( \frac{1}{6} + \frac{\pi}{8} \right) - \left( \frac{\sqrt{3}}{16} + \frac{\pi}{12} - \frac{\sqrt{3}}{16} \right) = \frac{1}{6} + \frac{\pi}{24}</math> both cao</p>	<p>M1  M1A1  M1  A1 cso  A1 cso (6)  M1  M1  A1 cso (3)  M1A1  M1  A1,A1 (5)  [14]</p>
	<p>Notes:  (a) So many ways <math>x</math> may be expressing e.g  <math>2 \sin 2\theta \cos^2 \theta, \sin 2\theta(1 + \cos 2\theta), \sin 2\theta + (1/2) \sin 4\theta</math>  leading to many results for <math>\frac{dx}{d\theta}</math>  Some relevant equations in solving  <math>[(1 - 4 \sin^2 \theta) = 0, (4 \cos^2 \theta - 3) = 0, (1 - 3 \tan^2 \theta) = 0, \cos 3\theta = 0]</math>  Showing that <math>\theta = \frac{\pi}{6}</math> satisfies <math>\frac{dx}{d\theta} = 0</math>, allow M1A1 providing <math>\frac{dx}{d\theta}</math> correct  Starting with <math>x = r \sin \theta</math> can gain M0M1M1 in (a)  (b) First M1 for use of double angle formula for <math>\sin 2A</math>  Second M1 for use of <math>\cos 2A = 2 \cos^2 A - 1</math>  Answer given: must be intermediate step, as shown, and no incorrect work  (c) For first M, of the form <math>a \sin^3 2\theta + \frac{\theta}{2} \pm b \sin 4\theta</math> (Allow if two of correct form)  On ePen the order of the As in answer is as written</p>	