





Question 1 continued

Lined writing area for the answer.


(Total 6 marks)

Q1



2.

Figure 1



A uniform solid right circular cone has base radius  $a$  and semi-vertical angle  $\alpha$ , where  $\tan \alpha = \frac{1}{3}$ . The cone is freely suspended by a string attached at a point  $A$  on the rim of its base, and hangs in equilibrium with its axis of symmetry making an angle of  $\theta^\circ$  with the upward vertical, as shown in Figure 1.

Find, to one decimal place, the value of  $\theta$ .

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

Question 2 continued

Lined writing area for Question 2 continued.

(Total 5 marks)

Q2




N 2 3 8 2 2 A 0 5 2 4





4.

**Figure 2**



A particle  $P$  of mass  $m$  is attached to one end of a light inextensible string of length  $a$ . The other end of the string is attached to a point  $O$ . The point  $A$  is vertically below  $O$ , and  $OA = a$ . The particle is projected horizontally from  $A$  with speed  $\sqrt{3ag}$ . When  $OP$  makes an angle  $\theta$  with the upward vertical through  $O$  and the string is still taut, the tension in the string is  $T$  and the speed of  $P$  is  $v$ , as shown in Figure 2.

(a) Find, in terms of  $a$ ,  $g$  and  $\theta$ , an expression for  $v^2$ . (3)

(b) Show that  $T = (1 - 3 \cos \theta)mg$ . (3)

The string becomes slack when  $P$  is at the point  $B$ .

(c) Find, in terms of  $a$ , the vertical height of  $B$  above  $A$ . (2)

After the string becomes slack, the highest point reached by  $P$  is  $C$ .

(d) Find, in terms of  $a$ , the vertical height of  $C$  above  $B$ . (5)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---





**Question 4 continued**

Lined writing area for the answer to Question 4.



**Question 4 continued**


A large rectangular area consisting of approximately 35 horizontal lines for writing.





5.

Figure 3



One end of a light inextensible string is attached to a fixed point  $A$ . The other end of the string is attached to a fixed point  $B$ , vertically below  $A$ , where  $AB = h$ . A small smooth ring  $R$  of mass  $m$  is threaded on the string. The ring  $R$  moves in a horizontal circle with centre  $B$ , as shown in Figure 3. The upper section of the string makes a constant angle  $\theta$  with the downward vertical and  $R$  moves with constant angular speed  $\omega$ . The ring is modelled as a particle.

(a) Show that  $\omega^2 = \frac{g}{h} \left( \frac{1 + \sin \theta}{\sin \theta} \right)$ . (7)

(b) Deduce that  $\omega > \sqrt{\frac{2g}{h}}$ . (2)

Given that  $\omega = \sqrt{\frac{3g}{h}}$ ,

(c) find, in terms of  $m$  and  $g$ , the tension in the string. (4)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---







**Question 5 continued**

Lined writing area for the answer to Question 5.


**Q5**

**(Total 13 marks)**



6.

Figure 4




The shaded region  $R$  is bounded by the curve with equation  $y = \frac{1}{2x^2}$ , the  $x$ -axis and the lines  $x = 1$  and  $x = 2$ , as shown in Figure 4. The unit of length on each axis is 1 m. A uniform solid  $S$  has the shape made by rotating  $R$  through  $360^\circ$  about the  $x$ -axis.

(a) Show that the centre of mass of  $S$  is  $\frac{2}{7}$  m from its larger plane face.

(6)

Figure 5



A sporting trophy  $T$  is a uniform solid hemisphere  $H$  joined to the solid  $S$ . The hemisphere has radius  $\frac{1}{2}$  m and its plane face coincides with the larger plane face of  $S$ , as shown in Figure 5. Both  $H$  and  $S$  are made of the same material.

(b) Find the distance of the centre of mass of  $T$  from its plane face.

(7)

---



---



---



---









### Question 6 continued

Lined writing area for student response.

(Total 13 marks)

Q6



7. A particle  $P$  of mass  $0.25$  kg is attached to one end of a light elastic string. The string has natural length  $0.8$  m and modulus of elasticity  $\lambda$  N. The other end of the string is attached to a fixed point  $A$ . In its equilibrium position,  $P$  is  $0.85$  m vertically below  $A$ .

(a) Show that  $\lambda = 39.2$ .

(2)

The particle is now displaced to a point  $B$ ,  $0.95$  m vertically below  $A$ , and released from rest.

(b) Prove that, while the string remains stretched,  $P$  moves with simple harmonic motion of period  $\frac{\pi}{7}$  s.

(6)

(c) Calculate the speed of  $P$  at the instant when the string first becomes slack.

(3)

The particle first comes to instantaneous rest at the point  $C$ .

(d) Find, to 3 significant figures, the time taken for  $P$  to move from  $B$  to  $C$ .

(5)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



Question 7 continued

Lined writing area for the answer to Question 7.



**Question 7 continued**

Lined writing area for the answer to Question 7.



Question 7 continued

Lined area for writing the answer to Question 7.

(Total 16 marks)

TOTAL FOR PAPER: 75 MARKS

END

Q7



**BLANK PAGE**

