

Mark Scheme (Results)


Summer 2007


GCE


GCE Mathematics

Mechanics M4 (6680)


June 2007
6680 Mechanics M4
Mark Scheme

Question Number	Scheme	Marks
1(a)	 $u \cos 60^\circ = v \cos 30^\circ$ $u = v\sqrt{3}$ $\text{KE lost} = \frac{1}{2}m(u^2 - v^2)$ $\text{Fraction of KE lost} = 1 - \left(\frac{v}{u}\right)^2$ $= 1 - \frac{1}{3} = \frac{2}{3} \text{ or at least 3sf ending in 7}$ $\text{or } \frac{3}{4}(1 - e^2)$	<p>M1A1 A1</p> <p>M1</p> <p>DM1</p> <p>A1</p> <p style="text-align: right;">(6)</p>
(b)	$e = \frac{v \sin 30^\circ}{u \sin 60^\circ}$ $= \frac{v}{u} \cdot \frac{1}{\sqrt{3}}$ $= \frac{1}{3}$	<p>M1A1</p> <p>DM1</p> <p>A1</p> <p style="text-align: right;">(4)</p>
a)	<p>M1 Resolve parallel to the wall <i>Alt: reasonable attempt at equation connecting two variables</i></p> <p>A1 Correct as above or equivalent <i>equation correct</i></p> <p>A1 u in terms of v or $v.v.$ - not necessarily simplified. <i>or ratio of the two variables correct</i></p> <p>M1 expression for KE lost</p> <p>DM1 expression in one variable for fraction of KE lost – could be u/v as above</p> <p>A1 cao</p>	<p><i>The first three marks can be awarded in (b) if not seen in (a)</i></p>
b)	<p>M1 Use NIL perpendicular to the wall and form equation in e</p> <p>A1 Correct unsimplified expression as above or $eu \sin 60^\circ = v \sin 30^\circ$ or equivalent</p> <p>DM1 Substitute values for trig functions or use relationship from (a) and rearrange to $e = \dots$</p> <p>A1 cao accept decimals to at least 3sf</p>	<p><i>The first two marks can be awarded in (a)</i></p>

<p>2(a)</p> <p>(b)</p>	 <p> $F = \frac{Ru}{v}$ $R(\square), \frac{Ru}{v} - R = M \frac{dv}{dt}$ $R(u - v) = Mv \frac{dv}{dt} *$ </p> <p> $\int_0^T dt = \frac{M}{R} \int_{\frac{1}{4}U}^{\frac{1}{3}U} \frac{v dv}{u - v}$ $\square T = \frac{M}{R} \int_{\frac{1}{4}U}^{\frac{1}{3}U} -1 + \frac{u}{u - v} dv$ $\square \frac{M}{R} [-v - u \ln(u - v)]_{\frac{1}{4}U}^{\frac{1}{3}U}$ $\square \frac{M}{R} \left[-\frac{u}{3} - u \ln\left(\frac{2u}{3}\right) + \frac{u}{4} + u \ln\left(\frac{3u}{4}\right) \right] \left(C = -\frac{Mu}{R} \left(\ln \frac{3u}{4} + \frac{1}{4} \right) \right)$ $\square \frac{Mu}{R} \left(-\frac{1}{12} + \ln \frac{9}{8} \right)$ <p>Hence $k = \ln \frac{9}{8} - \frac{1}{12}$</p> </p>	<p>B1</p> <p>M1</p> <p>A1</p> <p>(3)</p> <p>M1A1</p> <p>DM1</p> <p>A1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>(7)</p>
<p>a)</p> <p>b)</p>	<p>B1 Correct expression involving the driving force.</p> <p>M1 Use of $F = ma$ to form a differential equation. Condone sign errors. a must be expressed as a derivative, but could be any valid form.</p> <p>A1 Rearrange to given form.</p> <p>M1 Separate the variables</p> <p>A1 Separation correct (limits not necessarily seen at this stage)</p> <p>DM1 Attempt a complete integration process</p> <p>A1 Integration correct</p> <p>M1 Correct use of both limits – substitute and subtract. Condone wrong order.</p> <p>M1 Simplify to find k from an expression involving a logarithm</p> <p>A1 Answer as given, or exact equivalent. Need to see $k = \ln A + B$</p>	

Question Number	Scheme	Marks
<p>3. (a)</p> <p>(b)</p> <p>(c)</p>	 <p> $V = -mga \cos \theta - mg(2a \cos \theta + a \sin \theta)$ $= -mga(3 \cos \theta + \sin \theta) \quad (+const) *$ </p> <p> $\frac{dV}{d\theta} = -mga(-3 \sin \theta + \cos \theta)$ </p> <p> $= 0 \quad \square \quad \tan \theta = \frac{1}{3}$ </p> <p> $\square \quad \square = 0.32(1)^\circ \text{ or } 18.4^\circ \text{ accept awrt}$ </p> <p> $\frac{d^2V}{d\theta^2} = -mga(-3 \cos \theta - \sin \theta)$ $= mga(3 \cos \theta + \sin \theta)$ </p> <p> Hence, when $\square = 0.32^\circ$, $\frac{d^2V}{d\theta^2} > 0$ i.e. stable </p>	<p>M1A1A1 A1 (4)</p> <p>M1A1</p> <p>M1</p> <p>A1 (4)</p> <p>M1A1</p> <p>M1 A1 (4)</p>
<p>a)</p> <p>b)</p> <p>c)</p>	<p>M1 Expression for the potential energy of the two rods. Condone trig errors. Condone sign errors. BC term in two parts</p> <p>A1 correct expression for AB</p> <p>A1 correct expression for BC</p> <p>A1 Answer <u>as given</u> .</p> <p>M1 Attempt to differentiate V. Condone errors in signs and in constants.</p> <p>A1 Derivative correct</p> <p>M1 Set derivative = 0 and rearrange to a single trig function in \square</p> <p>A1 Solve for \square</p> <p>or M1A1 find the position of the center of mass</p> <p>M1A1 form and solve trig equation for \square</p> <p>M1 Differentiate to obtain the second derivative</p> <p>A1 Derivative correct</p> <p>M1 Determine the sign of the second derivative</p> <p>A1 Correct conclusion. cso</p> <p>Or: M1 Find the value of $\frac{dV}{d\theta}$ on both sides of the minimum point</p> <p>A1 signs correct</p> <p>M1 Use the results to determine the nature of the turning point</p> <p>A1 Correct conclusion, cso.</p>	<p><i>These 4 marks are dependent on the use of derivatives</i></p>

4 (a)



Fix A

$$v_{\min} = 15 \sin 50^\circ$$

$$= 11.5 \text{ km h}^{-1} \text{ (3 s.f.)}$$

or: triangle without the right angle

identified and $\frac{15}{\sin \theta} = \frac{v_B}{\sin 50}$


$$\Rightarrow v_B = \frac{15 \sin 50}{\sin \theta}$$

minimum value $\Rightarrow \theta = 90$ for M1

As above for A1A1

M1A1
A1
(3)

(b)



Ambiguous Sine Rule:

2 possible solutions for \square

B1B1
(2)

$$\frac{\sin \alpha}{15} = \frac{\sin 50}{13}$$

$$\square = 62.1^\circ \text{ (or } 118^\circ)$$

(smaller value gives larger relative velocity)

\square either

$$v = 13 \cos 62.1 + 15 \cos 50 = 15.72 \text{ kmh}^{-1}$$

M1A1

A1

M1A1

Or

$$v^2 = 15^2 + 13^2 - 390 \cos 67.9 = 247.27$$

$$v = 15.7 \text{ kmh}^{-1}$$

M1
A1

$$\text{Time} = \frac{20}{\text{their } 15.72 \dots}$$

$$= 1.272 \dots \text{ hrs}$$

M1 A1

Earliest time is 13.16 hrs or 13.17 hrs
accept 1.16 (pm) or 1.17 (pm)

A1
(8)

(c)

a) M1 Velocity of B relative to A is in the direction of the line joining AB.
 Minimum V requires a right angled triangle.
 Convincing attempt to find the correct side.
 A1 $15 \times \sin(\text{their } \hat{})$
 A1 Q specifies 3sf, so 11.5 only

b) B1B1 Convincing argument
 B1B0 Argument with some merit


c) M1 Use of Sine Rule
 A1 Correct expression
 A1 (2 possible values,) pick the correct value.
 M1 Use trig. to form an equation in v
 A1 correct equation
 M1 $time = \frac{distance}{speed}$
 A1ft correct expression with their v (not necessarily evaluated)
 A1 correct time in hours & minutes

Or:

M1 Use of cosine rule
 A1 $13^2 = 15^2 + v^2 - 2 \times 15 \times v \times \cos 50$
 A1 (Award after the next two marks) 15.72 or awrt 15.72
 M1 Attempt to solve the equation for v
 A1
$$\frac{30 \cos 50 \pm \sqrt{(30 \cos 50)^2 - 4 \times 56}}{2}$$

 (15.72 or 3.562)
 Finish as above

5. (a)



CLM: $2v_2 - v_1 = 1 - 2 = -1$

NIL: $1 + 1 = \frac{1}{2}(v_1 + v_2)$

$\square v_2 = 1, v_1 = 3$

Horizontal components unchanged (i.e. 2 & 3)
 $\mathbf{v}_A = 3\mathbf{i} + \mathbf{j}; \mathbf{v}_B = 2\mathbf{i} - 3\mathbf{j}$

Dependent on both M's above

Independent of all other marks

M1A1
M1A1

DM1

A1
A1

(7)

(b)

For B: $I = m(1 - (-3)) = 4m$

(Or For A: $-I = 2m(-1 - 1) \square I = 4m$)

M1A1

(2)

(c)

$\begin{pmatrix} 3 \\ 1 \end{pmatrix} \cdot \begin{pmatrix} 3 \\ -1 \end{pmatrix} = \sqrt{3^2 + 1^2} \cdot \sqrt{3^2 + (-1)^2} \cos \theta$

$\Rightarrow 8 = 10 \cos \theta$


$\square = 37^\circ$

M1A1

M1
A1

Alternative:

(4)




where $\tan \theta = \frac{1}{3} \square 1$

$\square 1$

required angle is $2\square$

M1A1

<p>a)</p> <p>b)</p> <p>c)</p>	<p>M1 Conservation of momentum along the line of centres. Condone sign errors A1 equation correct</p> <p>M1 Impact law along the line of centres. e must be used correctly, but condone sign errors. A1 equation correct. The signs need to be consistent between the two equations</p> <p>M1 Solve the simultaneous equations for their v_1 and v_2. A1 \mathbf{i} components correct – independent mark A1 \mathbf{v}_A & \mathbf{v}_B correct</p> <p>M1 Impulse = change in momentum for one sphere. Condone order of subtraction. A1 Magnitude correct.</p> <p>M1 Any complete method to find the trig ratio of a relevant angle. A1 $\cos\theta = \frac{4}{5}$, $\tan\frac{\theta}{2} = \frac{1}{3}$, ...</p> <p>Or M1 find angle of approach to the line of centres and angle after collision. A1 values correct. (both 71.56</p> <p>M1 solve for \square A1 37° (Q specifies nearest degree)</p> <p>Special case: candidates who act as if the line of centres is in the direction of \mathbf{i}:</p> <p>CLM $u+2v = 8$ NIL $v-u = 2$</p> <p>$u=4/3, v=10/3$</p> <p>$4/3\mathbf{i} + \mathbf{j}; 10/3\mathbf{i} - \mathbf{j}$</p> <p>Impulse $2m-4/3m = 2/3m$</p> $\frac{10+1}{\sqrt{10}\sqrt{\frac{109}{9}}} = \cos\theta \quad \square\square 1.70^\circ$ <p>Work is equivalent, so treat as a MR: M1A0M1A0M1A1A1 M1A1 M1A1M1A1</p>	
-------------------------------	---	--

6 (a)	 <p>At E, $\frac{2.45e}{0.5} = 0.1g$ $\square e=0.2$</p> <p>\square $0.5(ori) + 0.2 + y = 2 \sin 2t + 0.5(ori) + x$</p> <p>$\square 0.2 + y = 2 \sin 2t + x^*$</p>	M1 A1 B1 (3)
(b)	$0.1g - T = 0.1\ddot{y}$ $R(\square) \quad 0.1g - \frac{2.45x}{0.5} = 0.1\ddot{y}$ $0.98 - 4.9(0.2 + y - 2 \sin 2t) = 0.1\ddot{y}$ $(-4.9y + 9.8 \sin 2t = 0.1\ddot{y})$ $\square \frac{d^2y}{dt^2} + 49y = 98 \sin 2t^*$	M*1 M1 DM*1A1 A1 cso (5)
(c)	<p>CF is $y = A \cos 7t + B \sin 7t$</p> <p>Hence GS is $y = A \cos 7t + B \sin 7t + \frac{98}{45} \sin 2t$</p> <p>$t = 0, y = 0: 0 = A$ so, $y = B \sin 7t + \frac{98}{45} \sin 2t$</p> $\dot{y} = 7B \cos 7t + \frac{196}{45} \cos 2t$ <p>$t = 0, \dot{y} = 0: 0 = 7B + \frac{196}{45} \quad \square B = -\frac{28}{45}$</p> $\square y = \frac{14}{45}(7 \sin 2t - 2 \sin 7t)$	M1 A1 B1 M1 A1 (5)
(d)	$\dot{y} = \frac{14}{45}(14 \cos 2t - 14 \cos 7t)$ $\dot{y} = 0 \quad \square \cos 2t = \cos 7t$ $\square 7t = 2k\pi \pm 2t$ $k=1 \quad \square 9t = 2 \quad (\text{or } 5t = 2 \quad \square$ $t = \frac{2\pi}{9} \dots \text{accept } 0.698s, 0.70s.$	B1 M1 M1 A1 (4)

<p>a)</p> <p>b)</p> <p>c)</p> <p>d)</p>	<p>M1 Hooke's law to find extension at equilibrium A1 cao B1 Q specifies reference to a diagram. Correct reasoning leading to <u>given answer</u>.</p> <p>M1 Use of F=ma. Weight, tension and acceleration. Condone sign errors. M1 Substitute for tension in terms of x M1 Use given result to substitute for x in terms of y A1 Correct unsimplified equation A1 Rearrange to <u>given form</u> cso.</p> <p>M1 Correct form for CF A1 GS for y correct B1 Deduce coefficient of $\cos \square = 0$ M1 Differentiate their y and substitute $t=0$, $\dot{y} = 0$ A1 y in terms of t. Any exact equivalent.</p> <p>B1 \dot{y} correct M1 set $\dot{y} = 0$ M1 solve for general solution for t: $7t = 2k\pi \pm 2t$ or: $\sin \frac{9t}{2} \times \sin \frac{5t}{2} = 0 \Rightarrow \sin \frac{9t}{2} = 0$ or $\sin \frac{5t}{2} = 0$</p> <p>A1 Select smallest value</p>	