








3.



**Figure 1**

Figure 1 shows a sketch of the curve with equation  $y = f(x)$ . The curve passes through the point  $(0, 7)$  and has a minimum point at  $(7, 0)$ .

On separate diagrams, sketch the curve with equation

(a)  $y = f(x) + 3$ , **(3)**

(b)  $y = f(2x)$ . **(2)**

On each diagram, show clearly the coordinates of the minimum point and the coordinates of the point at which the curve crosses the  $y$ -axis.



**Question 3 continued**

Leave  
blank

**(Total 5 marks)**

**Q3**

5

**Turn over**





**Question 4 continued**

Handwritten notes area consisting of approximately 35 horizontal lines for writing.

**(Total 5 marks)**

Leave  
blank

**Q4**

7

**Turn over**







**Question 5 continued**

Leave blank

Lined area for writing the answer to Question 5 continued.

**(Total 6 marks)**

**Q5**




Leave  
blank

6. The curve  $C$  has equation  $y = \frac{3}{x}$  and the line  $l$  has equation  $y = 2x + 5$ .

(a) On the axes below, sketch the graphs of  $C$  and  $l$ , indicating clearly the coordinates of any intersections with the axes. (3)

(b) Find the coordinates of the points of intersection of  $C$  and  $l$ . (6)





7. Sue is training for a marathon. Her training includes a run every Saturday starting with a run of 5 km on the first Saturday. Each Saturday she increases the length of her run from the previous Saturday by 2 km.

(a) Show that on the 4th Saturday of training she runs 11 km. (1)

(b) Find an expression, in terms of  $n$ , for the length of her training run on the  $n$ th Saturday. (2)

(c) Show that the total distance she runs on Saturdays in  $n$  weeks of training is  $n(n + 4)$  km. (3)

On the  $n$ th Saturday Sue runs 43 km.

(d) Find the value of  $n$ . (2)

(e) Find the total distance, in km, Sue runs on Saturdays in  $n$  weeks of training. (2)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---















9. The curve  $C$  has equation  $y = kx^3 - x^2 + x - 5$ , where  $k$  is a constant.

(a) Find  $\frac{dy}{dx}$ . **(2)**

The point  $A$  with  $x$ -coordinate  $-\frac{1}{2}$  lies on  $C$ . The tangent to  $C$  at  $A$  is parallel to the line with equation  $2y - 7x + 1 = 0$ .

Find

(b) the value of  $k$ , **(4)**

(c) the value of the  $y$ -coordinate of  $A$ . **(2)**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---







**Question 9 continued**

Leave  
blank

*(This section contains 33 horizontal lines for writing answers.)*

**(Total 8 marks)**

**Q9**

21

Turn over



10.




Figure 2

The points  $Q(1, 3)$  and  $R(7, 0)$  lie on the line  $l_1$ , as shown in Figure 2.

The length of  $QR$  is  $a\sqrt{5}$ .

- (a) Find the value of  $a$ . (3)

The line  $l_2$  is perpendicular to  $l_1$ , passes through  $Q$  and crosses the  $y$ -axis at the point  $P$ , as shown in Figure 2.

Find

- (b) an equation for  $l_2$ , (5)
- (c) the coordinates of  $P$ , (1)
- (d) the area of  $\triangle PQR$ . (4)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---





Leave  
blank

**Question 10 continued**

Handwritten response area consisting of 25 horizontal lines.



H 2 9 9 9 2 A 0 2 3 2 8

23

Turn over











Leave  
blank

**Question 11 continued**

Lined area for writing the answer to Question 11.



