

C4 June 2009

$$\begin{aligned}
 \textcircled{1} \quad f(x) &= \frac{1}{\sqrt{4+x}} = (4+x)^{-\frac{1}{2}} \\
 &= \left[4 \left(1 + \frac{x}{4} \right) \right]^{-\frac{1}{2}} = \frac{1}{2} \left(1 + \frac{x}{4} \right)^{-\frac{1}{2}} \\
 &= \frac{1}{2} \left(1 + (-\frac{1}{2})(\frac{x}{4}) + \frac{(-\frac{1}{2})(-\frac{3}{2})(\frac{x}{4})^2}{2!} + \frac{(-\frac{1}{2})(-\frac{3}{2})(-\frac{5}{2})(\frac{x}{4})^3}{3!} + \dots \right) \\
 &= \frac{1}{2} \left(1 - \frac{x}{8} + \frac{3}{128}x^2 - \frac{5}{1024}x^3 + \dots \right) \\
 &= \frac{1}{2} - \frac{x}{16} + \frac{3}{256}x^2 - \frac{5}{2048}x^3 + \dots
 \end{aligned}$$

$$\textcircled{2} \quad y = 3 \cos\left(\frac{x}{3}\right), \quad 0 \leq x \leq \frac{3\pi}{2}.$$

(a)	x	0	$\frac{3\pi}{8}$	$\frac{3\pi}{4}$	$\frac{9\pi}{8}$	$\frac{3\pi}{2}$
	y	3	2.77164	2.12132	1.14805	0

$$\begin{aligned}
 \text{(b) Area of } R &\approx \frac{\left(\frac{3\pi}{8}\right)}{2} \left(3 + 0 + 2(2.77164 + 2.12132 + 1.14805) \right) \\
 &= \underline{\underline{8.884}}
 \end{aligned}$$

$$\begin{aligned}
 \text{(c) Exact area of } R &= \int_0^{\frac{3\pi}{2}} 3 \cos\left(\frac{x}{3}\right) dx = \left[9 \sin\left(\frac{x}{3}\right) \right]_0^{\frac{3\pi}{2}} \\
 &= \underline{\underline{9}}
 \end{aligned}$$

$$(3)(a) \quad f(x) = \frac{4-2x}{(2x+1)(x+1)(x+3)} = \frac{A}{2x+1} + \frac{B}{x+1} + \frac{C}{x+3}$$

$$\Rightarrow 4-2x = A(x+1)(x+3) + B(2x+1)(x+3) + C(2x+1)(x+1).$$

$$x=-1 \Rightarrow 6 = -2B \Rightarrow B = -3$$

$$x=-3 \Rightarrow 10 = 10C \Rightarrow C=1$$

$$x = -\frac{1}{2} \Rightarrow 5 = \frac{5}{4}A \Rightarrow A=4$$

$$(b)(i) \quad \int f(x) dx = \int \frac{4}{2x+1} - \frac{3}{x+1} + \frac{1}{x+3} dx$$

$$= 4 \frac{\ln|2x+1|}{2} - 3 \ln|x+1| + \ln|x+3| + C.$$

$$= \underline{2 \ln|2x+1| - 3 \ln|x+1| + \ln|x+3| + C}$$

$$(ii) \quad \int_0^2 f(x) dx = (2 \ln 5 - 3 \ln 3 + \ln 5) - (\ln 3)$$

$$= 3 \ln 5 - 4 \ln 3 = \underline{\underline{\ln \frac{125}{81}}}$$

$$(4) \text{ (a)} \quad y e^{-2x} = 2x + y^2$$

Differentiate both sides w.r.t. x

$$\Rightarrow y(-2e^{-2x}) + \frac{dy}{dx} \cdot e^{-2x} = 2 + 2y \frac{dy}{dx}$$

$$\Rightarrow \frac{dy}{dx} (e^{-2x} - 2y) = 2 + 2ye^{-2x}$$

$$\Rightarrow \frac{dy}{dx} = \frac{2 + 2ye^{-2x}}{e^{-2x} - 2y}$$

$$(b) \quad P(0, 1).$$

$$\text{At } P, \quad \frac{dy}{dx} = \frac{2 + 2}{1 - 2} = -4$$

\Rightarrow gradient of normal is $\frac{1}{4}$.

$$\text{Equation of normal is } y - 1 = \frac{1}{4}(x - 0)$$

$$\Rightarrow \underline{x - 4y + 4 = 0}$$

$$(5) \text{ (a)} \quad x = 2 \cos 2t, \quad y = 6 \sin t, \quad 0 \leq t \leq \frac{\pi}{2}.$$

$$\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{\frac{6 \cos t}{-4 \sin 2t}}{\frac{2 \cos 2t}{2 \sin 2t}} = \frac{-3 \cos t}{2 \sin 2t}$$

$$\text{When } t = \frac{\pi}{3}, \quad \frac{dy}{dx} = \frac{-3 \cos \frac{\pi}{3}}{2 \sin \frac{2\pi}{3}} = \frac{-\frac{\sqrt{3}}{2}}{\underline{\underline{2}}}$$

$$(b) \quad x = 2 \cos 2t = 2(1 - 2 \sin^2 t) = 2 - 4 \sin^2 t$$

$$= 2 - 4 \left(\frac{y}{6}\right)^2 = 2 - \frac{y^2}{9} \Rightarrow y^2 = 18 - 9x$$

From sketch we want +ve square root,

$$\text{so } y = \sqrt{18 - 9x}, \quad -2 \leq x \leq 2 \quad \begin{array}{l} \text{N.B. } 0 \leq t \leq \frac{\pi}{2} \\ \Rightarrow 0 \leq 2t \leq \pi \end{array}$$

$$\Rightarrow -1 \leq \cos 2t \leq 1 \Rightarrow -2 \leq 2 \cos 2t \leq 2.$$

(c) Range is . $0 \leq f(x) \leq 6$

Since maximum value is $\sqrt{18 - 9(-2)} = 6$.

$$\begin{aligned} (6)(a) \int \sqrt{5-x} \, dx &= \int (5-x)^{\frac{1}{2}} \, dx \\ &= \frac{(5-x)^{\frac{3}{2}}}{\frac{3}{2}(-1)} + C = -\frac{2}{3}(5-x)^{\frac{3}{2}} + C \end{aligned}$$

$$\begin{aligned} (b) (i) \int (x-1)\sqrt{5-x} \, dx &\quad \text{Using integration by parts,} \\ &= (x-1)\left(-\frac{2}{3}(5-x)^{\frac{3}{2}}\right) - \int 1 \cdot \left(-\frac{2}{3}(5-x)^{\frac{3}{2}}\right) \, dx \\ &= -\frac{2}{3}(x-1)(5-x)^{\frac{3}{2}} + \frac{2}{3} \int (5-x)^{\frac{3}{2}} \, dx \\ &= -\frac{2}{3}(x-1)(5-x)^{\frac{3}{2}} - \frac{4}{15}(5-x)^{\frac{5}{2}} + C \end{aligned}$$

$$\begin{aligned} \text{Method 2} \quad \int (x-1)\sqrt{5-x} \, dx &\quad \text{Put } u^2 = 5-x \\ &\quad 2u \frac{du}{dx} = -1 \\ &\quad dx = -2u \, du \\ &= \int (4-u^2)(u)(-2u) \, du \\ &= \int 2u^4 - 8u^2 \, du \\ &= \frac{2u^5}{5} - \frac{8u^3}{3} + C \\ &= \frac{2}{5}(5-x)^{\frac{5}{2}} - \frac{8}{3}(5-x)^{\frac{3}{2}} + C \end{aligned}$$

$$\begin{aligned}
 & \text{(ii)} \int_1^5 (x-1)\sqrt{5-x} dx \\
 &= \left[-\frac{2}{3}(x-1)(5-x)^{\frac{3}{2}} - \frac{4}{15}(5-x)^{\frac{5}{2}} \right]_1^5 \\
 &= 0 - \left(0 - \frac{4}{15}(4^{\frac{5}{2}}) \right) = \frac{4}{15} \times 2^5 = \frac{128}{15}
 \end{aligned}$$

Method 2


$$\text{Using } u^2 = 5-x, \quad x=1 \Rightarrow u=2 \\
 x=5 \Rightarrow u=0$$

$$\begin{aligned}
 & \rightarrow \int_2^0 \left[\frac{2}{5}u^5 - \frac{8}{3}u^3 \right] = 0 - \left(\frac{2}{5} \times 2^5 - \frac{8}{3} \times 2^3 \right) \\
 &= \frac{64}{3} - \frac{64}{5} = \frac{128}{15}
 \end{aligned}$$

$$\textcircled{7} \quad \text{(a) Equation of } l : \underline{l} = \begin{pmatrix} 8 \\ 13 \\ -2 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ 1 \\ -2 \end{pmatrix}$$

$$\begin{aligned}
 & \text{(b) } |\overrightarrow{CB}| = |\underline{l} - \underline{c}| = \left| \begin{pmatrix} 1 \\ 5 \\ -10 \end{pmatrix} \right| = \sqrt{1^2 + 5^2 + (-10)^2} \\
 &= \sqrt{126} = \underline{3\sqrt{14}}
 \end{aligned}$$

(C)



$$\cos \theta = \frac{\vec{BC} \cdot \vec{BA}}{|\vec{BC}| |\vec{BA}|} = \frac{\begin{pmatrix} -1 \\ -5 \\ 10 \end{pmatrix} \cdot \begin{pmatrix} -2 \\ -1 \\ 2 \end{pmatrix}}{3\sqrt{14} \times 3}$$

$$= \frac{27}{9\sqrt{14}} \Rightarrow \theta = 36.7^\circ.$$

(d) CX is shortest distance from C to the line,

where

$$\frac{CX}{CB} = \sin \theta \Rightarrow CX = 3\sqrt{14} \times \sin \theta$$

$$= \underline{6.708}$$

$$\begin{aligned} (\text{e}) \quad \text{Area of } \triangle CXB &= \frac{1}{2} \cdot CX \cdot CB \cdot \sin(90 - \theta) \\ &= \frac{1}{2} \times (6.708 \dots) \times 3\sqrt{14} \times \sin(90 - \theta) \\ &= \underline{30.2 \text{ units}^2} \end{aligned}$$

$$\textcircled{8} \quad (a) \quad \int \sin^2 \theta \, d\theta = \int \frac{1}{2}(1 - \cos 2\theta) \, d\theta$$

$$= \frac{1}{2} \left(\theta - \frac{\sin 2\theta}{2} \right) + C = \underline{\underline{\frac{\frac{1}{2}\theta - \frac{1}{4}\sin 2\theta + C}{}}}$$

$$(b) \quad x = \tan \theta, \quad y = 2 \sin 2\theta, \quad 0 \leq \theta < \frac{\pi}{2}.$$

$$V = \int_0^{\frac{\sqrt{3}}{2}} \pi y^2 \, dx$$

$$x = 0 \Rightarrow \theta = 0$$

$$x = \frac{1}{\sqrt{2}} \Rightarrow \theta = \frac{\pi}{4}$$

$$\text{So } V = \int_0^{\frac{\pi}{6}} \pi y^2 \frac{dx}{d\theta} \, d\theta$$

$$= \int_0^{\frac{\pi}{6}} \pi (4 \sin^2 2\theta) \sec^2 \theta \, d\theta$$

$$= \int_0^{\frac{\pi}{6}} \pi \left(4 (2 \sin \theta \cos \theta)^2 \right) \times \frac{1}{\cos^2 \theta} \, d\theta$$

$$= \int_0^{\frac{\pi}{6}} 16\pi \sin^2 \theta \, d\theta \quad \text{so } k = 16\pi.$$

$$(c) \quad = \left[16\pi \left(\frac{1}{2}\theta - \frac{1}{4}\sin 2\theta \right) \right]_0^{\frac{\pi}{6}}$$

$$= 16\pi \left(\left(\frac{\pi}{12} - \frac{\sqrt{3}}{8} \right) - 0 \right) = \underline{\underline{\frac{4\pi^2 - 2\pi\sqrt{3}}{3}}}$$