

June 2006
6675 Further Pure Mathematics FP2
Mark Scheme

Question number	Scheme Marks	
1.	$5\left(\frac{e^x + e^{-x}}{2}\right) - 2\left(\frac{e^x - e^{-x}}{2}\right) = 11$ $3e^{2x} - 22e^x + 7 = 0 \quad \text{M: Simplify to form } \underline{\text{quadratic}} \text{ in } e^x$ $(3e^x - 1)(e^x - 7) = 0 \quad e^x = \frac{1}{3}, \quad e^x = 7 \quad \text{M: Solve 3 term quadratic.}$ $x = \ln \frac{1}{3} \text{ (or } -\ln 3\text{)} \quad x = \ln 7$	B1 M1 A1 M1 A1 A1 (6) 6 Marks
2. (a)	Using $b^2 = a^2(1-e^2)$ or equiv. to find e or ae : ($a = 2$ and $b = 1$) $e = \frac{\sqrt{3}}{2}$	M1 A1
(b)	Using $y^2 = 4(ae)x$ $y^2 = 4\sqrt{3}x$ (M requires <u>values</u> for a and e) $x = -\sqrt{3}$	M1 A1 (4) B1ft (1) 5 Marks
3.	$\rho = \frac{ds}{d\psi} \quad s = \int e^{\sin \psi} \cos \psi d\psi = e^{\sin \psi} (+k)$ <p style="text-align: center;">(M mark may be scored by a full substitution method)</p> $s = 0 \text{ at } \psi = 0: \quad k = -1 \quad s = e^{\sin \psi} - 1$ <p style="text-align: center;">(M mark requires a k, or use of limits)</p>	M1 A1 M1 A1cso (4) 4 Marks

Question number	Scheme Marks	
4.	$\frac{dy}{dx} = \frac{2x}{1+(x^2)^2}$ <p style="text-align: right;">M: $\frac{dy}{dx} = \frac{*}{1+(x^2)^2}$ or $\frac{dy}{dx} = \frac{2x}{1+x^2}$</p> $\frac{d^2y}{dx^2} = \frac{2(1+x^4) - 2x \cdot 4x^3}{(1+x^4)^2} \quad \left(= \frac{2-6x^4}{(1+x^4)^2} \right)$ $\rho = \frac{\left(1 + \left(\frac{dy}{dx} \right)^2 \right)^{\frac{3}{2}}}{\frac{d^2y}{dx^2}} = \frac{(1+1)^{\frac{3}{2}}}{-1}$ $= -2\sqrt{2} \quad (\text{or } 2\sqrt{2}) \quad (\text{or exact equivalent, e.g. } \sqrt{8}, 2^{\frac{3}{2}})$	M1 A1 M1 M1 A1cso A1cso (6) 6 Marks
	<p>2nd M: Quotient or product rule attempt. (Chain rule, if used, must be ‘good’.)</p> <p>3rd M: Attempt ρ with derivative values</p> <p>A: Correct derivative values (1 and -1) seen or implied by working. <u>Alternative:</u> (involving implicit differentiation).</p> $\sec^2 y \frac{dy}{dx} = 2x \quad [\text{M1 A1}] \text{ (allow one ‘slip’ for M1)}$ $\sec^2 y \frac{d^2y}{dx^2} + \frac{dy}{dx} \cdot 2 \sec y (\sec y \tan y) \frac{dy}{dx} = 2$ <p>(or alternative e.g. $\frac{dy}{dx} = 2x \cos^2 y$, so $\frac{d^2y}{dx^2} = 2 \cos^2 y + 2x \cdot 2 \cos y (-\sin y) \frac{dy}{dx}$)</p> <p>Then marks as in main scheme (n.b. $\tan y = 1$, $\sec^2 y = 2$).</p>	[M1]

Question number	Scheme Marks	
5. (a)	$\frac{dy}{dx} = 4 \operatorname{sech}^2 4x - 1$ Put $\frac{dy}{dx} = 0$ $(\cosh^2 4x = 4 \quad \cosh 4x = 2)$ $4x = \ln(2 \pm \sqrt{3})$ or $8x = \ln(7 \pm 4\sqrt{3})$ or $e^{4x} = 2 \pm \sqrt{3}$ or $e^{4x} = 7 \pm 4\sqrt{3}$ $(\pm \text{ or } +)$ $x = \frac{1}{4} \ln(2 + \sqrt{3}) \quad \text{or} \quad x = \frac{1}{8} \ln(7 + 4\sqrt{3}) \quad (\text{or equiv.})$ (b) $y = -\frac{1}{4} \ln(2 + \sqrt{3}) + \tanh(\dots)$ (Substitute for x) $\operatorname{sech} 4x = \frac{1}{2} = \sqrt{1 - \tanh^2 4x}, \quad \tanh 4x = \frac{\sqrt{3}}{2}$ $y = \frac{\sqrt{3}}{2} - \frac{1}{4} \ln(2 + \sqrt{3}) = \frac{1}{4} \left\{ 2\sqrt{3} - \ln(2 + \sqrt{3}) \right\}$ (*)	B1 M1 A1 A1 (4) M1 M1 A1 (3) 7 Marks
	(a) ‘Second solution’, if seen, must be rejected to score the final mark. (b) 2 nd M requires an expression in terms of $\sqrt{3}$ without hyperbolics, exponentials and logarithms.	

Question number	Scheme Marks	
6. (a)	$\frac{dx}{dt} = 1 - \frac{1}{t} \quad \frac{dy}{dt} = 2t^{-\frac{1}{2}}$ $\sqrt{\left(1 - \frac{1}{t}\right)^2 + \left(2t^{-\frac{1}{2}}\right)^2}, \quad = \sqrt{1 + \frac{2}{t} + \frac{1}{t^2}} = 1 + \frac{1}{t} \text{ or } \frac{t+1}{t}$ $\text{Length} = \int_1^4 \left(1 + \frac{1}{t}\right) dt = [t + \ln t]_1^4 = (4 + \ln 4) - 1 = 3 + \ln 4 \quad (*)$	B1 B1 M1, A1
(b)	$\text{Surface area} =$ $2\pi \int_1^4 4\sqrt{t} \sqrt{\left(1 - \frac{1}{t}\right)^2 + \left(2t^{-\frac{1}{2}}\right)^2} dt \quad \left(= 8\pi \int_1^4 \left(t^{\frac{1}{2}} + t^{-\frac{1}{2}}\right) dt\right)$ $= (8\pi) \left[\frac{2t^{\frac{3}{2}}}{3} + 2t^{\frac{1}{2}} \right]_1^4 = (8\pi) \left\{ \left(\frac{16}{3} + 4\right) - \left(\frac{2}{3} + 2\right) \right\} = \frac{160\pi}{3} \quad \left(53\frac{1}{3}\pi\right)$	M1 M1 M1 A1 (4) 11 Marks
	<p>Note dependence of M's.</p> <p>(a) 2nd M: Complete integration attempt. 3rd M: Subs. correct limits and subtract.</p> <p>(b) 2nd M: Complete integration attempt. 3rd M: Subs. correct limits and subtract.</p> <p>In (b), missing 2 or π or 2π throughout could score M0 M1 M1 A0. If 2π is there initially, then lost, could score M1 M1 M1 A0.</p>	

Question number	Scheme Marks	
7.	$\int x^2 \operatorname{arsinh} x dx = \frac{x^3}{3} \operatorname{arsinh} x - \int \frac{x^3}{3\sqrt{x^2+1}} dx$ $\left[\frac{x^3}{3} \operatorname{arsinh} x \right]_0^3 = 9 \operatorname{arsinh} 3 \quad (\text{or } 9 \ln(3 + \sqrt{10}))$ <p>Let $u = x^2 + 1 \quad \frac{du}{dx} = 2x \quad \left[u^2 = x^2 + 1 \quad 2u \frac{du}{dx} = 2x \right]$</p> $\frac{1}{3} \int \frac{x^3}{u^{\frac{1}{2}} \cdot 2x} du = \frac{1}{6} \int \frac{u-1}{u^{\frac{1}{2}}} du = \frac{1}{6} \int \left(u^{\frac{1}{2}} - u^{-\frac{1}{2}} \right) du \quad \left[\frac{1}{3} \int (u^2 - 1) du \right]$ $= \frac{1}{6} \left[\frac{2u^{\frac{3}{2}}}{3} - 2u^{\frac{1}{2}} \right] \quad \left[= \frac{1}{3} \left[\frac{u^3}{3} - u \right] \right]$ <p>When $x = 0, u = 1$ and when $x = 3, u = 10 \quad [.....u = \sqrt{10}]$</p> $\frac{1}{6} \left[\frac{2u^{\frac{3}{2}}}{3} - 2u^{\frac{1}{2}} \right]_1^{10} = \frac{1}{6} \left\{ \left(\frac{20\sqrt{10}}{3} - 2\sqrt{10} \right) - \left(\frac{2}{3} - 2 \right) \right\}$ $\text{Area} = 9 \operatorname{arsinh} 3 - \frac{1}{6} \left(\frac{14\sqrt{10}}{3} + \frac{4}{3} \right) = 9 \ln(3 + \sqrt{10}) - \frac{1}{9} (7\sqrt{10} + 2) \quad (*)$	M1 A1 A1 B1 M1 M1 M1 M1 A1 A1cso (10) 10 Marks
	<u>Dependent M marks:</u> M: Choose an appropriate substitution & find $\frac{du}{dx}$ or 'Set up' integration by parts. M: Get <u>all</u> in terms of ' u ' or Use integration by parts. M: Sound integration. M: Substitute both limits (for the correct variable) and subtract.	

Question number	Scheme Marks	
7.	<p><u>Alternative solution:</u></p> <p>Let $x = \sinh \theta \quad \frac{dx}{d\theta} = \cosh \theta$</p> $\int x^2 \operatorname{arsinh} x dx = \int \theta \sinh^2 \theta \cosh \theta d\theta$ $= \left[\frac{\theta \sinh^3 \theta}{3} \right] - \int \frac{1}{3} \sinh \theta (\cosh^2 \theta - 1) d\theta$ $\left[\frac{\theta \sinh^3 \theta}{3} \right]_0^{\operatorname{arsinh} 3} = 9 \operatorname{arsinh} 3$ $\int \frac{1}{3} \sinh \theta (\cosh^2 \theta - 1) d\theta = \frac{1}{3} \left[\frac{\cosh^3 \theta}{3} - \cosh \theta \right]$ $\frac{1}{3} \left[\frac{\cosh^3 \theta}{3} - \cosh \theta \right]_0^{\operatorname{arsinh} 3} = \frac{1}{3} \left\{ \left(\frac{10\sqrt{10}}{3} - \sqrt{10} \right) - \left(\frac{1}{3} - 1 \right) \right\}$ $\text{Area} = 9 \operatorname{arsinh} 3 - \frac{1}{3} \left(\frac{7\sqrt{10}}{3} + \frac{2}{3} \right) = 9 \ln(3 + \sqrt{10}) - \frac{1}{9} (7\sqrt{10} + 2) \quad (*)$	M1 M1 M1 A1 A1 B1 M1 M1 A1 A1cs (10) 10 Marks

Question number	Scheme Marks	
7. (i)	<p><u>A few alternatives for:</u> $\int \frac{x^3}{\sqrt{x^2 + 1}} dx$.</p> <p>Let $u = x^2 \quad \frac{du}{dx} = 2x$</p> $\int \frac{u^{\frac{3}{2}}}{\sqrt{1+u}} \cdot \frac{1}{2u^{\frac{1}{2}}} du = \frac{1}{2} \int \frac{u}{\sqrt{1+u}} du$ <p>No marks yet... needs another substitution, or parts, or perhaps...</p> $\frac{u}{\sqrt{1+u}} = \sqrt{1+u} - \frac{1}{\sqrt{1+u}}$ $\frac{1}{2} \int \sqrt{1+u} du - \frac{1}{2} \int \frac{1}{\sqrt{1+u}} du$ $\frac{1}{3}(1+u)^{\frac{3}{2}} - (1+u)^{\frac{1}{2}}$ <p>Limits (0 to 9)</p>	M1 M1 M1 M1
(ii)	<p>Let $x = \sinh \theta \quad \frac{dx}{d\theta} = \cosh \theta$</p> $\int \frac{\sinh^3 \theta}{\cosh \theta} \cdot \cosh \theta d\theta = \int \sinh \theta (\cosh^2 \theta - 1) d\theta$ <p>Then, as in the alternative solution,</p> $\int \frac{1}{3} \sinh \theta (\cosh^2 \theta - 1) d\theta = \frac{1}{3} \left[\frac{\cosh^3 \theta}{3} - \cosh \theta \right]$ <p>Limits (0 to $\text{arsinh}3$)</p>	M1 M1 M1 M1

Question number	Scheme Marks	
7. (iii)	<p>Let $u = \tan \theta \quad \frac{du}{d\theta} = \sec^2 \theta$</p> $\int \frac{\tan^3 \theta}{\sec \theta} \cdot \sec^2 \theta d\theta = \int \tan \theta \sec \theta (\sec^2 \theta - 1) d\theta$ $= \int \sec^2 \theta (\sec \theta \tan \theta) d\theta - \int (\sec \theta \tan \theta) d\theta = \frac{\sec^3 \theta}{3} - \sec \theta$ <p>Limits ($\sec \theta = 1$ to $\sec \theta = \sqrt{10}$)</p>	M1 M1 M1 M1
(iv)	<p>(By parts... must be the ‘right way round’, not integrating x^2)</p> $u = x^2, \quad \frac{du}{dx} = 2x \quad \frac{dv}{dx} = \frac{x}{\sqrt{1+x^2}}, \quad v = \sqrt{1+x^2}$ $x^2 \sqrt{1+x^2} - \int 2x \sqrt{1+x^2} dx$ $x^2 \sqrt{1+x^2} - \frac{2}{3} (x^2 + 1)^{\frac{3}{2}}$ <p>Limits</p>	M1 M1 M1 M1 M1 M1
(v)	<p>(By parts)</p> $u = x^3, \quad \frac{du}{dx} = 3x^2 \quad \frac{dv}{dx} = \frac{1}{\sqrt{1+x^2}}, \quad v = \text{arsinh } x$	No progress
(vi)	$\frac{x^3}{\sqrt{1+x^2}} = \frac{x(x^2+1)-x}{\sqrt{1+x^2}} = \frac{x(x^2+1)}{\sqrt{1+x^2}} - \frac{x}{\sqrt{1+x^2}}$ $\int x \sqrt{1+x^2} dx - \int \frac{x}{\sqrt{1+x^2}} dx$ $= \frac{1}{3} (1+x^2)^{\frac{3}{2}} - (1+x^2)^{\frac{1}{2}}$ <p>Limits</p>	M0 M1 M1 M1 M1

Question number	Scheme Marks	
8. (a)	$\int x^n \cosh x dx = x^n \sinh x - \int nx^{n-1} \sinh x dx$ $= x^n \sinh x - nx^{n-1} \cosh x + n(n-1) \int x^{n-2} \cosh x dx$ $I_n = x^n \sinh x - nx^{n-1} \cosh x + n(n-1)I_{n-2} \quad (*)$	M1 A1
(b)	$I_4 = x^4 \sinh x - 4x^3 \cosh x + 12I_2$ $I_4 = x^4 \sinh x - 4x^3 \cosh x + 12(x^2 \sinh x - 2x \cosh x + 2I_0)$ <p>(This M may also be scored by finding I_2 by integration.)</p> $I_0 = \int \cosh x dx = \sinh x + k$ $I_4 = (x^4 + 12x^2 + 24)\sinh x + (-4x^3 - 24x)\cosh x \quad (+C)$	M1 M1 A1 (4)
(c)	$[(x^4 + 12x^2 + 24)\sinh x + (-4x^3 - 24x)\cosh x]_0^1$ $= 37\sinh 1 - 28\cosh 1 \quad \text{M: } x = 1 \text{ substituted throughout (at some stage)}$	B1 A1, A1(5) M1
	$= 37\left(\frac{e - e^{-1}}{2}\right) - 28\left(\frac{e + e^{-1}}{2}\right)$ <p>M: Use of exp. Definitions (can be in terms of x)</p> $= \frac{1}{2}(9e - 65e^{-1})$ <p>(b) Integration constant missing <u>throughout</u> loses the B mark</p>	M1 A1 (3) 12 Marks

•

Question number	Scheme Marks	
9. (a)	$\frac{x^2}{a^2} + \frac{(mx+c)^2}{b^2} = 1$ $b^2x^2 + a^2(mx+c)^2 = a^2b^2$	M1
(b)	$(b^2 + a^2m^2)x^2 + 2a^2mcx + a^2(c^2 - b^2) = 0$ $(2a^2mc)^2 = 4(b^2 + a^2m^2)a^2(c^2 - b^2)$ $4a^4m^2c^2 = 4a^2(b^2c^2 - b^4 + a^2m^2c^2 - a^2m^2b^2)$ $c^2 = b^2 + a^2m^2$	A1 (2) M1 A1 (2)
(c)	Find height and base of triangle (perhaps in terms of c). $OB = c \quad (= \sqrt{b^2 + a^2m^2})$ and $AO = \frac{c}{m} \quad (= \frac{\sqrt{b^2 + a^2m^2}}{m})$ Area of triangle $OAB = \frac{c^2}{2m} = \frac{b^2 + a^2m^2}{2m}$ M: Find area and subs. for c .	M1 A1 M1 A1 (4)
(d)	$\Delta = \frac{b^2 + a^2m^2}{2m} = \frac{b^2}{2}m^{-1} + \frac{a^2}{2}m$ $\frac{d\Delta}{dm} = -\frac{b^2}{2}m^{-2} + \frac{a^2}{2} = 0$ $\frac{b^2}{m^2} = a^2$ $m = \frac{b}{a}$ $\Delta = \left(\frac{b^2}{2}\right)\left(\frac{a}{b}\right) + \left(\frac{a^2}{2}\right)\left(\frac{b}{a}\right) = ab$	A1 (3)
(e)	Root of quadratic: $x = \frac{-a^2mc}{b^2 + a^2m^2}$ (Should be <u>correct</u> if quoted directly) Using $m = \frac{b}{a}$ and $c = \sqrt{b^2 + a^2m^2}$: $x = -\frac{a}{\sqrt{2}}$ (The 2 nd M is dependent on using the quadratic equation).	M1 M1 A1 (3) 14 Marks
(d)	Alternative: $b^2 + a^2m^2 \geq 2bam$ (since $(b-am)^2 \geq 0$) $\frac{b^2 + a^2m^2}{2m} \geq ab$	[M1] [A1]
(e)	Conclusion [A1] <u>Alternative:</u> Begin with full eqn. $(b^2 + a^2m^2)x^2 + 2a^2mcx + a^2(c^2 - b^2) = 0$. In the eqn., use conditions $m = \frac{b}{a}$ and $c = \sqrt{b^2 + a^2m^2}$ ($= b\sqrt{2}$) Simplify and solve eqn., e.g. $2x^2 + 2a\sqrt{2}x + a^2 = 0$ $x = -\frac{a}{\sqrt{2}}$	[M1]