



Mark Scheme (Unused)

January 2022

Pearson Edexcel International A Level
In Statistics S3 (WST03) Paper 01

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

January 2022

Question Paper Log Number P71859A

All the material in this publication is copyright

© Pearson Education Ltd 2022

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

EDEXCEL IAL MATHEMATICS

General Instructions for Marking

1. The total number of marks for the paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:
 - **M** marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
 - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
 - **B** marks are unconditional accuracy marks (independent of M marks)
 - Marks should not be subdivided.
3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod – benefit of doubt
 - ft – follow through
 - the symbol \surd will be used for correct ft
 - cao – correct answer only
 - cso - correct solution only. There must be no errors in this part of the question to obtain this mark
 - isw – ignore subsequent working
 - awrt – answers which round to
 - SC: special case
 - oe – or equivalent (and appropriate)
 - dep – dependent
 - indep – independent
 - dp decimal places
 - sf significant figures
 - * The answer is printed on the paper
 - \square The second mark is dependent on gaining the first mark
4. All A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.
 5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.
 6. Ignore wrong working or incorrect statements following a correct answer.

Question Number	Scheme		Marks
1 (a)	Number the 1200 students (1 – 1200)		B1
	Use a random starting point between 1 and 20		B1
	Select every 20 th person on the list		B1
			(3)
(b)(i)	They only need to generate one random number		B1
			(1)
(b)(ii)	It is not random as the list is ordered alphabetically or not all combinations of sampling units are possible		M1
	e.g. unlikely siblings would be selected		A1
			(2)
(c)	Number of Y9 students = $\frac{200}{1200} \times 60 [= 10]$		M1
	The stratified sample gives a better proportion or is more representative of		A1
			(2)
Notes			Total 8
1 (a)	B1	numbering the students (Allow 0 – 1199).	
	B1	using a random starting point. Must be between 1 and 20 (Allow 0 – 19).	
	B1	selecting every 20 th person.	
(b)(i)	B1	a suitable comment.	
(b)(ii)	M1	a suitable comment.	
	A1	a suitable example.	
(c)	M1	a suitable calculation to find the number of Y9 students e.g. $\frac{200}{1200} \times 60$	
	A1	a correct explanation.	

Question Number	Scheme		Marks
2 (a)	Use of $\bar{x} \pm z \times \frac{1.9}{\sqrt{10}}$; $z = 1.96$		M1;B1
	(52.54..., 54.897...)	awrt 52.5 and 54.9	A1 A1
			(4)
(b)	Use of $1.5 > 2 \times z \times \frac{1.9}{\sqrt{n}}$ oe ; $z = 2.5758$ (or better)		M1;B1
	$1.5 > \frac{9.78804}{\sqrt{n}}$		dM1
	$n > 42.58...$ So $n = 43$		A1
			(4)
Notes			Total 8
2 (a)	M1	for use of correct expression with 1.9, 10 and $1 < z < 3$	
	B1	for $z = 1.96$	
	A1	for awrt 52.5	
	A1	for awrt 54.9	
(b)	M1	use of $z \times \frac{1.9}{\sqrt{n}}$ in a correct inequality with 0.75 or 1.5 and $2 < z < 3$ (allow written as an equation)	
	B1	for $z = 2.5758$ (or better)	
	dM1	dependent on 1 st M1, for solving a correct inequality for the width of the 99% CI (allow an equation rather than an inequality)	
	A1	cao	

Question Number	Scheme											Marks
3 (a)	Driver	<i>A</i>	<i>B</i>	<i>C</i>	<i>D</i>	<i>E</i>	<i>F</i>	<i>G</i>	<i>H</i>	<i>I</i>	<i>J</i>	M1
	Rank FQL	1	5	3	2	6	4	8	9	10	7	
	FP	1	2	3	4	5	6	7	8	9	10	
	$\sum d^2 = 0 + 9 + 0 + 4 + 1 + 4 + 1 + 1 + 1 + 9 [=30]$											M1
	$r_s = 1 - \frac{6(30)}{10(99)}$											dM1
= 0.8181818...											awrt 0.818	A1
												(4)
(b)	$H_0: \rho = 0, H_1: \rho > 0$											B1
	Critical Value $r_s = 0.7455$ or CR: $r_s \dots 0.7455$											B1
	Reject H_0 or significant or lies in the critical region											M1
	There is sufficient evidence of a positive correlation between fastest qualifying lap time and finishing position for these Formula One racing drivers											A1
												(4)
Notes											Total 8	
3 (a)	M1	attempt to rank fastest qualifying lap (at least four correct).										
	M1	finding the difference between each of the ranks and evaluating $\sum d^2$										
	dM1	dependent on 1 st M1. Using $1 - \frac{6 \sum d^2}{10(99)}$ with their $\sum d^2$										
	A1	$\frac{9}{11}$ or awrt 0.818										
(b)	B1	both hypotheses correct. Must be in terms of ρ . Must be attached to H_0 and H_1										
	B1	critical value of 0.7455										
	M1	A correct statement comparing their CV with their r_s - no context needed but do not allow contradicting non contextual comments.										
	A1	correct conclusion which is rejecting H_0 , which must mention lap time and finishing position .										

Question Number	Scheme				Marks	
4	H_0 : There is no association between type of property and the time taken to sell it H_1 : There is an association between type of property and the time taken to sell it				B1	
	Expected	Bungalow	Flat	House	Total	M1 A1
	Within 3 months	10.496	31.488	40.016	(82)	
	More than 3 months	5.504	16.512	20.984	(43)	
	Total	(16)	(48)	(61)	(125)	
	Observed	Expected	$\frac{(O - E)^2}{E}$	$\frac{O^2}{E}$		dM1 A1
	7	10.496	1.1644...	4.6684...		
	29	31.488	0.1965...	26.7085...		
	46	40.016	0.8948...	52.8788...		
	9	5.504	2.2205...	14.7165...		
	19	16.512	0.3748...	21.8628...		
	15	20.984	1.7064...	10.7224...		
	Totals		6.557...	131.557...		
	$[X^2 =] \sum \frac{(O - E)^2}{E}$ or $\sum \frac{O^2}{E} - 125$ = 6.557...				awrt 6.56	dM1 A1
$\nu = (2 - 1)(3 - 1) = 2$					B1	
$c^2_2(0.05) = 5.991 \Rightarrow CR: X^2 \dots 5.991$					B1	
[in the CR/significant/Reject H_0] There is sufficient evidence to suggest that there is an association between type of property and the time taken to sell it.					A1	
				(10)		
Notes					Total 10	
4	B1	Both hypotheses correct. Must mention “type of property” and “time taken” at least once. (may be written in terms of independence)				
	M1	Some attempt at $\frac{(\text{Row Total})(\text{Column Total})}{(\text{Grand Total})}$ Can be implied by at least one correct E_i to 1dp				
	A1	All expected frequencies correct				
	dM1	Dependent on 1 st M1 for at least 2 correct terms for $\frac{(O - E)^2}{E}$ or $\frac{O^2}{E}$ or correct expressions with their E_i Accept 2 sf accuracy.				
	A1	At least 3 correct $\frac{(O - E)^2}{E}$ or $\frac{O^2}{E}$ terms to 2dp or better. Allow truncated answers.				
	dM1	Dependent on 2 nd M1 For applying either $\sum \frac{(O - E)^2}{E}$ or $\sum \frac{O^2}{E} - 125$				
	A1	awrt 6.56				
	B1	$\nu = 2$ This mark can be implied by a correct critical value of 5.991				
	B1	5.991				
	A1	Dependent on the 3 rd M1 and 3 rd B1. A correct contextualised conclusion which is rejecting H_0 Must mention type and time . Contradictory statements score A0. e.g. “significant, do not reject H_0 ”. Condone “relationship” or “connection” here but not “correlation”.				

Question Number	Scheme		Marks
5 (a)(i)	$\left[\bar{x} = \frac{3610}{50} \Rightarrow \right] \bar{x} = 72.2 \quad s_x^2 = \frac{260955.6 - 50(72.2)^2}{50 - 1} = 6.4$		B1; M1 A1
5(a)(ii)	$\left[\bar{y} = \frac{2585}{50} \Rightarrow \right] \bar{y} = 51.7 \quad s_y^2 = \frac{133757.2 - 50(51.7)^2}{50 - 1} = 2.3$		B1 A1
			(5)
(b)	$H_0 : \mu_x - \mu_y = 20$		B1
	$H_1 : \mu_x - \mu_y > 20$		
	$z = \frac{'72.2' - '51.7' - 20}{\sqrt{\frac{'6.4'}{50} + \frac{'2.3'}{50}}}$		M1 M1
	= 1.1986...		awrt 1.20
	One tailed c.v. $Z = 1.6449$ or CR: $Z \dots 1.6449$		B1
	Not in CR/Not significant/Do not reject H_0		M1
	No significant evidence to support Tammy's belief		A1
			(7)
(c)	Since the sample is large the CLT applies.		M1
	No need to assume (the weights) are normally distributed.		A1
			(2)
(d)	Assumed that $s^2 = \sigma^2$		B1
			(1)
Notes			Total 15
5 (a)(i)	B1	$\bar{x} = 72.2$	
	M1	A correct method for finding an unbiased estimate of the variance e.g. $\frac{\sum x^2 - n(\bar{x})^2}{n - 1}$ (May be seen in (i) or (ii))	
	A1	6.4	
5(a)(ii)	B1	$\bar{y} = 51.7$	
	A1	2.3	
(b)	B1	Both hypotheses correct. Allow equivalent hypotheses. Must be in terms of μ	
	M1	For correct standard error. Follow through their values from (a)	
	M1	An attempt at $\frac{a - b - 20}{\sqrt{\frac{c}{50} + \frac{d}{50}}}$ with at least 2 of a, b, c or d correct. Allow \pm	
	A1	awrt 1.20 Allow 1.2 if no incorrect working shown	
	B1	1.6449 or better (seen)	
	M1	A correct statement – need not be contextual but do not allow contradicting non contextual comments.	
	A1	A correct contextual statement. Allow the difference in mean weights is not greater than 20 kg	
(c)	M1	A suitable comment that mentions large and CLT	
	A1	A correct answer, context not required.	
(d)	B1	for the assumption that sample variance = population variance	

Question Number	Scheme				Marks																																								
6 (a)	$\frac{0 \times 1 + 1 \times 10 + 2 \times 23 + 3 \times 15 + 4 \times 19 + 5 \times 9 + 6 \times 3}{80} = 3 *$				B1 (1)																																								
(b)	$r = e^{-3} \times 80 = 3.983 \quad s = \frac{e^{-3} \times 3^5}{5!} \times 80 ; = 8.066$				M1 ; A1																																								
	$t = 80 - (r + 11.949 + 17.923 + 17.923 + 13.443 + s) ; = 6.713$				M1 ; A1 (4)																																								
(c)	<p>H_0 : Poisson (distribution) is a reasonable/suitable/ sensible (model) H_1 : Poisson (distribution) is not a /reasonable/suitable/ sensible (model).</p> <table border="1"> <thead> <tr> <th>Number of emails</th> <th>Combined Observed</th> <th>Combined Expected</th> <th>$\frac{(O - E)^2}{E}$</th> <th>$\frac{O^2}{E}$</th> </tr> </thead> <tbody> <tr> <td>≤ 1</td> <td>11</td> <td>15.932</td> <td>1.5267...</td> <td>7.5947...</td> </tr> <tr> <td>2</td> <td>23</td> <td>17.923</td> <td>1.4381...</td> <td>29.5151...</td> </tr> <tr> <td>3</td> <td>15</td> <td>17.923</td> <td>0.4767...</td> <td>12.5537...</td> </tr> <tr> <td>4</td> <td>19</td> <td>13.443</td> <td>2.2971...</td> <td>26.8541...</td> </tr> <tr> <td>5</td> <td>9</td> <td>8.065</td> <td>0.1083...</td> <td>10.0433...</td> </tr> <tr> <td>≥ 6</td> <td>3</td> <td>6.714</td> <td>2.0544...</td> <td>1.3404...</td> </tr> <tr> <td colspan="2" style="text-align: right;">Totals</td> <td></td> <td>7.901...</td> <td>87.901...</td> </tr> </tbody> </table>				Number of emails	Combined Observed	Combined Expected	$\frac{(O - E)^2}{E}$	$\frac{O^2}{E}$	≤ 1	11	15.932	1.5267...	7.5947...	2	23	17.923	1.4381...	29.5151...	3	15	17.923	0.4767...	12.5537...	4	19	13.443	2.2971...	26.8541...	5	9	8.065	0.1083...	10.0433...	≥ 6	3	6.714	2.0544...	1.3404...	Totals			7.901...	87.901...	B1 M1
Number of emails	Combined Observed	Combined Expected	$\frac{(O - E)^2}{E}$	$\frac{O^2}{E}$																																									
≤ 1	11	15.932	1.5267...	7.5947...																																									
2	23	17.923	1.4381...	29.5151...																																									
3	15	17.923	0.4767...	12.5537...																																									
4	19	13.443	2.2971...	26.8541...																																									
5	9	8.065	0.1083...	10.0433...																																									
≥ 6	3	6.714	2.0544...	1.3404...																																									
Totals			7.901...	87.901...																																									
	$X^2 = \sum \frac{(O - E)^2}{E} \quad \text{or} \quad \sum \frac{O^2}{E} - 80$				M1																																								
	$= 7.901...$ awrt 7.90				A1																																								
	$v = 6 - 1 - 1 = 4$				B1																																								
	$c^2_4(0.10) = 7.779 \Rightarrow \text{CR: } X^2 \dots 7.779$				B1																																								
	[since $X^2 = 7.90$ does lie in CR, then there is sufficient evidence to reject H_0]																																												
	Sufficient evidence to say that Poisson is not a reasonable model				A1																																								
					(7)																																								
Notes					Total 12																																								
6 (a)	B1	For a correct method to shown that the mean is 3																																											
(b)	M1	Use of $\frac{e^{-\lambda} \times \lambda^r}{r!} \times 80$ or May be implied by a correct answer for either r or s																																											
	A1	$r = 3.983$ and $s = 8.066$ (allow $r = 3.984$ and $s = 8.064$ as these come from tables)																																											
	M1	A correct method that ensures that expected totals = 80																																											
	A1	$t = 6.713$ (allow $t = 6.714$ if tables used)																																											
(c)	B1	Both hypotheses correct. Must mention Poisson at least once.																																											
	M1	Combining 0 emails and 1 email. Must have both observed and expected frequencies																																											
	M1	An attempt at the test statistic, at least 2 correct expressions/values (to awrt 2dp)																																											
	A1	awrt 7.90 Accept 7.9 if no incorrect working seen																																											
	B1	$v = 4$ This mark can be implied by a correct critical value of 7.779																																											
	B1	7.779																																											
	A1	A correct conclusion based on their X^2 value and their χ^2 critical value																																											

Question Number	Scheme		Marks
7 (a)	Let X represent $B_1 + B_2 - C_1$		
	$X \sim N(0.268, 0.015633)$ awrt 0.0156		M1 A1
	$P(X < 0) = P\left(Z < \frac{0 - 0.268}{\sqrt{0.015633}} (= -2.14)\right)$		M1
	$(= 1 - 0.9838) = 0.0162$		A1 (4)
(b)	Let Y represent $2.5B_1 + 3C_1 + 3C_2$		
	$Y \sim N(6.918, 0.071478)$ awrt 6.92, 0.0715		M1 A1
	$P(Y > 7) = P\left(Z > \frac{7 - 6.918}{\sqrt{0.071478}} (= 0.31)\right)$		M1
	$(= 1 - 0.6217) = 0.3783$ (Calculator gives 0.3795...) 0.378 – 0.380		A1 (4)
(c)	Mean = $2.94w$		B1
	Standard deviation = $0.084\sqrt{5} w$ (= $0.188w$)		B1
			(2)
(d)	$\frac{6 - 2.94w}{0.084\sqrt{5} w} = -1.2816$		M1;B1
	$-1.2816 \times 0.084\sqrt{5} w + 2.94w = 6$		dM1
	$w = 2.22\dots$ So $w = 2.23$		A1
			(4)
Notes			Total 14
7 (a)	M1	for setting up normal distribution with mean 0.268	
	A1	for a correct expression for variance (= 0.015633) or for standard deviation (= 0.125...)	
(b)	M1	for standardising with 0, 0.268 and their standard deviation	
	A1	awrt 0.0162 (Allow awrt 0.0160 as this comes from a calculator)	
	M1	for setting up normal distribution with mean awrt 6.92	
(c)	A1	for a correct expression for variance (= 0.071478) or for standard deviation (= 0.267...)	
	M1	for standardising with 7, 0.071478 and their standard deviation	
	A1	for answer between 0.378 – 3.80	
(d)	B1	for $2.94w$	
	B1	for $0.084\sqrt{5}w$ or awrt 0.188w	
(d)	M1	for standardising using their mean and their standard deviation = z where $1 < z < 1.5$	
	B1	for -1.28	
	dM1	dependent on M1, for solving their inequality	
	A1	awrt (£)2.23	

