

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise A, Question 1

##### Question:

Write out each of the following as a sum of terms, and hence calculate the sum of the series.

**a**  $\sum_{r=1}^{10} r$

**b**  $\sum_{p=3}^8 p^2$

**c**  $\sum_{r=1}^{10} r^3$

**d**  $\sum_{p=1}^{10} (2p^2 + 3)$

**e**  $\sum_{r=0}^5 (7r + 1)^2$

**f**  $\sum_{i=1}^4 2i(3 - 4i^2)$

##### Solution:

**a**  $1 + 2 + 3 + 4 + 5 + 6 + 7 + 8 + 9 + 10 = 55$

**b**  $3^2 + 4^2 + 5^2 + 6^2 + 7^2 + 8^2 = 199$

**c**  $1^3 + 2^3 + 3^3 + 4^3 + 5^3 + 6^3 + 7^3 + 8^3 + 9^3 + 10^3 = 3025$

{notice that this result is the square of the result for (a)}

**d**  $5 + 11 + 21 + 35 + 53 + 75 + 101 + 131 + 165 + 203 = 800$

**e**  $1 + 64 + 225 + 484 + 841 + 1296 = 2911$

**f**  $-2 - 52 - 198 - 488 = -740$

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise A, Question 2

##### Question:

Write each of the following as a sum of terms, showing the first three terms and the last term.

a  $\sum_{r=1}^n (7r - 1)$

b  $\sum_{r=1}^n (2r^3 + 1)$

c  $\sum_{j=1}^n (j-4)(j+4)$

d  $\sum_{p=3}^k p(p+3)$

##### Solution:

a  $6 + 13 + 20 + \dots + (7n - 1)$

b  $3 + 17 + 55 + \dots + (2n^3 + 1)$

c  $-15 - 12 - 7 + \dots + (n-4)(n+4)$

d  $18 + 28 + 40 + \dots + k(k+3)$

© Pearson Education Ltd 2008

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise A, Question 3

##### Question:

In each part of this question write out, as a sum of terms, the two series defined by  $\sum f(r)$ ; for example, in part c, write out the series  $\sum_{r=1}^{10} r^2$  and  $\sum_{r=1}^{10} r$ . Hence, state whether the given statements relating their sums are true or not.

a  $\sum_{r=1}^n (3r+1) = \sum_{r=2}^{n+1} (3r-2)$

b  $\sum_{r=1}^n 2r = \sum_{r=0}^n 2r$

c  $\sum_{r=1}^{10} r^2 = \left( \sum_{r=1}^{10} r \right)^2$

d  $\sum_{r=1}^4 r^3 = \left( \sum_{r=1}^4 r \right)^2$

e  $\sum_{r=1}^n (3r^2 + 4) = 3 \sum_{r=1}^n r^2 + 4$

##### Solution:

a The two series are exactly the same,  $4 + 7 + 10 + \dots + (3n + 1)$ , and so their sums are the same.

b The two series are exactly the same,  $2 + 4 + 6 + \dots + 2n$ , and so their sums are the same.

c The statement is not true.

$$\sum_{r=1}^{10} r^2 = 1^2 + 2^2 + 3^2 + \dots + 10^2 = 385 \text{ (using your calculator)}$$

$$\left( \sum_{r=1}^{10} r \right)^2 = (1 + 2 + 3 + \dots + 10)^2 = 55^2 = 3025.$$

[This one example is enough to prove  $\sum_{r=1}^n r^2 = \left( \sum_{r=1}^n r \right)^2$  for all  $n$  is not true]

d This statement is true.

$$\sum_{r=1}^4 r^3 = 1^3 + 2^3 + 3^3 + 4^3 = 100$$

$$\left( \sum_{r=1}^4 r \right)^2 = (1 + 2 + 3 + 4)^2 = 10^2 = 100$$

[This does not prove that  $\sum_{r=1}^n r^3 = \left(\sum_{r=1}^n r\right)^2$  for all  $n$ ; but it is true and this will be proved in Chapter 6]

e The statement is not true.

$$\begin{aligned}\sum_{r=1}^n (3r^2 + 4) &= \{3 \times 1^2 + 4\} + \{3 \times 2^2 + 4\} + \{3 \times 3^2 + 4\} + \dots + \{3n^2 + 4\} \\&= 3\{1^2 + 2^2 + 3^2 + \dots + n^2\} + 4n \\3 \sum_{r=1}^n r^2 + 4 &= 3\{1^2 + 2^2 + 3^2 + \dots + n^2\} + 4\end{aligned}$$

© Pearson Education Ltd 2008

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise A, Question 4

##### Question:

Express these series using  $\Sigma$  notation.

**a**  $3 + 4 + 5 + 6 + 7 + 8 + 9 + 10$

**b**  $1 + 8 + 27 + 64 + 125 + 216 + 243 + 512$

**c**  $11 + 21 + 35 + \dots + (2n^2 + 3)$

**d**  $11 + 21 + 35 + \dots + (2n^2 - 4n + 5)$

**e**  $3 \times 5 + 5 \times 7 + 7 \times 9 + \dots + (2r - 1)(2r + 1) + \dots$  to  $k$  terms.

##### Solution:

Answers are not unique (two examples are given, and any letter may be used for  $r$ )

**a**  $\sum_{r=3}^{10} r, \sum_{r=1}^8 (r+2)$

**b**  $\sum_{r=1}^8 r^3, \sum_{r=2}^9 (r-1)^3$

**c**  $\sum_{r=2}^n (2r^2 + 3), \sum_{r=3}^{n+1} (2r^2 - 4r + 5)$

**d**  $\sum_{r=3}^n (2r^2 - 4r + 5), \sum_{r=2}^{n-1} (2r^2 + 3)$

**e**  $\sum_{r=2}^{k+1} (2r - 1)(2r + 1), \sum_{r=1}^k (2r + 1)(2r + 3)$

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise B, Question 1

##### Question:

Use the result for  $\sum_{r=1}^n r$  to calculate

a  $\sum_{r=1}^{36} r$

b  $\sum_{r=1}^{99} r$

c  $\sum_{p=10}^{55} p$

d  $\sum_{r=100}^{200} r$

e  $\sum_{r=1}^k r + \sum_{r=k+1}^{80} r$ , where  $k < 80$ .

##### Solution:

a  $\frac{36 \times 37}{2} = 666$

b  $\frac{99 \times 100}{2} = 4950$

c  $\sum_{p=1}^{55} p - \sum_{p=1}^9 p = \frac{55 \times 56}{2} - \frac{9 \times 10}{2} = 1540 - 45 = 1495$

d  $\sum_{r=1}^{200} r - \sum_{r=1}^{99} r = \frac{200 \times 201}{2} - \frac{99 \times 100}{2} = 20100 - 4950 = 15150$

e  $\sum_{r=1}^k r + \sum_{r=k+1}^{80} r = \sum_{r=1}^{80} r = \frac{80 \times 81}{2} = 3240$

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise B, Question 2

##### Question:

Given that  $\sum_{r=1}^n r = 528$ ,

**a** show that  $n^2 + n - 1056 = 0$

**b** find the value of  $n$ .

##### Solution:

**a**  $\frac{n}{2}(n+1) = 528 \Rightarrow n(n+1) = 1056 \Rightarrow n^2 + n - 1056 = 0$

**b** Factorising:  $(n - 32)(n + 33) = 0$  (or use “the formula”)  $\Rightarrow n = 32$ , as  $n$  cannot be negative.

© Pearson Education Ltd 2008

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise B, Question 3

##### Question:

a Find  $\sum_{k=1}^{2n-1} k$ .

b Hence show that  $\sum_{k=n+1}^{2n-1} k = \frac{3n}{2}(n-1), n \geq 2$ .

##### Solution:

a 
$$\frac{(2n-1)\{(2n-1)+1\}}{2} = \frac{(2n-1)(2n)}{2} = n(2n-1)$$

b

$$\begin{aligned} \sum_{k=1}^{2n-1} k - \sum_{k=1}^n k &= n(2n-1) - \frac{n}{2}(n+1) = \frac{n}{2}\{2(2n-1) - (n+1)\} = \frac{n}{2}(3n-3) \\ &= \frac{3n}{2}(n-1) \end{aligned}$$

© Pearson Education Ltd 2008

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise B, Question 4

##### Question:

Show that  $\sum_{r=k-1}^{2k} r = \frac{(k+2)(3k-1)}{2}, k \geq 1$

##### Solution:

$$\begin{aligned}\sum_{r=1}^{2k} r - \sum_{r=1}^{k-2} r &= \frac{2k}{2}(2k+1) - \frac{(k-2)}{2}(k-1) = \frac{(4k^2+2k)-(k^2-3k+2)}{2} \\ &= \frac{3k^2+5k-2}{2} = \frac{(3k-1)(k+2)}{2}\end{aligned}$$

© Pearson Education Ltd 2008

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise B, Question 5

**Question:**

a Show that  $\sum_{r=1}^{n^2} r - \sum_{r=1}^n r = \frac{n(n^3 - 1)}{2}$ .

b Hence evaluate  $\sum_{r=10}^{81} r$ .

**Solution:**

a  $\frac{n^2(n^2 + 1)}{2} - \frac{n(n + 1)}{2} = \frac{n}{2} \left\{ n(n^2 + 1) - (n + 1) \right\} = \frac{n}{2}(n^3 - 1)$

b  $\sum_{r=10}^{81} r = \sum_{r=1}^{9^2} r - \sum_{r=1}^9 r = \frac{9}{2}(9^3 - 1)$  [using part (a)] = 3276.

© Pearson Education Ltd 2008

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise C, Question 1

##### Question:

(In this exercise use the results for  $\sum_{r=1}^n r$  and  $\sum_{r=1}^n 1$ .)

Calculate the sum of the series:

a  $\sum_{r=1}^{55} (3r - 1)$

b  $\sum_{r=1}^{90} (2 - 7r)$

c  $\sum_{r=1}^{46} (9 + 2r)$

##### Solution:

a  $3 \sum_{r=1}^{55} r - \sum_{r=1}^{55} 1 = 3 \times \frac{55 \times 56}{2} - 55 = 4565$

b  $2 \sum_{r=1}^{90} 1 - 7 \sum_{r=1}^{90} r = 2 \times 90 - 7 \times \frac{90 \times 91}{2} = -28485$

c  $9 \sum_{r=1}^{46} 1 + 2 \sum_{r=1}^{46} r = 9 \times 46 + 2 \times \frac{46 \times 47}{2} = 2576$

© Pearson Education Ltd 2008

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise C, Question 2

##### Question:

Show that

**a**  $\sum_{r=1}^n (3r + 2) = \frac{n}{2}(3n + 7)$

**b**  $\sum_{i=1}^{2n} (5i - 4) = n(10n - 3)$

**c**  $\sum_{r=1}^{n+2} (2r + 3) = (n + 2)(n + 6)$

##### d

$$\sum_{p=3}^n (4p + 5) = (2n + 11)(n - 2)$$

##### Solution:

**a**  $3 \sum_{r=1}^n r + 2 \sum_{r=1}^n 1 = 3 \times \frac{n}{2}(n + 1) + 2n = \frac{n}{2}(3n + 3 + 4) = \frac{n}{2}(3n + 7)$

**b**  $5 \sum_{i=1}^{2n} i - 4 \sum_{i=1}^{2n} 1 = 5 \times \frac{2n}{2}(2n + 1) - 4(2n) = n(10n + 5 - 8) = n(10n - 3)$

**c**  $2 \sum_{r=1}^{n+2} r + 3 \sum_{r=1}^{n+2} 1 = 2 \times \frac{(n+2)}{2}(n+3) + 3(n+2) = (n+2)(n+3+3) = (n+2)(n+6)$

##### d

$$\begin{aligned} \left\{ 4 \sum_{p=1}^n p + 5 \sum_{p=1}^n 1 \right\} - \sum_{p=1}^2 (4p + 5) &= \left\{ 4 \times \frac{n}{2}(n + 1) + 5n \right\} - (9 + 13) \\ &= 2n^2 + 7n - 22 = (2n + 11)(n - 2) \end{aligned}$$

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise C, Question 3

##### Question:

a Show that  $\sum_{r=1}^k (4r - 5) = 2k^2 - 3k.$

b Find the smallest value of  $k$  for which  $\sum_{r=1}^k (4r - 5) > 4850.$

##### Solution:

a  $4 \sum_{r=1}^k r - 5 \sum_{r=1}^k 1 = 4 \times \frac{k}{2}(k+1) - 5k = 2k^2 - 3k$

b  $2k^2 - 3k > 4850 \Rightarrow 2k^2 - 3k - 4850 > 0 \Rightarrow (2k + 97)(k - 50) > 0,$

so  $k > 50$  [ $k$  is positive]  $\Rightarrow k = 51$

© Pearson Education Ltd 2008

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise C, Question 4

##### Question:

Given that  $u_r = ar + b$  and  $\sum_{r=1}^n u_r = \frac{n}{2}(7n + 1)$ , find the constants  $a$  and  $b$ .

##### Solution:

$$\sum_{r=1}^n (ar + b) = \frac{an}{2}(n + 1) + bn = \frac{an^2 + (a + 2b)n}{2}$$

$$\text{Comparing with } \frac{7n^2 + n}{2} \Rightarrow a = 7 \text{ and } a + 2b = 1$$

$$\text{So } a = 7, b = -3$$

© Pearson Education Ltd 2008

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise C, Question 5

##### Question:

a Show that  $\sum_{r=1}^{4n-1} (1+3r) = 24n^2 - 2n - 1$   $n \geq 1$ .

b Hence calculate  $\sum_{r=1}^{99} (1+3r)$ .

##### Solution:

a  $\sum_{r=1}^{4n-1} 1 + 3 \sum_{r=1}^{4n-1} r = (4n-1) + 3 \times \frac{(4n-1)(4n)}{2} = (4n-1)(1+6n) = 24n^2 - 2n - 1$

b Substituting  $n = 25$  into above result gives 14949

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise C, Question 6

##### Question:

Show that  $\sum_{r=1}^{2k+1} (4 - 5r) = -(2k + 1)(5k + 1)$ ,  $k \geq 0$

##### Solution:

$$\begin{aligned} 4 \sum_{r=1}^{2k+1} 1 - 5 \sum_{r=1}^{2k+1} r &= 4(2k+1) - 5 \frac{(2k+1)}{2}(2k+2) = (2k+1)\{4 - 5(k+1)\} \\ &= (2k+1)(-1 - 5k) = -(2k+1)(5k+1) \end{aligned}$$

© Pearson Education Ltd 2008

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise D, Question 1

##### Question:

Verify that  $\sum_{r=1}^n r^2 = \frac{n}{6}(n+1)(2n+1)$  is true for  $n = 1, 2$  and  $3$ .

##### Solution:

$$\text{For } n = 1, \quad \sum_{r=1}^n r^2 = 1^2 = 1, \quad \frac{n}{6}(n+1)(2n+1) = \frac{1}{6}(1+1)(2+1) = 1$$

$$\text{For } n = 2, \quad \sum_{r=1}^n r^2 = 1^2 + 2^2 = 5, \quad \frac{n}{6}(n+1)(2n+1) = \frac{2}{6}(2+1)(4+1) = 5$$

$$\text{For } n = 3, \quad \sum_{r=1}^n r^2 = 1^2 + 2^2 + 3^2 = 14, \quad \frac{n}{6}(n+1)(2n+1) = \frac{3}{6}(3+1)(6+1) = 14$$

© Pearson Education Ltd 2008

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise D, Question 2

##### Question:

a By writing out each series, evaluate  $\sum_{r=1}^n r$  for  $n = 1, 2, 3$  and  $4$ .

b By writing out each series, evaluate  $\sum_{r=1}^n r^3$  for  $n = 1, 2, 3$  and  $4$ .

c What do you notice about the corresponding results for each value of  $n$ ?

##### Solution:

a  $\sum_{r=1}^1 r = 1; \quad \sum_{r=1}^2 r = 1 + 2 = 3; \quad \sum_{r=1}^3 r = 1 + 2 + 3 = 6; \quad \sum_{r=1}^4 r = 1 + 2 + 3 + 4 = 10$

b  $\sum_{r=1}^1 r^3 = 1; \quad \sum_{r=1}^2 r^3 = 1^3 + 2^3 = 9; \quad \sum_{r=1}^3 r^3 = 1^3 + 2^3 + 3^3 = 36; \quad \sum_{r=1}^4 r^3 = 1^3 + 2^3 + 3^3 + 4^3 = 100$

c The results for (b) are the square of the results for (a)

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

**Series****Exercise D, Question 3****Question:**

Using the appropriate formula, evaluate

a  $\sum_{r=1}^{100} r^2$

b  $\sum_{r=20}^{40} r^2$

c  $\sum_{r=1}^{30} r^3$

d  $\sum_{r=25}^{45} r^3$

**Solution:**

a  $\frac{100}{6} \times 101 \times 201 = 338350$

b  $\sum_{r=1}^{40} r^2 - \sum_{r=1}^{19} r^2 = \frac{40}{6} \times 41 \times 81 - \frac{19}{6} \times 20 \times 39 = 22140 - 2470 = 19670$

c  $\frac{30^2 \times 31^2}{4} = 216225$

d  $\sum_{r=1}^{45} r^3 - \sum_{r=1}^{24} r^3 = \frac{45^2 \times 46^2}{4} - \frac{24^2 \times 25^2}{4} = 1071225 - 90000 = 981225$

© Pearson Education Ltd 2008

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise D, Question 4

##### Question:

Use the formula for  $\sum_{r=1}^n r^2$  or  $\sum_{r=1}^n r^3$  to find the sum of

**a**  $1^2 + 2^2 + 3^2 + 4^2 + \dots + 52^2$

**b**  $2^3 + 3^3 + 4^3 + \dots + 40^3$

**c**  $26^2 + 27^2 + 28^2 + 29^2 + \dots + 100^2$

**d**  $1^2 + 2^2 + 3^2 + \dots + (k+1)^2$

**e**  $1^3 + 2^3 + 3^3 + \dots + (2n-1)^3$

##### Solution:

**a**  $\sum_{r=1}^{52} r^2 = \frac{52}{6} \times 53 \times 105 = 48230$

**b**  $\sum_{r=1}^{40} r^3 - 1 = \frac{40^2 \times 41^2}{4} - 1 = 672399$

**c**  $\sum_{r=1}^{100} r^2 - \sum_{r=1}^{25} r^2 = \frac{100}{6} \times 101 \times 201 - \frac{25}{6} \times 26 \times 51 = 338350 - 5525 = 332825$

**d**  $\sum_{r=1}^{k+1} r^2 = \frac{(k+1)}{6}(k+2)(2k+3)$

**e**  $\sum_{r=1}^{2n-1} r^3 = \frac{(2n-1)^2(2n)^2}{4} = n^2(2n-1)^2$

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise D, Question 5

##### Question:

For each of the following series write down, in terms of  $n$ , the sum, giving the result in its simplest form

**a**  $\sum_{r=1}^{2n} r^2$

**b**  $\sum_{r=1}^{n^2-1} r^2$

**c**  $\sum_{i=1}^{2n-1} i^2$

**d**  $\sum_{r=1}^{n+1} r^3$

**e**  $\sum_{k=n+1}^{3n} k^3, n > 0.$

##### Solution:

**a**  $\frac{(2n)}{6}(2n+1)(4n+1) = \frac{n}{3}(2n+1)(4n+1)$

**b**  $\frac{(n^2-1)n^2(2n^2-1)}{6}$

**c**  $\frac{(2n-1)}{6}(2n)[2(2n-1)+1] = \frac{(2n-1)}{6}(2n)(4n-1) = \frac{n}{3}(2n-1)(4n-1)$

**d**  $\frac{(n+1)^2(n+2)^2}{4}$

**e**

$$\begin{aligned} \sum_{r=1}^{3n} k^3 - \sum_{r=1}^n k^3 &= \frac{(3n)^2(3n+1)^2}{4} - \frac{n^2(n+1)^2}{4} = \frac{n^2}{4}\{9(3n+1)^2 - (n+1)^2\} \\ &= \frac{n^2}{4}\{3(3n+1) - (n+1)\}\{3(3n+1) + (n+1)\} [\text{using } a^2 - b^2 = (a-b)(a+b)] \\ &= \frac{n^2}{4}\{(8n+2)(10n+4)\} \\ &= n^2(4n+1)(5n+2) \end{aligned}$$

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise D, Question 6

**Question:**

Show that

$$\mathbf{a} \sum_{r=2}^n r^2 = \frac{1}{6}(n-1)(2n^2 + 5n + 6)$$

$$\mathbf{b} \sum_{r=n}^{2n} r^2 = \frac{n}{6}(n+1)(14n+1)$$

**Solution:**

$$\mathbf{a} \frac{n}{6}(n+1)(2n+1) - 1 = \frac{2n^3 + 3n^2 + n - 6}{6} = \frac{(n-1)(2n^2 + 5n + 6)}{6} \quad [\text{use factor theorem}]$$

**b**

$$\begin{aligned} \sum_{r=1}^{2n} r^2 - \sum_{r=1}^{n-1} r^2 &= \frac{2n}{6}(2n+1)(4n+1) - \frac{(n-1)}{6}n(2n-1) \\ &= \frac{n}{6}\{2(2n+1)(4n+1) - (n-1)(2n-1)\} \\ &= \frac{n}{6}\{(16n^2 + 12n + 2) - (2n^2 - 3n + 1)\} = \frac{n}{6}(14n^2 + 15n + 1) \\ &= \frac{n}{6}(14n+1)(n+1) \end{aligned}$$

© Pearson Education Ltd 2008

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise D, Question 7

**Question:**

a Show that  $\sum_{k=n}^{2n} k^3 = \frac{3n^2(n+1)(5n+1)}{4}$

b Find  $\sum_{k=30}^{60} k^3$ .

**Solution:**

a

$$\begin{aligned}\sum_{k=1}^{2n} k^3 - \sum_{k=1}^{n-1} k^3 &= \frac{(2n)^2(2n+1)^2}{4} - \frac{(n-1)^2n^2}{4} \\&= \frac{n^2}{4} \{4(2n+1)^2 - (n-1)^2\} \\&= \frac{n^2}{4} [\{2(2n+1) + (n-1)\} \{2(2n+1) - (n-1)\}] \text{ "Difference of two squares"} \\&= \frac{n^2}{4} (5n+1)(3n+3) = \frac{3n^2}{4} (5n+1)(n+1)\end{aligned}$$

b Substituting  $n = 30$  into (a) gives 3 159 675

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise D, Question 8

**Question:**

a Show that  $\sum_{r=1}^{2n} r^3 = n^2(2n+1)^2$ .

b By writing out the series for  $\sum_{r=1}^n (2r)^3$ , show that  $\sum_{r=1}^n (2r)^3 = 8 \sum_{r=1}^n r^3$ .

c Show that  $1^3 + 3^3 + 5^3 + \dots + (2n-1)^3$  can be written as  $\sum_{r=1}^{2n} r^3 - \sum_{r=1}^n (2r)^3$ .

d Hence show that the sum of the cubes of the first  $n$  odd natural numbers,  $1^3 + 3^3 + 5^3 + \dots + (2n-1)^3$ , is  $n^2(2n^2-1)$ .

**Solution:**

a  $\sum_{r=1}^{2n} r^3 = \frac{(2n)^2(2n+1)^2}{4} = n^2(2n+1)^2$ .

b  $\sum_{r=1}^n (2r)^3 = 2^3 + 4^3 + 6^3 + \dots + (2n)^3 = 2^3 \{1^3 + 2^3 + 3^3 + \dots + n^3\} = 8 \sum_{r=1}^n r^3$ .

c

$$\begin{aligned} 1^3 + 3^3 + 5^3 + \dots + (2n-1)^3 &= \{1^3 + 2^3 + 3^3 + \dots + (2n-1)^3 + (2n)^3\} - \{2^3 + 4^3 + 6^3 + \dots + (2n)^3\} \\ &= \sum_{r=1}^{2n} r^3 - \sum_{r=1}^n (2r)^3. \end{aligned}$$

d Using the results in parts (b) and (c),  $1^3 + 3^3 + 5^3 + \dots + (2n-1)^3 = \sum_{r=1}^{2n} r^3 - 8 \sum_{r=1}^n r^3$

$$\begin{aligned} &= n^2(2n+1)^2 - 8 \sum_{r=1}^n r^3 \text{ (using(a))} \\ &= n^2(2n+1)^2 - \frac{8n^2(n+1)^2}{4} \\ &= n^2[(2n+1)^2 - 2(n+1)^2] \\ &= n^2[(4n^2 + 4n + 1) - 2(n^2 + 2n + 1)] \\ &= n^2(2n^2 - 1) \end{aligned}$$

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise E, Question 1

##### Question:

Use the formulae for  $\sum_{r=1}^n r^3$ ,  $\sum_{r=1}^n r^2$ ,  $\sum_{r=1}^n r$  and  $\sum_{r=1}^n 1$ , where appropriate, to find

a  $\sum_{m=1}^{30} (m^2 - 1)$

b  $\sum_{r=1}^{40} r(r+4)$

c  $\sum_{r=1}^{80} r(r^2 + 3)$

d  $\sum_{r=11}^{35} (r^3 - 2)$ .

##### Solution:

a  $\sum_{m=1}^{30} m^2 - 30 = \frac{30 \times 31 \times 61}{6} - 30 = 9425$

b  $\sum_{r=1}^{40} r^2 + 4 \sum_{r=1}^{40} r = \frac{40 \times 41 \times 81}{6} + 4 \times \frac{40 \times 41}{2} = 22140 + 3280 = 25420$

c  $\sum_{r=1}^{80} r^3 + 3 \sum_{r=1}^{80} r = \frac{80^2 \times 81^2}{4} + 3 \times \frac{80 \times 81}{2} = 10497600 + 9720 = 10507\ 320$

d  $\sum_{r=1}^{35} (r^3 - 2) - \sum_{r=1}^{10} (r^3 - 2) = \sum_{r=1}^{35} r^3 - 2(35) - \left[ \sum_{r=1}^{10} r^3 - 2(10) \right]$

$$\sum_{r=1}^{35} r^3 - \sum_{r=1}^{10} r^3 - 2(35 - 10) = \frac{35^2 \times 36^2}{4} - \frac{10^2 \times 11^2}{4} - 50 = 396900 - 3025 - 50 = 393825.$$

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise E, Question 2

**Question:**

Use the formulae for  $\sum_{r=1}^n r^3$ ,  $\sum_{r=1}^n r^2$ , and  $\sum_{r=1}^n r$ , where appropriate, to find

a  $\sum_{r=1}^n (r^2 + 4r)$

b  $\sum_{r=1}^n r(2r^2 - 1)$

c  $\sum_{r=1}^{2n} r^2(1+r)$ , giving your answer in its simplest form.

**Solution:**

$$\mathbf{a} \sum_{r=1}^n r^2 + 4 \sum_{r=1}^n r = \frac{n(n+1)(2n+1)}{6} + \frac{4n(n+1)}{2} = \frac{n(n+1)\{(2n+1)+12\}}{6} = \frac{n}{6}(n+1)(2n+13)$$

$$\mathbf{b} 2 \sum_{r=1}^n r^3 - \sum_{r=1}^n r = \frac{2n^2(n+1)^2}{4} - \frac{n(n+1)}{2} = \frac{n(n+1)\{n(n+1)-1\}}{2} = \frac{n}{2}(n+1)(n^2+n-1)$$

**c**

$$\begin{aligned} \sum_{r=1}^{2n} r^2 + \sum_{r=1}^{2n} r^3 &= \frac{2n(2n+1)(4n+1)}{6} + \frac{(2n)^2(2n+1)^2}{4} = \frac{n(2n+1)\{(4n+1)+3n(2n+1)\}}{3} \\ &= \frac{n}{3}(2n+1)(6n^2+7n+1) = \frac{n}{3}(n+1)(2n+1)(6n+1) \end{aligned}$$

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise E, Question 3

##### Question:

a Write out  $\sum_{r=1}^n r(r+1)$  as a sum, showing at least the first three terms and the final term.

b Use the results for  $\sum_{r=1}^n r$  and  $\sum_{r=1}^n r^2$  to calculate

$$1 \times 2 + 2 \times 3 + 3 \times 4 + 4 \times 5 + 5 \times 6 + \dots + 60 \times 61.$$

##### Solution:

a  $1 \times 2 + 2 \times 3 + 3 \times 4 + \dots + n(n+1)$

b Putting  $n = 60$ :  $\sum_{r=1}^{60} r^2 + \sum_{r=1}^{60} r = \frac{60 \times 61 \times 121}{6} + \frac{60 \times 61}{2} = 73810 + 1830 = 75640$

© Pearson Education Ltd 2008

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise E, Question 4

##### Question:

a Show that  $\sum_{r=1}^n (r+2)(r+5) = \frac{n}{3}(n^2 + 12n + 41)$ .

b Hence calculate  $\sum_{r=10}^{50} (r+2)(r+5)$ .

##### Solution:

a

$$\begin{aligned}\sum_{r=1}^n (r^2 + 7r + 10) &= \sum_{r=1}^n r^2 + 7 \sum_{r=1}^n r + 10 \sum_{r=1}^n 1 \\&= \frac{n}{6}(n+1)(2n+1) + 7 \frac{n}{2}(n+1) + 10n \\&= \frac{n}{6}\{(2n^2 + 3n + 1) + 21(n+1) + 60\} \\&= \frac{n}{6}(2n^2 + 24n + 82) = \frac{n}{3}(n^2 + 12n + 41)\end{aligned}$$

b Substituting  $n = 50$  and  $n = 9$  in the formula in (a), and subtracting, gives 51 660.

© Pearson Education Ltd 2008

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise E, Question 5

**Question:**

a Show that  $\sum_{r=2}^n (r-1)r(r+1) = \frac{(n-1)n(n+1)(n+2)}{4}$ .

b Hence find the sum of the series  $13 \times 14 \times 15 + 14 \times 15 \times 16 + 15 \times 16 \times 17 + \dots + 44 \times 45 \times 46$ .

**Solution:**

a

$$\begin{aligned}\sum_{r=2}^n (r^3 - r) &= \sum_{r=1}^n (r^3 - r) = \sum_{r=1}^n r^3 - \sum_{r=1}^n r = \frac{n^2(n+1)^2}{4} - \frac{n(n+1)}{2} \\ &= \frac{n(n+1)}{4}(n^2 + n - 2) \\ &= \frac{n}{4}(n+1)\{n^2 + n - 2\} \\ &= \frac{n}{4}(n+1)(n+2)(n-1) = \frac{(n-1)n(n+1)(n+2)}{4}\end{aligned}$$

$$\begin{aligned}\mathbf{b} \quad \sum_{r=14}^{45} (r-1)r(r+1) &= \sum_{r=2}^{45} (r-1)r(r+1) - \sum_{r=2}^{13} (r-1)r(r+1) = \frac{44 \times 45 \times 46 \times 47}{4} - \frac{12 \times 13 \times 14 \times 15}{4} \\ &= 1\,062\,000\end{aligned}$$

© Pearson Education Ltd 2008

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise E, Question 6

##### Question:

Find the following sums, and check your results for the cases  $n = 1, 2$  and  $3$ .

**a**  $\sum_{r=1}^n (r^3 - 1)$

**b**  $\sum_{r=1}^n (2r - 1)^2$

**c**  $\sum_{r=1}^n r(r+1)^2$

##### Solution:

$$\mathbf{a} \sum_{r=1}^n r^3 - \sum_{r=1}^n 1 = \frac{n^2(n+1)^2}{4} - n = \frac{n}{4}\{n(n+1)^2 - 4\} = \frac{n}{4}(n^3 + 2n^2 + n - 4)$$

$$\text{When } n = 1 : \sum_{r=1}^1 (r^3 - 1) = 0; \quad \frac{n}{4}(n^3 + 2n^2 + n - 4) = \frac{1 \times 0}{4} = 0$$

$$\text{When } n = 2 : \sum_{r=1}^2 (r^3 - 1) = 0 + 7 = 7; \quad \frac{n}{4}(n^3 + 2n^2 + n - 4) = \frac{2 \times 14}{4} = 7$$

$$\text{When } n = 3 : \sum_{r=1}^3 (r^3 - 1) = 0 + 7 + 26 = 33; \quad \frac{n}{4}(n^3 + 2n^2 + n - 4) = \frac{3 \times 44}{4} = 33$$

##### b

$$\begin{aligned} \sum_{r=1}^n (4r^2 - 4r + 1) &= 4 \sum_{r=1}^n r^2 - 4 \sum_{r=1}^n r + \sum_{r=1}^n 1 = \frac{4n(n+1)(2n+1)}{6} - \frac{4n(n+1)}{2} + n \\ &= \frac{n}{3} \left\{ 2(2n^2 + 3n + 1) - 6(n+1) + 3 \right\} = \frac{n}{3}(4n^2 - 1) \end{aligned}$$

$$\text{When } n = 1 : \sum_{r=1}^1 (4r^2 - 4r + 1) = 1; \quad \frac{n}{3}(4n^2 - 1) = \frac{1 \times 3}{3} = 1$$

$$\text{When } n = 2 : \sum_{r=1}^2 (4r^2 - 4r + 1) = 1 + 9 = 10; \quad \frac{n}{3}(4n^2 - 1) = \frac{2 \times 15}{3} = 10$$

$$\text{When } n = 3 : \sum_{r=1}^3 (4r^2 - 4r + 1) = 1 + 9 + 25 = 35; \quad \frac{n}{3}(4n^2 - 1) = \frac{3 \times 35}{3} = 35$$

##### c

$$\begin{aligned}
 \sum_{r=1}^n (r^3 + 2r^2 + r) &= \sum_{r=1}^n r^3 + 2 \sum_{r=1}^n r^2 + \sum_{r=1}^n r = \frac{n^2(n+1)^2}{4} + \frac{2n(n+1)(2n+1)}{6} + \frac{n(n+1)}{2} \\
 &= \frac{n(n+1)}{12} \{3n(n+1) + 4(2n+1) + 6\} = \frac{n(n+1)}{12} \{3n^2 + 11n + 10\} \\
 &= \frac{n}{12}(n+1)(n+2)(3n+5)
 \end{aligned}$$

When  $n = 1$  :  $\sum_{r=1}^1 r(r+1)^2 = 1 \times 4 = 4$ ;  $\frac{n}{12}(n+1)(n+2)(3n+5) = \frac{1 \times 2 \times 3 \times 8}{12} = 4$

When  $n = 2$  :  $\sum_{r=1}^2 r(r+1)^2 = 4 + 2 \times 9 = 22$ ;  $\frac{n}{12}(n+1)(n+2)(3n+5) = \frac{2 \times 3 \times 4 \times 11}{12} = 22$

When  $n = 3$  :  $\sum_{r=1}^3 r(r+1)^2 = 22 + 3 \times 16 = 70$ ;  $\frac{n}{12}(n+1)(n+2)(3n+5) = \frac{3 \times 4 \times 5 \times 14}{12} = 70$

© Pearson Education Ltd 2008

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise E, Question 7

**Question:**

a Show that  $\sum_{r=1}^n r^2(r-1) = \frac{n}{12}(n^2-1)(3n+2)$ .

b Deduce the sum of  $1 \times 2^2 + 2 \times 3^2 + 3 \times 4^2 + \dots + 30 \times 31^2$ .

**Solution:**

a

$$\begin{aligned}\sum_{r=1}^n r^3 - \sum_{r=1}^n r^2 &= \frac{n^2(n+1)^2}{4} - \frac{n(n+1)(2n+1)}{6} \\&= \frac{n(n+1)}{12} \{3n(n+1) - 2(2n+1)\} \\&= \frac{n(n+1)}{12} (3n^2 - n - 2) \\&= \frac{n(n+1)(n-1)(3n+2)}{12} = \frac{n(n^2-1)(3n+2)}{12}\end{aligned}$$

b As  $\sum_{r=2}^{31} r^2(r-1) = \sum_{r=1}^{31} r^2(r-1)$ , substitute  $n = 31$  in (a); sum = 235 600

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise E, Question 8

**Question:**

a Show that  $\sum_{r=2}^n (r-1)(r+1) = \frac{n}{6}(2n+5)(n-1)$ .

b Hence sum the series  $1 \times 3 + 2 \times 4 + 3 \times 5 + \dots + 35 \times 37$ .

**Solution:**

a [  $\sum_{r=2}^n (r^2 - 1) = \sum_{r=1}^n (r^2 - 1)$  as when  $r = 1$  the term is zero ]

$$\begin{aligned}\sum_{r=1}^n (r^2 - 1) &= \sum_{r=1}^n r^2 - \sum_{r=1}^n 1 = \frac{n}{6}(n+1)(2n+1) - n \\ &= \frac{n}{6}\{(2n^2 + 3n + 1) - 6\} \\ &= \frac{n}{6}(2n^2 + 3n - 5) \\ &= \frac{n}{6}(2n+5)(n-1)\end{aligned}$$

b  $1 \times 3 + 2 \times 4 + 3 \times 5 + \dots + 35 \times 37 = \sum_{r=1}^{36} (r-1)(r+1)$

Substituting  $n = 36$  into result in (a) gives 16 170

© Pearson Education Ltd 2008

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise E, Question 9

**Question:**

a Write out the series defined by  $\sum_{r=7}^{12} r(2 + 3r)$ , and hence find its sum.

b Show that  $\sum_{r=n+1}^{2n} r(2 + 3r) = \frac{n}{2}(14n^2 + 15n + 3)$ .

c By substituting the appropriate value of  $n$  into the formula in b, check that your answer for a is correct.

**Solution:**

a  $7 \times 23 + 8 \times 26 + 9 \times 29 + 10 \times 32 + 11 \times 35 + 12 \times 38 = 1791$ .

b 
$$\sum_{r=n+1}^{2n} (2r + 3r^2) = \sum_{r=1}^{2n} (2r + 3r^2) - \sum_{r=1}^n (2r + 3r^2)$$

$$\begin{aligned} \sum_{r=1}^n (2r + 3r^2) &= 2 \sum_{r=1}^n r + 3 \sum_{r=1}^n r^2 = n(n+1) + \frac{n}{2}(n+1)(2n+1) \\ &= \frac{n}{2}(n+1)\{2 + (2n+1)\} \\ &= \frac{n}{2}(n+1)(2n+3) \end{aligned}$$

$$\Rightarrow \sum_{r=1}^{2n} (2r + 3r^2) = n(2n+1)(4n+3)$$

$$\begin{aligned} \sum_{r=n+1}^{2n} (2r + 3r^2) &= n(2n+1)(4n+3) - \frac{n}{2}(n+1)(2n+3) \\ &= \frac{n}{2}\{2(2n+1)(4n+3) - (n+1)(2n+3)\} \\ &= \frac{n}{2}\{(16n^2 + 20n + 6) - (2n^2 + 5n + 3)\} \\ &= \frac{n}{2}(14n^2 + 15n + 3) \end{aligned}$$

c Substituting  $n = 6$  gives 1791

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise E, Question 10

##### Question:

Find the sum of the series  $1 \times 1 + 2 \times 3 + 3 \times 5 + \dots$  to  $n$  terms.

##### Solution:

Series can be written as  $\sum_{r=1}^n r(2r - 1)$

$$\begin{aligned}\sum_{r=1}^n r(2r - 1) &= 2 \sum_{r=1}^n r^2 - \sum_{r=1}^n r = 2 \times \frac{n}{6}(n+1)(2n+1) - \frac{n}{2}(n+1) \\ &= \frac{n(n+1)\{2(2n+1)-3\}}{6} \\ &= \frac{n(n+1)(4n-1)}{6}\end{aligned}$$

© Pearson Education Ltd 2008

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise F, Question 1

##### Question:

a Write down the first three terms and the last term of the series given by  $\sum_{r=1}^n (2r + 3^r)$ .

b Find the sum of this series.

c Verify that your result in b is correct for the cases  $n = 1, 2$  and  $3$ .

##### Solution:

a  $(2+3) + (4+3^2) + (6+3^3) + \dots + (2n+3^n) \quad [= 5 + 13 + 33 + \dots + (2n+3^n)]$

b  $\sum_{r=1}^n (2r + 3^r) = 2 \sum_{r=1}^n r + \sum_{r=1}^n 3^r = n(n+1) + \frac{3}{2}(3^n - 1) \quad [\text{AP} + \text{GP}]$

c

$n = 1$ : (b) gives  $2 + 3 = 5$ , agrees with (a)

$n = 2$ : (b) gives  $6 + 12 = 18$ , agrees with (a)

$n = 3$ : (b) gives  $12 + 39 = 51$ , agrees with (a)

© Pearson Education Ltd 2008

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise F, Question 2

##### Question:

Find

$$\mathbf{a} \sum_{r=1}^{50} (7r + 5)$$

$$\mathbf{b} \sum_{r=1}^{40} (2r^2 - 1)$$

$$\mathbf{c} \sum_{r=33}^{75} r^3.$$

##### Solution:

$$\mathbf{a} 7 \sum_{r=1}^{50} r + 5 \sum_{r=1}^{50} 1 = \frac{7 \times 50 \times 51}{2} + 5(50) = 9175$$

$$\mathbf{b} 2 \sum_{r=1}^{40} r^2 - \sum_{r=1}^{40} 1 = \frac{40(41)(81)}{3} - 40 = 44\,240$$

$$\mathbf{c} \sum_{r=1}^{75} r^3 - \sum_{r=1}^{32} r^3 = \frac{75^2 \times 76^2}{4} - \frac{32^2 \times 33^2}{4} = 7\,843\,716$$

© Pearson Education Ltd 2008

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise F, Question 3

##### Question:

Given that  $\sum_{r=1}^n U_r = n^2 + 4n$ ,

a find  $\sum_{r=1}^{n-1} U_r$ .

b Deduce an expression for  $U_n$ .

c Find  $\sum_{r=n}^{2n} U_r$ .

##### Solution:

a Replacing  $n$  with  $(n - 1)$  gives  $(n - 1)^2 + 4(n - 1) = n^2 + 2n - 3$

b  $U_n = \sum_{r=1}^n U_r - \sum_{r=1}^{n-1} U_r = n^2 + 4n - (n^2 + 2n - 3) = 2n + 3$

c  $\sum_{r=1}^{2n} U_r - \sum_{r=1}^{n-1} U_r = (4n^2 + 8n) - (n^2 + 2n - 3) = 3n^2 + 6n + 3 = 3(n + 1)^2$

© Pearson Education Ltd 2008

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

**Series****Exercise F, Question 4****Question:**

Evaluate  $\sum_{r=1}^{30} r(3r - 1)$

**Solution:**

$$3 \sum_{r=1}^{30} r^2 - \sum_{r=1}^{30} r = \frac{3 \times 30 \times 31 \times 61}{6} - \frac{30 \times 31}{2} = 28365 - 465 = 27900$$

© Pearson Education Ltd 2008

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

**Series****Exercise F, Question 5****Question:**

Find  $\sum_{r=1}^n r^2(r-3)$ .

**Solution:**

$$\begin{aligned}\sum_{r=1}^n r^3 - 3 \sum_{r=1}^n r^2 &= \frac{n^2}{4}(n+1)^2 - \frac{n}{2}(n+1)(2n+1) \\&= \frac{n}{4}(n+1)\{n(n+1) - 2(2n+1)\} \\&= \frac{n}{4}(n+1)(n^2 - 3n - 2)\end{aligned}$$

© Pearson Education Ltd 2008

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise F, Question 6

##### Question:

Show that  $\sum_{r=1}^{2n} (2r-1)^2 = \frac{2n}{3}(16n^2 - 1)$ .

##### Solution:

$$\begin{aligned} 4 \sum_{r=1}^{2n} r^2 - 4 \sum_{r=1}^{2n} r + \sum_{r=1}^{2n} 1 &= \frac{4}{3}n(2n+1)(4n+1) - 4n(2n+1) + 2n \\ &= \frac{n}{3}\{4(2n+1)(4n+1) - 12(2n+1) + 6\} \\ &= \frac{n}{3}\{32n^2 + 24n + 4 - 12(2n+1) + 6\} \\ &= \frac{n}{3}(32n^2 - 2) \\ &= \frac{2n}{3}(16n^2 - 1) \end{aligned}$$

© Pearson Education Ltd 2008

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise F, Question 7

**Question:**

a Show that  $\sum_{r=1}^n r(r+2) = \frac{n}{6}(n+1)(2n+7)$ .

b Using this result, or otherwise, find in terms of  $n$ , the sum of  $3\log 2 + 4\log 2^2 + 5\log 2^3 + \dots + (n+2)\log 2^n$ .

**Solution:**

**a**

$$\begin{aligned}\sum_{r=1}^n r^2 + 2 \sum_{r=1}^n r &= \frac{n}{6}(n+1)(2n+1) + 2 \cdot \frac{n}{2}(n+1) \\ &= \frac{n}{6}(n+1)\{(2n+1)+6\} \\ &= \frac{n}{6}(n+1)(2n+7)\end{aligned}$$

**b**

$$\begin{aligned}\text{The series is : } \sum_{r=1}^n (r+2)\log 2^r &= \sum_{r=1}^n r(r+2)\log 2 \quad \text{as } \log 2^r = r \log 2 \\ &= \log 2 \sum_{r=1}^n r(r+2) \quad \text{as } \log 2 \text{ is a constant} \\ &= \frac{n}{6}(n+1)(2n+7) \log 2\end{aligned}$$

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise F, Question 8

##### Question:

Show that  $\sum_{r=n}^{2n} r^2 = \frac{n}{6}(n+1)(an+b)$ , where  $a$  and  $b$  are constants to be found.

##### Solution:

$$\begin{aligned}\sum_{r=n}^{2n} r^2 &= \sum_{r=1}^{2n} r^2 - \sum_{r=1}^{n-1} r^2 = \frac{(2n)(2n+1)(4n+1)}{6} - \frac{(n-1)n(2n-1)}{6} \\&= \frac{n}{6}\{2(8n^2 + 6n + 1) - (2n^2 - 3n + 1)\} \\&= \frac{n}{6}(14n^2 + 15n + 1) \\&= \frac{n}{6}(n+1)(14n+1) \quad a = 14, \quad b = 1\end{aligned}$$

© Pearson Education Ltd 2008

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise F, Question 9

**Question:**

a Show that  $\sum_{r=1}^n (r^2 - r - 1) = \frac{n}{3}(n-2)(n+2)$ .

b Hence calculate  $\sum_{r=10}^{40} (r^2 - r - 1)$ .

**Solution:**

a

$$\begin{aligned}\sum_{r=1}^n r^2 - \sum_{r=1}^n r - \sum_{r=1}^n 1 &= \frac{n}{6}(n+1)(2n+1) - \frac{n}{2}(n+1) - n \\&= \frac{n}{6}\{(n+1)(2n+1) - 3(n+1) - 6\} \\&= \frac{n}{6}(2n^2 - 8) \\&= \frac{n}{3}(n^2 - 4) \\&= \frac{n}{3}(n-2)(n+2)\end{aligned}$$

b  $\sum_{r=10}^{40} (r^2 - r - 1) = \sum_{r=1}^{40} (r^2 - r - 1) - \sum_{r=1}^9 (r^2 - r - 1)$

Substitute  $n = 40$  and  $n = 9$  into the result for part (a), and subtract.

The result is  $21280 - 230 = 21049$

© Pearson Education Ltd 2008

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise F, Question 10

##### Question:

a Show that  $\sum_{r=1}^n r(2r^2 + 1) = \frac{n}{2}(n+1)(n^2+n+1)$ .

b Hence calculate  $\sum_{r=26}^{58} r(2r^2 + 1)$ .

##### Solution:

a

$$\begin{aligned} 2 \sum_{r=1}^n r^3 + \sum_{r=1}^n r &= \frac{n^2(n+1)^2}{2} + \frac{n}{2}(n+1) \\ &= \frac{n}{2}(n+1)\{n(n+1)+1\} \\ &= \frac{n}{2}(n+1)(n^2+n+1) \end{aligned}$$

b Substitute  $n = 58$  and  $n = 25$  into the result for (a), and subtract. The result = 5654178.

© Pearson Education Ltd 2008

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise F, Question 11

**Question:**

Find

a  $\sum_{r=1}^n r(3r-1)$

b  $\sum_{r=1}^n (r+2)(3r+5)$

c  $\sum_{r=1}^n (2r^3 - 2r + 1)$ .

**Solution:**

a  $3 \sum_{r=1}^n r^2 - \sum_{r=1}^n r = \frac{n(n+1)(2n+1)}{2} - \frac{n(n+1)}{2} = \frac{n(n+1)}{2}(2n+1-1) = n^2(n+1)$

b

$$\begin{aligned} 3 \sum_{r=1}^n r^2 + 11 \sum_{r=1}^n r + 10 \sum_{r=1}^n 1 &= \frac{n(n+1)(2n+1)}{2} + \frac{11n(n+1)}{2} + 10n \\ &= \frac{n}{2} \{(2n^2 + 3n + 1) + 11(n+1) + 20\} \\ &= \frac{n}{2} (2n^2 + 14n + 32) = n(n^2 + 7n + 16) \end{aligned}$$

c

$$\begin{aligned} 2 \sum_{r=1}^n r^3 - 2 \sum_{r=1}^n r + \sum_{r=1}^n 1 &= \frac{n^2(n+1)^2}{2} - n(n+1) + n \\ &= \frac{n}{2} \{n(n+1)^2 - 2(n+1) + 2\} \\ &= \frac{n}{2} \{n(n+1)^2 - 2n\} = \frac{n^2}{2} (n^2 + 2n - 1) \end{aligned}$$

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise F, Question 12

##### Question:

a Show that  $\sum_{r=1}^n r(r+1) = \frac{n}{3}(n+1)(n+2)$ .

b Hence calculate  $\sum_{r=31}^{60} r(r+1)$ .

##### Solution:

a

$$\begin{aligned}\sum_{r=1}^n r^2 + \sum_{r=1}^n r &= \frac{n}{6}(n+1)(2n+1) + \frac{n}{2}(n+1) = \frac{n}{6}(n+1)\{2n+1+3\} \\ &= \frac{n}{3}(n+1)(n+2)\end{aligned}$$

b Substitute  $n = 60$  and  $n = 30$  into the result for part (a), and subtract. The result = 65720.

© Pearson Education Ltd 2008

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise F, Question 13

**Question:**

a Show that  $\sum_{r=1}^n r(r+1)(r+2) = \frac{n}{4}(n+1)(n+2)(n+3)$ .

b Hence evaluate  $3 \times 4 \times 5 + 4 \times 5 \times 6 + 5 \times 6 \times 7 + \dots + 40 \times 41 \times 42$ .

**Solution:**

a

$$\begin{aligned}\sum_{r=1}^n r^3 + 3 \sum_{r=1}^n r^2 + 2 \sum_{r=1}^n r &= \frac{n^2}{4}(n+1)^2 + \frac{n}{2}(n+1)(2n+1) + n(n+1) \\ &= \frac{n}{4}(n+1)\{(n(n+1) + 2(2n+1) + 4\} \\ &= \frac{n}{4}(n+1)(n+2)(n+3)\end{aligned}$$

b  $3 \times 4 \times 5 + 4 \times 5 \times 6 + 5 \times 6 \times 7 + \dots + 40 \times 41 \times 42 = \sum_{r=3}^{40} r(r+1)(r+2)$

$$\begin{aligned}\sum_{r=3}^{40} r(r+1)(r+2) &= \sum_{r=1}^{40} r(r+1)(r+2) - \sum_{r=1}^2 r(r+1)(r+2) \\ &= \frac{40 \times 41 \times 42 \times 43}{4} - \frac{2 \times 3 \times 4 \times 5}{4} = 740430\end{aligned}$$

© Pearson Education Ltd 2008

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise F, Question 14

##### Question:

a Show that  $\sum_{r=1}^n r\{2(n-r)+1\} = \frac{n}{6}(n+1)(2n+1)$ .

b Hence sum the series  $(2n-1) + 2(2n-3) + 3(2n-5) + \dots + n$

##### Solution:

a Series can be written as  $(2n+1) \sum_{r=1}^n r - 2 \sum_{r=1}^n r^2$  as  $n$  is a constant.

$$\begin{aligned} &= (2n+1)\frac{n}{2}(n+1) - \frac{n}{3}(n+1)(2n+1) \\ &= \frac{n}{6}(n+1)(2n+1) \end{aligned}$$

b  $\sum_{r=1}^n r[2(n-r)+1] = (2n-1) + 2[(2n-4)+1] + 3[(2n-6)+1] + \dots + n[2(n-n)+1]$

$= (2n-1) + 2(2n-3) + 3(2n-5) + \dots + n$ , the series in part (b).

The sum, therefore, is  $\frac{n}{6}(n+1)(2n+1)$

© Pearson Education Ltd 2008

# Solutionbank FP1

## Edexcel AS and A Level Modular Mathematics

### Series

#### Exercise F, Question 15

**Question:**

a Show that when  $n$  is even,

$$\begin{aligned} 1^3 - 2^3 + 3^3 - \dots - n^3 &= 1^3 + 2^3 + 3^3 + \dots + n^3 - 16 \left\{ 1^3 + 2^3 + 3^3 + \dots + \left(\frac{n}{2}\right)^3 \right\} \\ &= \sum_{r=1}^n r^3 - 16 \sum_{r=1}^{\frac{n}{2}} r^3. \end{aligned}$$

b Hence show that, for  $n$  even,  $1^3 - 2^3 + 3^3 - \dots - n^3 = -\frac{n^2}{4}(2n+3)$

c Deduce the sum of  $1^3 - 2^3 + 3^3 - \dots - 40^3$ .

**Solution:**

a

$$\begin{aligned} 1^3 - 2^3 + 3^3 - \dots - n^3 &= (1^3 + 2^3 + 3^3 + \dots + n^3) - 2(2^3 + 4^3 + 6^3 + \dots + n^3) \\ &= (1^3 + 2^3 + 3^3 + \dots + n^3) - 2 \left\{ 2^3 (1^3 + 2^3 + 3^3 + \dots + \left(\frac{n}{2}\right)^3) \right\} \text{ as } n \text{ is even} \\ &= (1^3 + 2^3 + 3^3 + \dots + n^3) - 16 \left\{ 1^3 + 2^3 + 3^3 + \dots + \left(\frac{n}{2}\right)^3 \right\} \\ &= \sum_{r=1}^n r^3 - 16 \sum_{r=1}^{\frac{n}{2}} r^3 \text{ [As } n \text{ is even, } \frac{n}{2} \text{ is an integer]} \end{aligned}$$

b

$$\begin{aligned} \sum_{r=1}^n r^3 - 16 \sum_{r=1}^{\frac{n}{2}} r^3 &= \frac{n^2}{4} (n+1)^2 - 16 \frac{\left(\frac{n}{2}\right)^2 \left(\frac{n}{2}+1\right)^2}{4} \\ &= \frac{n^2}{4} (n+1)^2 - 4 \frac{n^2}{4} \frac{(n+2)^2}{4} \\ &= \frac{n^2}{4} (n+1)^2 - \frac{n^2}{4} (n+2)^2 \\ &= \frac{n^2}{4} \{(n+1)^2 - (n+2)^2\} \\ &= \frac{n^2}{4} (-2n-3) = -\frac{n^2}{4} (2n+3) \end{aligned}$$

c Substituting  $n = 40$ , gives  $-33200$ .