

GCE Examinations
Advanced Subsidiary

Core Mathematics C3

Paper H

MARKING GUIDE

This guide is intended to be as helpful as possible to teachers by providing concise solutions and indicating how marks could be awarded. There are obviously alternative methods that would also gain full marks.

Method marks (M) are awarded for knowing and using a method.

Accuracy marks (A) can only be awarded when a correct method has been used.

(B) marks are independent of method marks.



Written by Shaun Armstrong

© Solomon Press


These sheets may be copied for use solely by the purchaser's institute.

C3 Paper H – Marking Guide

1. (a) $f(2) = -2$ M1 A1
 (b) $gf(x) = g(2 - x^2) = \frac{3(2 - x^2)}{2(2 - x^2) - 1} = \frac{6 - 3x^2}{3 - 2x^2}$ M1 A1
 $\therefore \frac{6 - 3x^2}{3 - 2x^2} = \frac{1}{2}, \quad 2(6 - 3x^2) = 3 - 2x^2$
 $x^2 = \frac{9}{4}, \quad x = \pm \frac{3}{2}$ M1 A1 (6)

2. $5(\sec^2 2\theta - 1) - 13 \sec 2\theta = 1$ M1
 $5 \sec^2 2\theta - 13 \sec 2\theta - 6 = 0$
 $(5 \sec 2\theta + 2)(\sec 2\theta - 3) = 0$ M1
 $\sec 2\theta = -\frac{2}{5}$ or 3 A1
 $\cos 2\theta = -\frac{5}{2}$ (no solutions) or $\frac{1}{3}$
 $2\theta = 70.529, 360 - 70.529, 360 + 70.529, 720 - 70.529$ B1 M1
 $= 70.529, 289.471, 430.529, 649.471$
 $\theta = 35.3^\circ, 144.7^\circ, 215.3^\circ, 324.7^\circ$ (1dp) A2 (7)

3. (a) $\frac{(2x-3)(x+3)}{(2x-3)(x-2)} = \frac{x+3}{x-2}$ M1 A2
 (b) $\ln(2x^2 + 3x - 9) - \ln(2x^2 - 7x + 6) = 2, \quad \ln \frac{2x^2 + 3x - 9}{2x^2 - 7x + 6} = 2$ M1
 $\ln \frac{x+3}{x-2} = 2, \quad \frac{x+3}{x-2} = e^2$ A1
 $x+3 = e^2(x-2)$
 $3 + 2e^2 = x(e^2 - 1)$ M1
 $x = \frac{2e^2 + 3}{e^2 - 1}$ A1 (7)



5. (a) $t = 3, N = 18000 \Rightarrow 18000 = 2000e^{3k}, \quad e^{3k} = 9$ M1
 $k = \frac{1}{3} \ln 9 = 0.732$ (3sf) M1 A1
 (b) $4000 = 2000e^{0.7324t}$ M1
 $t = \frac{1}{0.7324} \ln 2 = 0.9464$ hours M1 A1
 \therefore doubles in 57 minutes (nearest minute) A1
 (c) $N = 2000e^{0.7324t}, \quad \frac{dN}{dt} = 0.7324 \times 2000e^{0.7324t} = 1465e^{0.7324t}$ M1 A1
 when $t = 3, \frac{dN}{dt} = 13200 \therefore$ increasing at rate of 13200 per hour (3sf) A1 (10)

6.	(a)	$\frac{d}{dx}(\sec x) = \frac{d}{dx}[(\cos x)^{-1}]$ $= -(\cos x)^{-2} \times (-\sin x)$ $= \frac{\sin x}{\cos^2 x} = \frac{1}{\cos x} \times \frac{\sin x}{\cos x}$ $= \sec x \tan x$	M1 A1 M1 A1
	(b)	$\frac{dy}{dx} = 2e^{2x} \times \sec x + e^{2x} \times \sec x \tan x = e^{2x} \sec x(2 + \tan x)$ $x = 0, y = 1, \text{ grad} = 2$ $\therefore y = 2x + 1$	M1 A1 M1 A1
	(c)	SP: $e^{2x} \sec x(2 + \tan x) = 0$ $\tan x = -2$ $x = -1.11 \text{ (2dp)}$	M1 M1 A1 (11)

7.	(a)	$f(x) = (x - 1)^2 - 1 + 5 = (x - 1)^2 + 4$	M1 A1
	(b)	$f(x) \geq 4$	B1
	(c)	$y = (x - 1)^2 + 4$ $(x - 1)^2 = y - 4$ $x - 1 = \pm \sqrt{y - 4}$ $x = 1 \pm \sqrt{y - 4}$ $f^{-1}(x) = 1 + \sqrt{x - 4}$	M1 M1 A1
	(d)	translation by 4 units in negative x direction translation by 1 unit in negative y direction (either first)	B2
	(e)	$\frac{dy}{dx} = \frac{1}{2}(x - 4)^{-\frac{1}{2}}$ $x = 8, y = 3, \text{ grad} = \frac{1}{4}$ $\therefore \text{grad of normal} = -4$ $\therefore y - 3 = -4(x - 8) \quad [y = 35 - 4x]$	M1 A1 M1 A1 (12)

8.	(a)	$\frac{dy}{dx} = -e^2 x^{-2} + e^x$	M1 A1
	(b)	SP: $-e^2 x^{-2} + e^x = 0$ let $f(x) = -e^2 x^{-2} + e^x$ $f(1.3) = -0.70, f(1.4) = 0.29$ sign change, $f(x)$ continuous \therefore root	M1 M1 A1
	(c)	$x = 2, y = \frac{3}{2} e^2, \text{ grad} = \frac{3}{4} e^2$ $\therefore y - \frac{3}{2} e^2 = \frac{3}{4} e^2(x - 2)$ $y = \frac{3}{4} e^2 x$ $\therefore x = 0 \Rightarrow y = 0$ so passes through origin	M1 M1 A1 A1
	(d)	$x_1 = -1.125589, x_2 = -1.125803, x_3 = -1.125804$ (7sf) $\therefore x\text{-coordinate of } B = -1.1258$ (5sf)	M1 A2 A1 (13)

Total (75)

Performance Record – C3 Paper H