

Question Number	Scheme	Marks
1.	$\int_1^3 x^2 \, dx = \left[\frac{x^3}{3} \ln x \right] - \int \frac{x^3}{3} \frac{1}{x} \, dx$ $= \left[\frac{x^3}{3} \ln x \right] - \int \frac{x^2}{3} \, dx$ $= \left[\frac{x^3}{3} \ln x \right] - \frac{1}{9} x^3$ $= 9 \ln 3 - 3 + \frac{1}{9} = 9 \ln 3 - 2\frac{8}{9}$	M1 A1 M1 A1 M1 A1 (6) (6 marks)
2. (a)	$\frac{dV}{dt} = \pm c\sqrt{V} \text{ or } \frac{dV}{dt} \propto \sqrt{V}$ As $V = Ah$, $\frac{dV}{dh} = A$ or $V \propto h$ $\therefore \text{use chain rule to obtain } \frac{dh}{dt} = -\frac{c}{A}\sqrt{V} = \frac{-c}{\sqrt{A}}\sqrt{h} = -k\sqrt{h}$	M1 M1 A1 (3)
(b)	$\int \frac{dh}{h} = - \int k dt$ $2h^{\frac{1}{2}} = A - kt$ $h^{\frac{1}{2}} = \frac{A}{2} - \frac{kt}{2}$ $h = (A - Bt)^2 \quad *$	M1 M1 A1 A1 (4)
(c)	$t = 0, h = 1: \quad A = 1$ $t = 5, h = 0.5: \quad 0.5 = (1 - 5B)^2$ $B = \frac{(1 - \sqrt{0.5})}{5} \quad (B = 0.0586)$ $h = 0, t = \frac{A}{B} = \frac{5}{1 - \sqrt{0.5}} = 17.1 \text{ min}$	B1 B1 B1 (3)
(d)	$h = \frac{A^2}{4} = 0.25 \text{ m}$	M1 A1 (2) (12 marks)

Question Number	Scheme	Marks
3. (a)	$\cos(A+A) = \cos^2 A - \sin^2 A$ $= \cos^2 A - (1 - \cos^2 A) = 2\cos^2 A - 1$	M1 A1 (2)
(b)	$[x = 2, \theta = \frac{\pi}{4}; x = \sqrt{6}, \theta = \frac{\pi}{3}]$ $x = 2\sqrt{2} \sin \theta, \frac{dx}{d\theta} = 2\sqrt{2} \cos \theta$ $\int \sqrt{8-x^2} dx = \int 2\sqrt{2} \cos \theta 2\sqrt{2} \cos \theta d\theta = \int 8 \cos^2 \theta d\theta$ Using $\cos 2\theta = 2\cos^2 \theta - 1$ to give $\int 4(1 + \cos 2\theta) d\theta = 4\theta + 2 \sin 2\theta$ Substituting limits to give $\frac{1}{3}\pi + \sqrt{3} - 2$ or given result	B1 B1 M1 A1 M1 A1 ft (7)
(c)	$\frac{dy}{d\theta} = \frac{-2 \sin 2\theta}{1 + \cos 2\theta}$ Using the chain rule, with $\frac{dx}{d\theta} = \sec \theta \tan \theta$ to give $\frac{dy}{dx} (= -2 \cos \theta)$ Gradient at the point where $\theta = \frac{\pi}{3}$ is -1 Equation of tangent is $y + \ln 2 = -(x - 2)$ (or equivalent answer)	B1 M1 A1 ft M1 A1 (5) (14 marks)
4. (a)	$A: y = 16, B: y = 2$	B1 (1)
(b)	$y(x-3) = 4, yx - 3y = 4$ $x = \frac{3y+4}{y}$ (*)	M1 (1)
(c)	$x^2 = \left(3 + \frac{4}{y}\right)^2 = 9 + \frac{24}{y} + \frac{16}{y^2}$ $\int x^2 dy = \int (9 + 24y^{-1} + 16y^{-2}) dy$ $= 9y - \frac{16}{y}, + 24 \ln y$ $\left[9y + 24 \ln y - 16y^{-1}\right]_2^{16} = (144 + 24 \ln 16 - 1) - (18 + 24 \ln 2 - 8)$ $V = \pi(133 + 24 \ln 8)$	M1 A1 M1 A1 ft, A1 ft M1 A1 (7)
(d)	$V \times 27 \approx 15500$ (*)	M1 A1 (2) (11 marks)

Question Number	Scheme	Marks
5. (a)	$\overrightarrow{AB} = (2\mathbf{i} - \mathbf{j} + 2\mathbf{k})$, $\overrightarrow{CB} = (-2\mathbf{i} + 2\mathbf{j} + \mathbf{k})$, (or $\overrightarrow{BA}, \overrightarrow{BC}$, or $\overrightarrow{AB}, \overrightarrow{BC}$ stated in above form or column vector form) $\cos A\hat{B}C = \frac{\overrightarrow{CB} \bullet \overrightarrow{AB}}{ \overrightarrow{CB} \overrightarrow{AB} } = -\frac{4}{9}$	M1 A1 M1 A1 (4)
(b)	Area of $\Delta ABC = \frac{1}{2} \times 3 \times 3 \times \sin B$ $\sin B = \sqrt{(1 - \frac{16}{81})} = \frac{\sqrt{65}}{9}$ $\therefore \text{Area} = \frac{1}{2} \sqrt{65}$	M1 M1 A1 (3)
(c)	$\overrightarrow{AC} = \begin{pmatrix} 4 \\ -3 \\ 1 \end{pmatrix}$ $\overrightarrow{DC} = \begin{pmatrix} 0 \\ -1 \\ -3 \end{pmatrix}$ or given in alternative form with attempt at scalar product $\overrightarrow{AC} \bullet \overrightarrow{DC} = 0$, therefore the lines are perpendicular.	M1 A1 (2)
(d)	$\overrightarrow{AD} = \begin{pmatrix} 4 \\ -2 \\ 4 \end{pmatrix}$ $\overrightarrow{DB} = \begin{pmatrix} -2 \\ 1 \\ -2 \end{pmatrix}$ and $AD:DB = 2:-1$ (allow 2:1)	M1 A1 (2)
		(11 marks)

Question Number	Scheme	Marks														
6. (a)	<table border="1"> <tr> <td>Distance from one side (m)</td><td>0</td><td>2</td><td>4</td><td>6</td><td>8</td><td>10</td></tr> <tr> <td>Height (m)</td><td>0</td><td>6.13</td><td>7.80</td><td>7.80</td><td>6.13</td><td>0</td></tr> </table> <p style="text-align: center;">“y” = 7.80 when “x” = 4 or 6 Symmetry</p>	Distance from one side (m)	0	2	4	6	8	10	Height (m)	0	6.13	7.80	7.80	6.13	0	
Distance from one side (m)	0	2	4	6	8	10										
Height (m)	0	6.13	7.80	7.80	6.13	0										
(b)	<p>Estimate area = $\frac{2}{2} [0 + 2(6.13 + 7.80 + 7.80 + 6.13)]$ $= 55.7 \text{ m}^2$</p>	B1 M1 A1ft														
(c)	$140 - (b) = 84.3 \text{ m}^2$	A1 (4)														
(d)	Over-estimate; reason, e.g. area under curve is under-estimate (due to curvature)	A1 ft (1) B1 B1 (2)														
		(9 marks)														
7. (a)	<p>Method using either</p> $\frac{A}{(1-x)} + \frac{B}{(2x+3)} + \frac{C}{(2x+3)^2} \text{ or } \frac{A}{1-x} + \frac{Dx+E}{(2x+3)^2}$ $A = 1, C = 10, B = 2 \quad \text{or} \quad D = 4 \text{ and } E = 16$	M1 B1 A1 A1 (4)														
(b)	$\int [\frac{1}{1-x} + \frac{2}{2x+3} + 10(2x+3)^{-2}] dx \text{ or } \int \frac{A}{1-x} + \frac{Dx+E}{(2x+3)^2} dx$ $- \ln 1-x + \ln 2x+3 - 5(2x+3)^{-1} (+c) \text{ or}$ $- \ln 1-x + \ln 2x+3 - (2x+8)(2x+3)^{-1} (+c)$	M1 M1 A1ft A1ft A1ft (5)														
(c)	$(1-x)^{-1} + 2(3+2x)^{-1} + 10(3+2x)^{-2} =$ $1+x+x^2+\dots$ $+ \frac{2}{3}(1-\frac{2x}{3}+\frac{4x^2}{9}\dots)$ $+ \frac{10}{9}(1+(-2)(\frac{2x}{3})+\frac{(-2)(-3)}{2}(\frac{2x}{3})^2+\dots)$ $= \frac{25}{9} - \frac{25}{27}x + \frac{25}{9}x^2\dots$	M1 A1 M1 A1 A1 M1 A1 (7)														
		(16 marks)														