

GCE Examinations
Advanced Subsidiary / Advanced Level
Pure Mathematics
Module P6

Paper G

MARKING GUIDE

This guide is intended to be as helpful as possible to teachers by providing concise solutions and indicating how marks should be awarded. There are obviously alternative methods that would also gain full marks.

Method marks (M) are awarded for knowing and using a method.

Accuracy marks (A) can only be awarded when a correct method has been used.

(B) marks are independent of method marks.



Written by Rosemary Smith & Shaun Armstrong

© Solomon Press

These sheets may be copied for use solely by the purchaser's institute.


P6 Paper G – Marking Guide

1. $\det \mathbf{A} = 3(0+k) - 1(0+2) - 4(k-4)$ M1
 $= 3k - 2 - 4k + 16 = 14 - k$ A1
 \therefore singular if $k = 14$ A1 **(3)**

2. $| -4 + 4\sqrt{3}i | = 4\sqrt{1+3} = 8$; $\arg(-4 + 4\sqrt{3}i) = \arctan(-\sqrt{3}) = \frac{2\pi}{3}$ M1
 $\therefore (re^{i\theta})^3 = 8e^{i\frac{2\pi}{3}}$ A1
 $r^3 = 8$ so $r = 2$ A1
 $3\theta = 2n\pi + \frac{2\pi}{3}$ M1
 $n = 0, 1, 2$ gives $\theta = \frac{2\pi}{9}, \frac{8\pi}{9}, \frac{14\pi}{9}$ A1
 $\therefore z = 2(\cos \frac{2\pi}{9} + i\sin \frac{2\pi}{9}), 2(\cos \frac{8\pi}{9} + i\sin \frac{8\pi}{9}), 2(\cos \frac{14\pi}{9} + i\sin \frac{14\pi}{9})$ A1 **(6)**

3. assume true for $n = k$ $\therefore f(k) = k(k^2 + 5)$ is divisible by 6 M1
 $f(k+1) = (k+1)[(k+1)^2 + 5]$
 $= (k+1)(k^2 + 2k + 6)$
 $f(k+1) - f(k) = (k+1)(k^2 + 2k + 6) - k(k^2 + 5)$ M1
 $= k^3 + 2k^2 + 6k + k^2 + 2k + 6 - k^3 - 5k$
 $= 3k^2 + 3k + 6 = 3k(k+1) + 6$
 $\therefore f(k+1) = 3k(k+1) + 6 + f(k)$ A1
 $k, (k+1)$ are consec. integers $\therefore k(k+1)$ is div. by 2 [one must be even] M1
 $\therefore 3k(k+1)$ is div. by 6 $\therefore f(k+1)$ is div. by 6 A1
 \therefore true for $n = k+1$ if true for $n = k$
if $n = 1$ $f(1) = 1 \times 6 = 6$ $\therefore f(1)$ is div. by 6 \therefore true for $n = 1$ B1
 \therefore by induction true for $n \in \mathbb{Z}^+$ A1 **(7)**

4. (a) $|z - (1 - 2i)| = 3$ \therefore circle, centre $1 - 2i$, radius 3 M1 A1



B1

- (b) T : enlargement s.f 4, centre O
giving circle, centre $4 - 8i$, radius 12 M1 A1
 U : translation through $5 - i$
giving circle centre $6 - 3i$, radius 3 M1 A1
 V : anticlockwise rotation through $\frac{\pi}{2}$ about O
giving circle centre $2 + i$, radius 3 M1 A1 **(9)**
-

5.	(a)	$f(x) = \cos x, f'(x) = -\sin x, f''(x) = -\cos x, f'''(x) = \sin x$	M1 A1
		$f\left(\frac{\pi}{6}\right) = \frac{\sqrt{3}}{2}, f'\left(\frac{\pi}{6}\right) = -\frac{1}{2}, f''\left(\frac{\pi}{6}\right) = -\frac{\sqrt{3}}{2}, f'''\left(\frac{\pi}{6}\right) = \frac{1}{2}$	A1
		$f(x) = \frac{\sqrt{3}}{2} - \frac{1}{2}(x - \frac{\pi}{6}) + \frac{1}{2!}(-\frac{\sqrt{3}}{2})(x - \frac{\pi}{6})^2 + \frac{1}{3!}(\frac{1}{2})(x - \frac{\pi}{6})^3 + \dots$	M1
		$f(x) = \frac{\sqrt{3}}{2} - \frac{1}{2}(x - \frac{\pi}{6}) - \frac{\sqrt{3}}{4}(x - \frac{\pi}{6})^2 + \frac{1}{12}(x - \frac{\pi}{6})^3 + \dots$	A1
	(b)	if $x = \frac{\pi}{4}, x - \frac{\pi}{6} = \frac{\pi}{12}$	M1
		$\therefore \cos \frac{\pi}{4} = \frac{\sqrt{3}}{2} - \frac{1}{2}(\frac{\pi}{12}) - \frac{\sqrt{3}}{4}(\frac{\pi}{12})^2 + \frac{1}{12}(\frac{\pi}{12})^3 + \dots$	M1
		$= 0.7069 \text{ (4dp)}$	A1
	(c)	$\% \text{ error} = \frac{\cos \frac{\pi}{4} - 0.7069}{\cos \frac{\pi}{4}} \times 100\% = 0.023 \% \text{ (2sf)}$	M1 A1 (10)
6.	(a)	$\frac{d^3y}{dx^3} = 2x + x \frac{dy}{dx} + y - 2y \frac{dy}{dx}$	M1 A1
		$x_0 = 0, y_0 = \frac{1}{2}, \left(\frac{dy}{dx}\right)_0 = -1; \left(\frac{d^2y}{dx^2}\right)_0 = 0 + 0 - \frac{1}{4} = -\frac{1}{4}$	A1
		$\left(\frac{d^3y}{dx^3}\right)_0 = 0 + 0 + \frac{1}{2} - [2 \times \frac{1}{2} \times (-1)] = \frac{3}{2}$	A1
		$\therefore y = \frac{1}{2} - 1x + \frac{1}{2!}(-\frac{1}{4})x^2 + \frac{1}{3!}(\frac{3}{2})x^3 + \dots$	M1
		$y = \frac{1}{2} - x - \frac{1}{8}x^2 + \frac{1}{4}x^3 + \dots$	A1
	(b)	$x = -0.1, y \approx 0.5 + 0.1 - 0.00125 - 0.00025 = 0.5985$	A1
	(c)	$\frac{y_1 - 2y_0 + y_{-1}}{0.01} = x_0^2 + x_0y_0 - y_0^2$	M1
		$y_1 - 2y_0 + y_{-1} = 0.01(x_0^2 + x_0y_0 - y_0^2)$	
		$y_1 = 2y_0 - y_{-1} + 0.01(x_0^2 + x_0y_0 - y_0^2)$	A1
		$x_{-1} = -0.1, x_0 = 0, x_1 = 0.1; y_{-1} = 0.5985, y_0 = 0.5, y_1 = ?$	
		$y_1 = 1 - 0.5985 + 0.01(0 + 0 - 0.25) = 0.399$	A1 (10)

7. (a)
$$\begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & -4 & 1 \\ 4 & -2 & 5 \end{vmatrix}$$

 $= \mathbf{i}(-20 + 2) - \mathbf{j}(10 - 4) + \mathbf{k}(-4 + 16) = -18\mathbf{i} - 6\mathbf{j} + 12\mathbf{k}$ M1 A1
 $|\mathbf{a} \times \mathbf{b}| = |\mathbf{a}| |\mathbf{b}| \sin \theta$
 $6\sqrt{9+1+4} = \sqrt{21}\sqrt{45} \sin \theta$ M1 A1
 $6\sqrt{7}\sqrt{2} = 3\sqrt{7}\sqrt{3}\sqrt{5}\sin \theta$
 $\sin \theta = \frac{2\sqrt{2}}{\sqrt{15}}$ or $\frac{2}{15}\sqrt{30}$ A1
- (b) $\mathbf{n} = 3\mathbf{i} + \mathbf{j} - 2\mathbf{k}$ M1
 $\mathbf{r} \cdot (3\mathbf{i} + \mathbf{j} - 2\mathbf{k}) = (2\mathbf{i} - \mathbf{j} + 2\mathbf{k}) \cdot (3\mathbf{i} + \mathbf{j} - 2\mathbf{k}) = 6 - 1 - 4 = 1$ M1 A1
 $\mathbf{r} \cdot (3\mathbf{i} + \mathbf{j} - 2\mathbf{k}) = 1 \therefore 3x + y - 2z - 1 = 0$ A1
- (c) $\mathbf{r} = \mathbf{i} - 2\mathbf{j} + u(2\mathbf{j} + \mathbf{k})$
 $u = 0, \mathbf{r} = \mathbf{i} - 2\mathbf{j}, (\mathbf{i} - 2\mathbf{j}) \cdot (3\mathbf{i} + \mathbf{j} - 2\mathbf{k}) = 3 - 2 = 1 \therefore$ in plane M1 A1
 $u = 1, \mathbf{r} = \mathbf{i} + \mathbf{k}, (\mathbf{i} + \mathbf{k}) \cdot (3\mathbf{i} + \mathbf{j} - 2\mathbf{k}) = 3 - 2 = 1 \therefore$ in plane M1
two points on line in plane \therefore line in plane A1 **(13)**
-

8. (a) $(\mathbf{AB})(\mathbf{B}^{-1}\mathbf{A}^{-1}) = \mathbf{AB}\mathbf{B}^{-1}\mathbf{A}^{-1} = \mathbf{A}\mathbf{I}\mathbf{A}^{-1} = \mathbf{AA}^{-1} = \mathbf{I}$ M1 A1
 $\therefore (\mathbf{B}^{-1}\mathbf{A}^{-1})$ is inverse of (\mathbf{AB}) i.e. $(\mathbf{AB})^{-1} = \mathbf{B}^{-1}\mathbf{A}^{-1}$ M1 A1
- (b) $S(a_1\mathbf{v}_1 + a_2\mathbf{v}_2) = S\begin{pmatrix} a_1x_1 + a_2x_2 \\ a_1y_1 + a_2y_2 \end{pmatrix}$ M1
 $= \begin{pmatrix} a_1y_1 + a_2y_2 - a_1x_1 - a_2x_2 \\ 2a_1x_1 + 2a_2x_2 + a_1y_1 + a_2y_2 \end{pmatrix}$ M1 A1
 $= \begin{pmatrix} a_1(y_1 - x_1) + a_2(y_2 - x_2) \\ a_1(2x_1 + y_1) + a_2(2x_2 + y_2) \end{pmatrix}$ M1
 $= a_1\begin{pmatrix} y_1 - x_1 \\ 2x_1 + y_1 \end{pmatrix} + a_2\begin{pmatrix} y_2 - x_2 \\ 2x_2 + y_2 \end{pmatrix}$ A1
 $= a_1S(\mathbf{v}_1) + a_2S(\mathbf{v}_2) \therefore S$ is a linear transformation M1 A1
- (c) $\mathbf{S} = \begin{pmatrix} -1 & 1 \\ 2 & 1 \end{pmatrix}, \mathbf{T} = \begin{pmatrix} 3 & 0 \\ 1 & 1 \end{pmatrix}$ M1 A1
 $\mathbf{ST} = \begin{pmatrix} -1 & 1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} 3 & 0 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} -2 & 1 \\ 7 & 1 \end{pmatrix}$ M1 A1
 $\det(\mathbf{ST}) = -2 - 7 = -9$ M1
 $\therefore (\mathbf{ST})^{-1} = -\frac{1}{9} \begin{pmatrix} 1 & -1 \\ -7 & -2 \end{pmatrix}$ or $\frac{1}{9} \begin{pmatrix} -1 & 1 \\ 7 & 2 \end{pmatrix}$ A1 **(17)**
-

Total **(75)**

Performance Record – P6 Paper G