

# GCE Examinations

# Pure Mathematics

# Module P4

Advanced Subsidiary / Advanced Level

## Paper B

Time: 1 hour 30 minutes

### *Instructions and Information*

---

Candidates may use any calculator except those with a facility for symbolic algebra and/or calculus.

Full marks may be obtained for answers to ALL questions.

Mathematical and statistical formulae and tables are available.

This paper has 7 questions.

### *Advice to Candidates*

---

You must show sufficient working to make your methods clear to an examiner.  
Answers without working will gain no credit.



*Written by Shaun Armstrong & Chris Huffer*

© *Solomon Press*

*These sheets may be copied for use solely by the purchaser's institute.*

1. Find the set of values of  $x$  for which

$$|2x^2 - 5x| < x. \quad \text{(6 marks)}$$

---

2. (a) Sketch the curve  $C$  with the polar equation

$$r^2 = a^2 \sin^2 2\theta, \quad 0 \leq \theta < 2\pi. \quad \text{(3 marks)}$$

- (b) Find the exact area of the region enclosed by one loop of the curve  $C$ . **(5 marks)**
- 

3. (a) Show that

$$\sum_{r=1}^n (r^2 + 1)(r - 1) = \frac{1}{12} n(n - 1)(3n^2 + 5n + 8). \quad \text{(6 marks)}$$

- (b) Hence evaluate

$$\sum_{r=5}^{25} (r^2 + 1)(r - 1). \quad \text{(3 marks)}$$

---

4. (a) Find the general solution of the differential equation

$$\frac{dy}{dx} - y \cot x = \sin 2x. \quad \text{(6 marks)}$$

- (b) Given also that  $y = 2$  when  $x = \frac{\pi}{6}$ , find the exact value of  $y$  when  $x = \frac{2\pi}{3}$ . **(3 marks)**
- 

5.  $f(x) \equiv x^3 - \ln(4 - x^2), \quad x \in \mathbb{R}, \quad -2 < x < 2.$

- (a) Show that one root,  $\alpha$ , of the equation  $f(x) = 0$  lies in the interval  $1.0 < \alpha < 1.1$

**(2 marks)**

- (b) Starting with  $x = 1.0$ , show that using the Newton-Raphson method twice gives an approximation to  $\alpha$  that is correct to 6 decimal places.

**(8 marks)**

---

6. The complex numbers  $z_1$ ,  $z_2$  and  $z_3$  are given by

$$z_1 = 7 - i, \quad z_2 = 1 + i\sqrt{3}, \quad z_3 = a + ib,$$

where  $a$  and  $b$  are rational constants.

Given that the modulus of  $z_1z_3$  is 50,

(a) find the modulus of  $z_3$ . **(3 marks)**

Given also that the argument of  $\frac{z_2}{z_3}$  is  $\frac{7\pi}{12}$ ,

(b) find the argument of  $z_3$ . **(3 marks)**

(c) Find the values of  $a$  and  $b$ . **(2 marks)**

(d) Show that  $\frac{z_1}{z_3} = \frac{1}{5}(4 + 3i)$ . **(3 marks)**

(e) Represent  $z_1$ ,  $z_3$  and  $\frac{z_1}{z_3}$  on the same Argand diagram. **(2 marks)**

(f) By considering the modulus and argument of  $z_1$  and  $z_3$ , explain why  $\frac{z_3}{z_1} = \left(\frac{z_1}{z_3}\right)^*$ .

**(2 marks)**

*Turn over*

7. (a) Given that  $x = e^t$ , find  $\frac{dy}{dx}$  in terms of  $\frac{dy}{dt}$  and show that

$$\frac{d^2y}{dx^2} = e^{-2t} \left( \frac{d^2y}{dt^2} - \frac{dy}{dt} \right). \quad \text{(5 marks)}$$

- (b) Show that the substitution  $x = e^t$  transforms the differential equation

$$x^2 \frac{d^2y}{dx^2} - x \frac{dy}{dx} - 3y = 6x^2$$

into the differential equation

$$\frac{d^2y}{dt^2} - 2 \frac{dy}{dt} - 3y = 6e^{2t}. \quad \text{(3 marks)}$$

- (c) Given that when  $x = 1$ ,  $y = 3$  and  $\frac{dy}{dx} = -5$ , solve the differential equation

$$x^2 \frac{d^2y}{dx^2} - x \frac{dy}{dx} - 3y = 6x^2. \quad \text{(10 marks)}$$

---

**END**