

## **STATISTICS 2 (A) TEST PAPER 6 : ANSWERS AND MARK SCHEME**

1.	(a) List of all drivers insured with that company (b) Individual drivers	B1 B1; B2	4
2.	(a) Continuous uniform distribution on $[0, 360]$ (b) Mean = 180      s.d. = $\sqrt{(360^2 \div 12)} = \sqrt{10800} = 103.9$	B1 B1 M1 A1	4
3.	(a) Mean = $322/150 = 2.15$ , variance = $982/150 - 2.147^2 = 1.94$ (b) Mean $\approx$ Variance, which suggests Poisson (c) Assuming $\lambda = 2.14666$ , no. of errors in 6 pages is $\text{Po}(12.88)$ then $P(X \geq 15) > P(X = 15) = 0.0868 > 5\%$ , so do not reject $H_0$	M1 A1 M1 A1 B1 B1 B1 M1 A1 A1	10
4.	(a) $X \sim B(5, 0.2)$ $P(X = 0) = 0.8^5 = 0.3277$ (b) $P(X > 2) = 1 - P(X \leq 2) = 1 - 0.9421 = 0.0579$ (from tables) (c) No of lates in 7 weeks is distributed $B(35, 0.2) \approx N(7, 5.6)$ $P(X > 10) = P(X > 10.5) = P(Z > 3.5/\sqrt{5.6}) = P(Z > 1.48)$ $= 1 - 0.9306 = 0.0694$	M1 A1 M1 A1 A1 M1 A1 M1 A1 A1 M1 A1	12
5.	(a) $X \sim \text{Po}(4)$ , so $P(X < 2) = 0.0916$ (b) $P(3 < X < 10) = 0.9919 - 0.4335 = 0.558$ (c) $H_0$ : mean number of bubbles is still 0.1 per $\text{cm}^3$ ; $H_1$ : mean $< 0.1$ Under $H_0$ , no. of bubbles in 60 $\text{cm}^3$ is $\text{Po}(6)$ Then $P(X \leq 1) = 0.0174$ , so do not reject $H_0$ at 1% level	B1 M1 A1 M1 M1 A1 B1 B1 B1 M1 A1 A1	12
6.	(a) $f'(x) = \frac{4}{27}(6x - 3x^2) = 0$ when $x = 2$ , so mode = 2 (b) $E(x) = \frac{4}{27} \int_0^3 3x^3 - x^4 dx = \frac{4}{27} \left[ \frac{3x^4}{4} - \frac{x^5}{5} \right]_0^3 = 1.8$ (c) $F(x) = 0$ ( $x < 0$ ), $F(x) = \frac{4}{27} \left( x^3 - \frac{x^4}{4} \right)$ ( $0 \leq x \leq 3$ ), $F(x) = 1$ ( $x > 3$ ) (d) For median $m$ , $F(m) = 0.5$ Now $F(1.84) = 0.498 < 0.5$ and $F(1.85) = 0.504 > 0.5$ , so $1.84 < m < 1.85$ (e) Mean $<$ median $<$ mode      Negative skew	M1 A1 A1 M1 A1 A1 B1 M1 A1 B1 M1 A1 A1 A1 B1	15
7.	(a) $\int_2^{10} f(x) dx = 1$ , so $k \int_2^{10} (-x^2 + 12x - 20) dx = 1$ $k \left[ -\frac{1}{3}x^3 + 6x^2 - 20x \right]_2^{10} = 1$ $\frac{256k}{3} = 1$ $k = \frac{3}{256}$ By symmetry of parabola, mean = 6 (b) $E(X^2) = \frac{3}{256} \int_2^{10} (-x^4 + 12x^3 - 20x^2) dx$ $= \frac{3}{256} \left[ -\frac{x^5}{5} + 3x^4 - \frac{20x^3}{3} \right]_2^{10} = 39.2$ $\text{Var}(X) = 3.2$ Standard deviation = $\sqrt{3.2} = 1.789$ , i.e. £1789 (c) $P(X > 8) = \int_8^{10} f(x) dx = \frac{3}{256} \left[ -\frac{1}{3}x^3 + 6x^2 - 20x \right]_8^{10} = 0.156$ (d) $0.156^4 = 0.000592$	M1 M1 A1 M1 A1 B1 M1 A1 M1 A1 A1 A1 M1 A1 M1 A1 M1 A1	18