

Calculus

Differentiation



Notation

- The function you get from differentiating y with respect to x is called the DERIVATIVE of y and it's written $\frac{dy}{dx}$.
- $\frac{dy}{dx}$ is the rate of change of y with respect to x .
It is the gradient of the curve/the tangent to the curve.
- The notation $f'(x)$ (f prime of x) is sometimes used instead of $\frac{dy}{dx}$.




Differentiation from first principle.

Consider two points on a curve $A(x, f(x))$ and a point B close to A $B(x+h, f(x+h))$ where h is "small".

The chord AB has gradient $\frac{f(x+h) - f(x)}{x+h-x} = \frac{f(x+h) - f(x)}{h}$.

When B get closer and closer to A, h tends to 0.

If $\frac{f(x+h) - f(x)}{h}$ has a value when h tends to 0, this value is the gradient of the curve at A: $f'(x)$.



Example: $f(x) = x^2$

Let's work out the gradient of the curve at $x = 3$.

- $A(3, 3^2)$ and $B(3+h, (3+h)^2)$

$$\text{the gradient of AB: } m = \frac{(3+h)^2 - 3^2}{3+h-3} = \frac{9+6h+h^2-9}{h} = 6+h$$

When h tends to 0, m tends to 6:

Conclusion: $\frac{dy}{dx}(x=3) = f'(3) = 6$



Differentiating polynomials

- if $y = x^n$ then $\frac{dy}{dx} = nx^{n-1}$
- if $y = x^n + x^p$ then $\frac{dy}{dx} = nx^{n-1} + px^{p-1}$
- if $y = k \times x^n$ then $\frac{dy}{dx} = k \times nx^{n-1}$ where $k \in \mathbb{R}$.

Example: $y = x^4$ $\frac{dy}{dx} = 4x^3$

$$y = 5x^6 \quad \frac{dy}{dx} = 5 \times 6x^5 = 30x^5$$

$$y = 3x^4 + 5x^3 + x \quad \frac{dy}{dx} = 12x^3 + 15x^2 + 1$$