

980/01

MATHEMATICS M1

Mechanics 1

P.M. TUESDAY, 6 June 2006

(1½ hours)

ADDITIONAL MATERIALS

In addition to this examination paper, you will need:

- a 12 page answer book;
- a Formula Booklet;
- a calculator.

INSTRUCTIONS TO CANDIDATES

Answer **all** questions.


Take g as 9.8 ms^{-2} .

INFORMATION FOR CANDIDATES

The number of marks is given in brackets at the end of each question or part-question.

You are reminded of the necessity for good English and orderly presentation in your answers.


1. A lift, starting from rest, descends with a uniform acceleration of 3 ms^{-2} until it reaches a speed of 9 ms^{-1} . It then travels at a constant speed of 9 ms^{-1} for a short time and finally, it is brought to rest with a uniform retardation of 2 ms^{-2} . An object, of mass 6 kg , is on the floor of the lift. Calculate the magnitude of the reaction of the floor on the object during each of the three stages of the motion. [5]
2. A train, starting from rest from station A , travels along a straight horizontal track until it stops at station B , which is 2400 m from A . Initially, the train accelerates at a uniform rate of 0.4 ms^{-2} until it reaches a speed of 16 ms^{-1} . It then maintains this speed of 16 ms^{-1} for $T \text{ s}$, before decelerating uniformly to rest in 20 s .
- (a) Calculate the time taken for accelerating. [2]
- (b) Draw a sketch of the v - t graph for the journey from A to B . [4]
- (c) Find the value of T . [4]
3. A box of mass 20 kg is at rest on a rough horizontal floor. The coefficient of friction between the box and the floor is 0.3 . The box is subjected to a horizontal force of $T \text{ N}$.
- (a) Given that $T = 65$, find the magnitude of the acceleration of the box. [5]
- (b) Given that $T = 45$, find the magnitude of the frictional force. [1]
4. The diagram shows two particles A and B , of mass 3.1 kg and 1.8 kg respectively, connected by a light inextensible string passing over a fixed smooth pulley. Initially, B is held at rest with the string taut. It is then released.



Calculate the magnitude of the acceleration of A and the tension in the string. [6]

5. A particle is projected vertically upwards with speed 22.05 ms^{-1} .
- (a) Calculate the time that elapses before the particle returns to the point of projection and the speed of the particle at that time. [4]
- (b) Find the greatest height of the particle above the point of projection. [3]
- (c) Determine the speed and direction of motion of the particle 3 s after projection. [4]

6. The diagram shows a uniform straight rod AB , of length 3.8 m, resting horizontally in equilibrium on two smooth supports at C and D with an object of mass 2.2 kg freely suspended from point B .



The mass of the rod is 4.4 kg, $AC = 0.4$ m and $AD = 2.6$ m. Calculate the magnitudes of the reactions at C and D . [7]

7. A particle A , of mass 0.1 kg, moving with speed 10 ms^{-1} on a smooth horizontal plane collides directly with another particle B , of mass 0.6 kg moving with speed 2 ms^{-1} in the same direction.

The coefficient of restitution between the two particles is $\frac{3}{4}$.


- (a) Find the speeds of A and B after the collision. [7]

After the collision between A and B , particle B strikes a smooth vertical wall at right-angles to its direction of motion. The coefficient of restitution between B and the wall is $\frac{1}{4}$.

- (b) Show that A and B do not collide again. [2]

- (c) Calculate the magnitude of the impulse of the wall on B . [2]

8. The diagram shows a uniform plane lamina $ABCDE$, in which $ABDE$ is a rectangle and BCD is a right-angled isosceles triangle with $DB = DC$. The lengths BD and DE are 6 cm and 10 cm respectively.



- (a) Find the distance of the centre of mass of the lamina $ABCDE$ from

- (i) AB ,
(ii) AE .

[9]

- (b) The lamina $ABCDE$ is freely suspended from the point X on AE such that it hangs in equilibrium with AB vertical. Write down the distance AX . [1]

9. A body of mass 15 kg is on a rough plane inclined at an angle of 25° to the horizontal. The coefficient of friction between the body and the plane is 0.4 . The body is held at rest by means of a light string, which is parallel to a line of greatest slope of the plane. The magnitude of the tension in the string is T N. Find the greatest and the least possible values of T , giving your answers correct to two decimal places. [9]