



**GCE AS/A level**

0980/01

**MATHEMATICS – M1**  
**Mechanics**

A.M. THURSDAY, 6 June 2013

1½ hours

#### **ADDITIONAL MATERIALS**

In addition to this examination paper, you will need:

- a 12 page answer book;
- a Formula Booklet;
- a calculator.

#### **INSTRUCTIONS TO CANDIDATES**

Use black ink or black ball-point pen.

Answer **all** questions.

Take  $g$  as  $9.8 \text{ ms}^{-2}$ .


Sufficient working must be shown to demonstrate the **mathematical** method employed.

#### **INFORMATION FOR CANDIDATES**

The number of marks is given in brackets at the end of each question or part-question.

You are reminded of the necessity for good English and orderly presentation in your answers.

1. A vehicle moves along a straight horizontal road. At time  $t = 0$  s, the vehicle passes a point  $A$  and is moving with a speed of  $20 \text{ ms}^{-1}$ . It continues with this constant speed of  $20 \text{ ms}^{-1}$  for 8 s. The vehicle then slows down with uniform deceleration for 10 s so that at time  $t = 18$  s, the speed of the vehicle is  $6 \text{ ms}^{-1}$ . This speed is maintained until the vehicle reaches the point  $B$  at time  $t = 40$  s.
- (a) Sketch a velocity-time graph for the motion of the vehicle between  $A$  and  $B$ . [3]
- (b) Find the magnitude of the deceleration between  $t = 8$  and  $t = 18$ . [3]
- (c) Calculate the distance  $AB$ . [3]
2. A person of mass  $64 \text{ kg}$  is standing in a lift which is of mass  $M \text{ kg}$ . When the lift is accelerating downwards at a constant rate of  $0.425 \text{ ms}^{-2}$ , the tension in the lift cable is  $7500 \text{ N}$ .
- (a) Calculate the value of  $M$ . [3]
- (b) Find the reaction between the person and the floor of the lift. [3]
3. An object is projected vertically upwards with speed  $u \text{ ms}^{-1}$  from a point  $A$  which is  $2.8 \text{ m}$  above horizontal ground. The object reaches its greatest height of  $18.225 \text{ m}$  above  $A$  before falling to the ground.
- (a) Show that the value of  $u$  is  $18.9$ . [3]
- (b) Find the time between the object being projected and the object hitting the ground. [4]
4. The diagram shows two bodies  $A$  and  $B$ , of mass  $9 \text{ kg}$  and  $5 \text{ kg}$  respectively, connected by a light inextensible string passing over a smooth light pulley fixed at the edge of a **rough** horizontal table. The heavier body  $A$  lies on the table and the lighter body  $B$  hangs freely below the pulley.




Initially, the system is held at rest with the string taut. The system is then released.

- (a) Given that the magnitude of the acceleration of the bodies is  $1.61 \text{ ms}^{-2}$ , calculate the tension in the string and the coefficient of friction between  $A$  and the table. [8]
- (b) Given that the coefficient of friction is  $0.6$ , determine whether the bodies will move or remain at rest and evaluate the tension in the string. [3]

5. The diagram shows a uniform plank  $AB$  of mass  $12\text{ kg}$  and length  $2\text{ m}$ . The plank rests horizontally in equilibrium on two supports at  $C$  and at  $D$ , where  $AC = 0.8\text{ m}$  and  $AD = x\text{ m}$ .




- (a) The reaction of the support on the plank at  $D$  has magnitude  $84\text{ N}$ .
- (i) Determine the reaction of the support on the plank at  $C$ .
- (ii) Calculate the value of  $x$ . [7]
- (b) A rock of mass  $M\text{ kg}$  is placed at  $A$  so that the plank is on the point of tilting about  $C$ . Calculate the value of  $M$ . [3]
6. A particle  $P$ , of mass  $2\text{ kg}$ , is moving with speed  $u\text{ ms}^{-1}$  in a straight line on a smooth horizontal surface. The particle  $P$  collides directly with another particle  $Q$ , of mass  $5\text{ kg}$ , which is at rest on the surface. Immediately after the collision,  $P$  moves with speed  $2\text{ ms}^{-1}$  in a direction opposite to the original direction of motion, and the speed of  $Q$  is  $3\text{ ms}^{-1}$ .
- (a) Find the value of  $u$ . [3]
- (b) Determine the coefficient of restitution between  $P$  and  $Q$ . [3]
- (c) Calculate the magnitude of the impulse exerted by  $P$  on  $Q$ . [2]
- (d) After the collision between  $P$  and  $Q$ , particle  $Q$  strikes a vertical wall which is perpendicular to its direction of motion. The coefficient of restitution between  $Q$  and the wall is  $0.25$ . Calculate the speed with which  $Q$  rebounds from the wall. [2]
7. Four coplanar horizontal forces of magnitude  $60\text{ N}$ ,  $85\text{ N}$ ,  $75\text{ N}$  and  $40\text{ N}$  act on a particle  $P$ , of mass  $5\text{ kg}$ , in the directions shown in the diagram, where  $\tan \alpha = \frac{3}{4}$ .



- (a) Calculate the magnitude of the resultant force and determine the angle it makes with the  $85\text{ N}$  force. [9]
- (b) Deduce the magnitude of the acceleration of the particle  $P$ . [2]

**TURN OVER**

8. The diagram shows a uniform lamina in the form of a trapezium  $ABCD$  with a circular hole, of radius 2 cm, removed. The angle  $\widehat{DAB}$  is  $90^\circ$ . The dimensions, in cm, are shown in the diagram. The centre  $O$  of the circular hole is 3 cm from  $AD$  and 3 cm from  $AB$ .



- (a) Find the distances of the centre of mass of the lamina from  $AD$  and  $AB$ . [10]
- (b) When the lamina is freely suspended from a point  $Q$  on  $AD$ , it hangs in equilibrium with  $AB$  vertical. Write down the distance of  $Q$  from  $A$ . [1]