



GCE AS/A level

0977/01

MATHEMATICS – FP1
Further Pure Mathematics

A.M. WEDNESDAY, 30 January 2013

1½ hours

ADDITIONAL MATERIALS

In addition to this examination paper, you will need:

- a 12 page answer book;
- a Formula Booklet;
- a calculator.

INSTRUCTIONS TO CANDIDATES

Use black ink or black ball-point pen.

Answer **all** questions.

Sufficient working must be shown to demonstrate the **mathematical** method employed.

INFORMATION FOR CANDIDATES

The number of marks is given in brackets at the end of each question or part-question.

You are reminded of the necessity for good English and orderly presentation in your answers.

1. Differentiate $\frac{1}{2+x^2}$ from first principles. [6]

2. Consider the equations

$$\begin{aligned}x + 2y + 3z &= 4, \\2x - y + z &= 2, \\x + 7y + 8z &= k.\end{aligned}$$

Given that these equations are consistent,

(a) find the value of the constant k , [4]

(b) find the general solution of the equations. [3]

3. The complex number z and its complex conjugate \bar{z} satisfy the equation

$$iz + 2\bar{z} = \frac{4+6i}{1+i}.$$

(a) Determine z in the form $x + iy$. [6]

(b) Find the modulus and the argument of z . [2]

4. The matrix \mathbf{A} is given by

$$\mathbf{A} = \begin{bmatrix} \lambda & 1 & 1 \\ 1 & 3 & \lambda \\ 4 & 7 & 5 \end{bmatrix}.$$

(a) Find the values of λ for which \mathbf{A} is singular. [5]

(b) Given that $\lambda = 1$,

(i) determine the adjugate matrix of \mathbf{A} ,

(ii) determine the inverse matrix \mathbf{A}^{-1} . [5]

5. The roots of the cubic equation $x^3 + 4x^2 + 3x + 2 = 0$ are denoted by α, β, γ .

(a) Show that

$$\frac{1}{\beta\gamma} + \frac{1}{\gamma\alpha} + \frac{1}{\alpha\beta} = 2. [3]$$

(b) Determine the cubic equation whose roots are $\frac{1}{\beta\gamma}, \frac{1}{\gamma\alpha}, \frac{1}{\alpha\beta}$. [6]

6. Use mathematical induction to prove that

$$\sum_{r=1}^n r^3 = \frac{n^2(n+1)^2}{4}$$

for all positive integers n .

[7]

7. The function f is defined for $x > 0$ by

$$f(x) = x^{\ln x}.$$

- (a) Obtain an expression for $f'(x)$.

[4]

- (b) Find the coordinates of the stationary point on the graph of f and determine whether it is a maximum or a minimum.

[5]

8. The transformation T in the plane consists of an anticlockwise rotation through 45° about the origin followed by a reflection in the line $x + y = 0$.

- (a) Show that the 2×2 matrix representing T is

$$\frac{1}{\sqrt{2}} \begin{bmatrix} -1 & -1 \\ -1 & 1 \end{bmatrix}.$$

[3]

- (b) (i) Find the equation of the image under T of the line $y = mx$.
 (ii) Given that the line $y = mx$ is transformed into itself under T , determine the possible values of m .

[6]

9. The complex numbers z and w are represented, respectively, by points $P(x, y)$ and $Q(u, v)$ in Argand diagrams and

$$w = z(z + 1).$$

- (a) Show that

$$v = (2x + 1)y$$

and obtain an expression for u in terms of x and y .

[3]

- (b) The point P moves along the line $y = x + 1$. Find the equation of the locus of Q , giving your answer in the form $v = au^2 + bu$ where a, b are positive integers.

[7]